

wwPDB NMR Structure Validation Summary Report (i)

PDFCREATIONDATE INFOmissingINFO

PDB ID : 2XKM

Title: Consensus structure of Pf1 filamentous bacteriophage from X-ray fibre diffrac-

tion and solid-state NMR

Authors: Straus, S.K.; P Scott, W.R.; Schwieters, C.D.; Marvin, D.A.

Deposited on : 2010-07-09

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

 ${\tt PERCENTILES\ INFOmissing INFO}$



1 Ensemble composition and analysis (i)

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.



2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 665 atoms, of which 343 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called CAPSID PROTEIN G8P.

Mol	Chain	Residues	Atoms				Trace		
1	Λ	46	Total	С	Н	N	О	S	0
1	A	46	665	205	343	52	63	2	0

SEQUENCE-PLOTS INFOmissingINFO



Refinement protocol and experimental data overview (i) 3



Of the? calculated structures, 1 were deposited, based on the following criterion:?.

The authors did not provide any information on software used for structure solution, optimization or refinement.

No chemical shift data was provided. Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.



4 Model quality (i)

4.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	322	343	343	6
All	All	322	343	343	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

5 of 6 clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	$\operatorname{Distance}(\text{\AA})$
1:A:22:ILE:O	1:A:26:ILE:HG13	0.52	2.03
1:A:40:TYR:C	1:A:40:TYR:CD1	0.51	2.83
1:A:3:ILE:HG22	1:A:4:ASP:N	0.50	2.22
1:A:16:GLN:O	1:A:20:LYS:HG2	0.43	2.13
1:A:3:ILE:HG22	1:A:4:ASP:H	0.42	1.73

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perc	entiles
1	A	44/46 (96%)	42 (95%)	1 (2%)	1 (2%)	9	48
All	All	44/46 (96%)	42 (95%)	1 (2%)	1 (2%)	9	48

All 1 Ramachandran outliers are listed below.

Mol	Iol Chain		Type	
1	1 A		VAL	

4.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	32/32 (100%)	30 (94%)	2 (6%)	21 70	
All	All	32/32 (100%)	30 (94%)	2 (6%)	21 70	

All 2 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	33	LEU
1	A	40	TYR

4.3.3 RNA (i)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.



4.6 Ligand geometry (i)

There are no ligands in this entry.

4.7 Other polymers (i)

There are no such molecules in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Chemical shift validation (i)

No chemical shift data were provided

