

# Full wwPDB X-ray Structure Validation Report (i)

#### Sep 18, 2023 - 06:02 PM EDT

PDB ID	:	4XNU
Title	:	X-ray structure of Drosophila dopamine transporter in complex with nisoxetine
Authors	:	Aravind, P.; Wang, K.; Gouaux, E.
Deposited on	:	2015-01-16
Resolution	:	2.98  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.35.1
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motrie	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
R <sub>free</sub>	130704	2754 (3.00-2.96)
Clashscore	141614	3103 (3.00-2.96)
Ramachandran outliers	138981	2993 (3.00-2.96)
Sidechain outliers	138945	2996 (3.00-2.96)
RSRZ outliers	127900	2644 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain							
1	А	532	84%	16%						
2	L	214	86%	14%						
3	Н	240	8% 72% 18%	10%						



#### 4XNU

# 2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 7470 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Integral membrane protein-dopamine transporter.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	532	Total 4192	C 2814	N 647	О 713	S 18	0	2	0

• Molecule 2 is a protein called antibody fragment light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	214	Total	С	Ν	0	$\mathbf{S}$	0	0	0
	Ц	211	1621	1007	270	336	8	0	Ū	U

• Molecule 3 is a protein called antibody fragment heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Н	216	Total 1598	C 1009	N 270	0 312	S 7	0	0	0

• Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	2	Total Na 2 2	0	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total Cl 1 1	0	0

• Molecule 6 is CHOLESTEROL (three-letter code: CLR) (formula:  $C_{27}H_{46}O$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	Δ	1	Total	С	0	0	0
0	Π	T	28	27	1	0	0

• Molecule 7 is nisoxetine (three-letter code: 41U) (formula:  $C_{17}H_{21}NO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	А	1	Total 20	C 17	N 1	O 2	0	0

• Molecule 8 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	8	Total O 8 8	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- 5%
   84%
   16%

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6888
   6440
   6440
   6440

   6440
   6440
   6440
   6440

   6440
   6440
   6440
   6440

   6440
   6440
   6440
   6440

   6440
   6440
   6440
   6440

   6440
   6440
   6440
   <td
- Molecule 1: Integral membrane protein-dopamine transporter





• Molecule 3: antibody fragment heavy chain



GLY PRO



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	96.90Å 133.67Å 161.68Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Bosolution (Å)	47.10 - 2.98	Depositor
Resolution (A)	47.10 - 2.97	EDS
% Data completeness	98.6 (47.10-2.98)	Depositor
(in resolution range)	93.3 (47.10-2.97)	EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.48 (at 2.96 \text{\AA})$	Xtriage
Refinement program	PHENIX	Depositor
P. P.	0.226 , $0.251$	Depositor
$n, n_{free}$	0.229 , $0.252$	DCC
$R_{free}$ test set	1973 reflections $(4.55\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	86.8	Xtriage
Anisotropy	0.487	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.30 , $36.6$	EDS
L-test for $twinning^2$	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	7470	wwPDB-VP
Average B, all atoms $(Å^2)$	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, CLR, NA,  $41\mathrm{U}$ 

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bo	ond angles
WIOI	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.29	0/4340	0.44	2/5932~(0.0%)
2	L	0.25	0/1659	0.46	0/2257
3	Н	0.28	0/1636	0.47	0/2230
All	All	0.28	0/7635	0.45	2/10419~(0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	414	LEU	CB-CG-CD2	-7.48	98.28	111.00
1	А	582	THR	C-N-CD	5.72	140.41	128.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4192	0	4087	62	0
2	L	1621	0	1523	23	0
3	Н	1598	0	1524	35	0
4	А	2	0	0	0	0
5	А	1	0	0	1	0
6	А	28	0	46	3	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (112) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
3:H:40:SER:HB3	3:H:41:PRO:HD2	1.53	0.91
3:H:91:THR:HG22	3:H:117:VAL:H	1.46	0.78
3:H:31:SER:O	3:H:100:GLU:OE1	2.08	0.72
1:A:585:SER:H	1:A:588:GLN:HB2	1.55	0.71
3:H:89:GLU:N	3:H:89:GLU:OE1	2.24	0.71
2:L:197:ALA:HB3	2:L:206:ILE:HD11	1.72	0.70
3:H:34:MET:SD	3:H:98:ARG:HB2	2.32	0.70
1:A:598:ARG:O	8:A:707:HOH:O	2.09	0.69
1:A:252:PHE:HA	1:A:255:TRP:CD1	2.30	0.67
1:A:502[A]:ARG:NH2	8:A:707:HOH:O	2.22	0.66
1:A:364:PHE:HA	1:A:367:LEU:HB2	1.78	0.66
3:H:39:GLN:O	3:H:92:ALA:HB1	1.95	0.66
1:A:33:LYS:NZ	1:A:339:ASN:OD1	2.30	0.65
3:H:48:VAL:HG13	3:H:64:VAL:HG21	1.79	0.65
1:A:502[B]:ARG:NH2	8:A:707:HOH:O	2.22	0.64
1:A:512:GLY:HA2	3:H:102:ARG:HG3	1.79	0.64
1:A:585:SER:O	1:A:589:ARG:HG2	1.97	0.64
3:H:40:SER:CB	3:H:41:PRO:HD2	2.27	0.62
1:A:513:PHE:HB3	3:H:101:VAL:HG13	1.83	0.60
1:A:94:GLY:HA3	1:A:432:ALA:HA	1.83	0.60
1:A:43:PHE:CZ	1:A:327:VAL:HG11	2.37	0.60
2:L:134:VAL:HG22	2:L:179:THR:HG23	1.85	0.59
2:L:40:LYS:HD3	2:L:85:ALA:HB2	1.83	0.59
1:A:93:LYS:HE3	1:A:435:ASP:HB3	1.84	0.59
2:L:62:ARG:HG3	2:L:76:ILE:HG23	1.84	0.58
3:H:71:SER:HB2	3:H:80:HIS:HB2	1.85	0.58
1:A:94:GLY:N	1:A:435:ASP:OD2	2.36	0.58
1:A:52:ARG:HH11	1:A:52:ARG:HG3	1.69	0.57
1:A:84:GLU:OE1	1:A:324:GLY:N	2.34	0.57
1:A:255:TRP:CZ3	1:A:256:LYS:HE2	2.40	0.56
2:L:101:SER:O	3:H:44:ARG:HD2	2.04	0.56



Chain Non-H H(added) Clashes Symm-Clashes Mol H(model) 7 0 А 200 211 8 А 8 0 0 4 0 All All 7470 0 0 7201 112

Continued from previous page...

		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
3:H:64:VAL:HG13	3:H:68:PHE:HB2	1.87	0.56
1:A:303:ALA:HB1	1:A:309:VAL:HG21	1.87	0.56
1:A:33:LYS:NZ	1:A:344:ASP:OD2	2.38	0.55
2:L:13:THR:HG21	2:L:79:VAL:HG21	1.89	0.55
3:H:39:GLN:O	3:H:92:ALA:CB	2.56	0.54
1:A:96:ILE:HA	1:A:110:GLY:HA3	1.89	0.54
1:A:287:PRO:HG2	1:A:373:THR:HG21	1.90	0.54
3:H:87:ARG:HB3	3:H:89:GLU:OE1	2.07	0.54
2:L:150:LYS:HB2	2:L:194:THR:HB	1.90	0.53
1:A:590:PHE:O	1:A:594:THR:N	2.43	0.52
3:H:51:ILE:HD13	3:H:72:ARG:HG3	1.91	0.52
1:A:235:ILE:HG12	1:A:464:THR:HG22	1.92	0.51
1:A:123:TYR:HB3	1:A:474:LEU:HD12	1.91	0.51
2:L:88:TYR:OH	3:H:43:LYS:O	2.26	0.51
1:A:286:LEU:HD13	1:A:369:TYR:CG	2.46	0.51
1:A:582:THR:O	1:A:589:ARG:NH1	2.44	0.50
2:L:164:TRP:CE2	2:L:176:MET:HG3	2.46	0.50
1:A:433:LEU:HB3	1:A:440:ILE:HD11	1.92	0.50
1:A:358:ILE:HD13	6:A:604:CLR:H263	1.94	0.50
1:A:109:ILE:HG21	1:A:490:GLU:HG2	1.93	0.50
2:L:191:ASN:ND2	2:L:213:ASN:OD1	2.45	0.49
1:A:512:GLY:CA	3:H:102:ARG:HG3	2.42	0.49
2:L:52:THR:HG21	2:L:72:TYR:CD2	2.47	0.49
3:H:32:TYR:CB	3:H:98:ARG:HD2	2.43	0.48
3:H:52:SER:HB3	3:H:57:TYR:HB2	1.95	0.48
2:L:34:LEU:HD13	2:L:72:TYR:CD1	2.49	0.48
2:L:92:PHE:CE1	2:L:97:LEU:HD11	2.49	0.47
1:A:52:ARG:HH11	1:A:52:ARG:CG	2.28	0.47
1:A:51:TRP:C	1:A:51:TRP:CD1	2.88	0.47
1:A:296:TYR:CZ	1:A:360:GLY:HA3	2.50	0.46
2:L:97:LEU:HB2	3:H:47:TRP:CD2	2.51	0.46
1:A:75:VAL:HB	1:A:526:VAL:HG11	1.97	0.46
3:H:6:GLU:N	3:H:6:GLU:OE1	2.49	0.46
1:A:367:LEU:HA	1:A:370:MET:HE2	1.98	0.46
1:A:582:THR:OG1	1:A:589:ARG:HD3	2.17	0.45
1:A:383:THR:OG1	1:A:384:GLU:N	2.50	0.45
1:A:489:PHE:CD2	1:A:571:ILE:HG21	2.52	0.45
1:A:494:VAL:HA	1:A:498:TYR:HB3	1.98	0.45
1:A:590:PHE:O	1:A:594:THR:OG1	2.21	0.45
1:A:351:ILE:HD13	6:A:604:CLR:H71	1.99	0.44
3:H:83:MET:HE2	3:H:86:LEU:HD21	1.99	0.44

Continued from previous page...



A 4 1		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:A:389:VAL:HB	1:A:414:LEU:HD21	1.98	0.44
2:L:92:PHE:CD1	3:H:102:ARG:HD3	2.52	0.44
1:A:292:GLY:HA3	1:A:364:PHE:O	2.17	0.44
1:A:582:THR:OG1	1:A:589:ARG:CD	2.66	0.44
2:L:92:PHE:CE2	2:L:97:LEU:HD21	2.53	0.43
1:A:69:TYR:HA	1:A:313:ALA:HB1	2.00	0.43
2:L:161:LEU:HD21	3:H:175:VAL:HB	2.01	0.43
3:H:12:VAL:O	3:H:117:VAL:HA	2.18	0.43
1:A:502[B]:ARG:HH21	3:H:56:ARG:HH11	1.66	0.43
1:A:370:MET:HG2	1:A:374:LEU:HD22	2.01	0.43
1:A:51:TRP:HD1	1:A:52:ARG:N	2.16	0.42
2:L:52:THR:HG21	2:L:72:TYR:CE2	2.54	0.42
3:H:29:PHE:O	3:H:72:ARG:NH2	2.52	0.42
1:A:501:ASN:OD1	1:A:501:ASN:N	2.53	0.42
2:L:171:ASP:N	2:L:171:ASP:OD1	2.53	0.42
1:A:410:PHE:CZ	1:A:414:LEU:HD11	2.55	0.42
3:H:44:ARG:NH1	3:H:44:ARG:HG2	2.34	0.42
1:A:140:THR:HG22	1:A:142:SER:H	1.85	0.42
1:A:572:PRO:O	1:A:576:ILE:HG12	2.20	0.42
2:L:60:PRO:HG2	2:L:63:PHE:CD2	2.55	0.42
3:H:19:LYS:HG3	3:H:82:GLU:HG3	2.01	0.41
3:H:217:GLU:HA	3:H:218:PRO:HD3	1.93	0.41
2:L:92:PHE:CE1	3:H:102:ARG:HD3	2.56	0.41
3:H:5:VAL:HG23	3:H:23:ALA:HB3	2.02	0.41
1:A:56:LEU:HD23	1:A:56:LEU:HA	1.87	0.41
1:A:595:THR:HA	1:A:596:PRO:HD3	1.89	0.41
1:A:134:PHE:HB3	1:A:411:PHE:CE2	2.56	0.41
7:A:605:41U:H8	7:A:605:41U:H7	1.86	0.41
2:L:116:VAL:HG21	2:L:206:ILE:HD13	2.03	0.41
1:A:95:ALA:HA	1:A:329:LEU:HD23	2.02	0.41
1:A:115:LEU:HB3	8:A:706:HOH:O	2.20	0.41
1:A:145:TRP:HB2	1:A:212:ALA:HA	2.03	0.41
3:H:19:LYS:NZ	3:H:82:GLU:HB2	2.34	0.41
6:A:604:CLR:H232	6:A:604:CLR:H211	1.78	0.41
3:H:161:ASN:OD1	3:H:200:THR:N	2.33	0.41
1:A:316:GLN:NE2	5:A:603:CL:CL	2.84	0.41
1:A:486:ALA:O	1:A:490:GLU:HG3	2.20	0.41
2:L:36:TRP:HB2	2:L:49:ILE:HB	2.01	0.40
1:A:420:ASP:HA	1:A:423:PHE:CD2	2.57	0.40
1:A:251:TYR:O	1:A:255:TRP:HD1	2.04	0.40

Continued from previous page...

There are no symmetry-related clashes.



### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	532/532~(100%)	516~(97%)	16 (3%)	0	100 100
2	L	212/214~(99%)	205~(97%)	7 (3%)	0	100 100
3	Н	212/240~(88%)	203~(96%)	9~(4%)	0	100 100
All	All	956/986~(97%)	924 (97%)	32 (3%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	425/437~(97%)	421 (99%)	4 (1%)	78 91
2	L	182/187~(97%)	181 (100%)	1 (0%)	88 95
3	Н	171/205~(83%)	169 (99%)	2 (1%)	71 89
All	All	778/829~(94%)	771 (99%)	7 (1%)	78 91

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	25	ASP
1	А	51	TRP
1	А	581	SER
1	А	582	THR
2	L	92	PHE



Continued from previous page...

Mol	Chain	Res	Type
3	Η	44	ARG
3	Н	215	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	158	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	n Res	bein Dec Link Bond lengths		B	ond ang	gles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	41U	А	605	-	21,21,21	0.57	0	24,26,26	1.34	5 (20%)
6	CLR	А	604	-	31,31,31	0.65	0	48,48,48	1.15	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings		
7	41U	А	605	-	-	12/14/14/14	0/2/2/2		
6	CLR	А	604	-	-	0/10/68/68	0/4/4/4		

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
7	А	605	41U	O-C7-C14	3.57	110.88	105.18
6	А	604	CLR	C7-C8-C9	3.09	113.46	109.71
6	А	604	CLR	C4-C5-C10	2.98	120.38	116.42
7	А	605	41U	C16-N-C15	2.79	120.02	111.95
7	А	605	41U	O1-C5-C	2.45	118.83	115.41
6	А	604	CLR	C14-C8-C9	-2.36	105.93	109.09
7	А	605	41U	C14-C7-C8	-2.09	108.84	112.62
7	А	605	41U	O1-C5-C4	-2.06	120.83	124.37

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
7	А	605	41U	C15-C14-C7-O
7	А	605	41U	C15-C14-C7-C8
7	А	605	41U	C-C5-O1-C6
7	А	605	41U	C4-C5-O1-C6
7	А	605	41U	C14-C15-N-C16
7	А	605	41U	O-C7-C8-C9
7	А	605	41U	С8-С7-О-С
7	А	605	41U	O-C7-C8-C13
7	А	605	41U	C14-C7-C8-C13
7	А	605	41U	C14-C7-C8-C9
7	А	605	41U	C7-C14-C15-N
7	А	605	41U	C1-C-O-C7

All (12) torsion outliers are listed below:

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	А	605	41U	1	0
6	А	604	CLR	3	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	532/532~(100%)	0.46	28 (5%) 26 15	66, 85, 109, 145	0
2	L	214/214~(100%)	0.66	24 (11%) 5 3	57, 77, 110, 146	0
3	Н	216/240~(90%)	0.45	18 (8%) 11 5	61, 85, 114, 134	0
All	All	962/986~(97%)	0.50	70 (7%) 15 7	57, 84, 111, 146	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	123	SER	7.1
2	L	126	LEU	5.4
2	L	127	THR	4.6
2	L	136	PHE	4.6
2	L	118	ILE	4.6
3	Н	129	PRO	4.6
1	А	437	PHE	4.5
1	А	439	LYS	4.5
2	L	119	PHE	4.5
1	А	291	LEU	4.5
2	L	134	VAL	4.3
1	А	290[A]	PHE	4.2
2	L	122	SER	4.2
3	Н	130	LEU	4.0
3	Н	144	LEU	3.9
3	Н	199	ILE	3.9
2	L	135	CYS	3.7
1	А	540	ILE	3.6
2	L	57	SER	3.6
2	L	213	ASN	3.5
2	L	130	GLY	3.5
3	Н	127	VAL	3.5
3	Н	79	LEU	3.5



Mol	Chain	Res	Type	RSRZ
3	Н	128	TYR	3.4
1	А	375	GLY	3.3
1	А	586	LEU	3.2
1	А	440	ILE	3.2
3	Н	131	ALA	3.1
3	Н	147	LEU	3.0
2	L	124	GLU	3.0
3	Н	138	THR	3.0
1	А	492	ILE	3.0
1	А	237	TRP	2.9
1	А	590	PHE	2.9
1	А	111	TYR	2.9
1	Α	105	LEU	2.9
1	A	438	PRO	2.8
2	L	137	LEU	2.8
3	Н	214	LYS	2.7
1	А	229	ILE	2.7
1	А	497	ILE	2.7
1	А	496	TRP	2.6
3	Н	100	GLU	2.6
1	А	481	TYR	2.6
1	А	346	LEU	2.5
2	L	151	ILE	2.5
1	А	461	ALA	2.5
1	А	580	LEU	2.5
2	L	95	TYR	2.4
3	Н	108	TYR	2.4
2	L	210	PHE	2.4
2	L	117	SER	2.4
3	Н	198	SER	2.4
1	А	579	LEU	2.3
2	L	128	SER	2.3
1	А	555	TRP	2.3
3	Н	142	VAL	2.3
1	A	433	LEU	2.3
1	А	372	HIS	2.3
1	А	576	ILE	2.3
3	Н	211	LYS	2.3
2	L	176	MET	2.3
2	L	212	ARG	2.2
3	Н	215	LYS	2.2
2	L	161	LEU	2.2

Continued from previous page...



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	94	GLY	2.1
3	Н	189	VAL	2.1
1	А	353	SER	2.0
1	А	96	ILE	2.0
2	L	133	VAL	2.0

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
6	CLR	А	604	28/28	0.88	0.73	84,97,109,113	0
4	NA	А	602	1/1	0.90	0.23	70,70,70,70	0
7	41U	А	605	20/20	0.93	0.33	88,94,111,113	0
4	NA	А	601	1/1	0.94	0.33	84,84,84,84	0
5	CL	А	603	1/1	0.96	0.15	74,74,74,74	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









## 6.5 Other polymers (i)

There are no such residues in this entry.

