



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2023 – 02:47 AM EDT

PDB ID : 6XNR  
Title : Crystal structure of Rhagium Mordax antifreeze protein  
Authors : Ye, Q.; Eves, R.; Campbell, R.L.; Davies, P.L.  
Deposited on : 2020-07-04  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	145	
1	BBB	145	
1	CCC	145	
1	DDD	145	
1	EEE	145	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
2	EDO	BBB	201	-	-	X	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4922 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antifreeze protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	139	Total 892	C 519	N 155	O 215	S 3	0	1	0
1	BBB	138	Total 880	C 512	N 154	O 211	S 3	0	0	0
1	CCC	138	Total 880	C 512	N 154	O 211	S 3	0	0	0
1	DDD	136	Total 864	C 501	N 152	O 209	S 2	0	0	0
1	EEE	138	Total 880	C 512	N 154	O 211	S 3	0	0	0

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	BBB	1	Total 4	C 2	O 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	CCC	1	Total C O 4 2 2	0	0
2	DDD	1	Total C O 4 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	117	Total O 117 117	0	0
3	BBB	116	Total O 116 116	0	0
3	CCC	123	Total O 123 123	0	0
3	DDD	89	Total O 89 89	0	0
3	EEE	69	Total O 69 69	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Antifreeze protein





- Molecule 1: Antifreeze protein




- Molecule 1: Antifreeze protein



- Molecule 1: Antifreeze protein



- Molecule 1: Antifreeze protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.46Å 91.14Å 117.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.41 – 2.05 49.36 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.41-2.05) 99.7 (49.36-2.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 2.05Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.213 , 0.251 0.217 , 0.252	Depositor DCC
$R_{free}$ test set	2230 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.2	Xtrriage
Anisotropy	0.757	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 42.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4922	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	AAA	0.70	0/899	0.87	0/1237
1	BBB	0.73	0/884	0.85	0/1217
1	CCC	0.68	0/884	0.84	0/1217
1	DDD	0.68	0/868	0.83	0/1196
1	EEE	0.67	0/884	0.85	0/1217
All	All	0.69	0/4419	0.85	0/6084

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	892	0	866	10	0
1	BBB	880	0	855	17	0
1	CCC	880	0	855	10	0
1	DDD	864	0	832	9	0
1	EEE	880	0	855	9	0
2	BBB	4	0	6	4	0
2	CCC	4	0	6	2	0
2	DDD	4	0	6	3	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	AAA	117	0	0	4	0
3	BBB	116	0	0	4	0
3	CCC	123	0	0	1	0
3	DDD	89	0	0	2	0
3	EEE	69	0	0	0	0
All	All	4922	0	4281	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:9:VAL:O	1:BBB:28:PRO:HA	1.69	0.91
1:BBB:96:ILE:HG23	2:BBB:201:EDO:H12	1.66	0.77
1:DDD:100:THR:HG21	1:EEE:78:ILE:HD11	1.70	0.72
1:BBB:78:ILE:HD11	1:CCC:100:THR:HG21	1.68	0.72
1:BBB:74:THR:OG1	2:CCC:301:EDO:H12	1.94	0.66
1:BBB:1:MET:HE2	3:BBB:416:HOH:O	1.95	0.64
1:BBB:74:THR:OG1	2:BBB:201:EDO:H21	1.98	0.64
1:DDD:125:ILE:HD11	1:EEE:56:THR:CG2	2.27	0.63
1:BBB:74:THR:OG1	2:CCC:301:EDO:C1	2.47	0.63
1:BBB:78:ILE:CD1	1:CCC:100:THR:HG21	2.30	0.60
1:BBB:138:LEU:C	3:BBB:347:HOH:O	2.39	0.60
1:AAA:1:MET:HE2	3:AAA:317:HOH:O	2.04	0.58
1:BBB:96:ILE:CG2	2:BBB:201:EDO:H12	2.33	0.57
1:BBB:122:ALA:HB3	1:BBB:134:VAL:HG22	1.87	0.56
1:BBB:22:HIS:HD2	3:BBB:400:HOH:O	1.90	0.55
1:DDD:100:THR:HG21	1:EEE:78:ILE:CD1	2.36	0.54
1:AAA:101:THR:HA	1:AAA:126:ALA:O	2.08	0.53
1:DDD:101:THR:HA	1:DDD:126:ALA:O	2.09	0.52
1:AAA:122:ALA:HB3	1:AAA:134:VAL:HG22	1.91	0.52
1:CCC:10:ASP:OD1	1:CCC:12:SER:HB3	2.09	0.52
1:BBB:101:THR:HA	1:BBB:126:ALA:O	2.09	0.52
1:CCC:101:THR:HA	1:CCC:126:ALA:O	2.10	0.52
1:AAA:112:LYS:HE2	3:AAA:215:HOH:O	2.10	0.52
1:DDD:91:ALA:O	1:DDD:92:ALA:HB3	2.09	0.51
1:EEE:101:THR:HA	1:EEE:126:ALA:O	2.09	0.51
1:DDD:76:ASN:HB3	1:DDD:78:ILE:HD11	1.93	0.51
1:BBB:96:ILE:HG23	2:BBB:201:EDO:C1	2.40	0.51
1:AAA:112:LYS:CE	3:AAA:215:HOH:O	2.59	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:DDD:201:EDO:H21	1:EEE:74:THR:CB	2.41	0.51
1:BBB:91:ALA:O	1:BBB:92:ALA:HB3	2.11	0.49
3:BBB:323:HOH:O	1:CCC:123:THR:HG21	2.12	0.49
2:DDD:201:EDO:H21	1:EEE:74:THR:HB	1.95	0.49
1:EEE:122:ALA:HB3	1:EEE:134:VAL:HG22	1.95	0.49
1:CCC:115:SER:HA	1:CCC:135:THR:O	2.12	0.48
1:CCC:91:ALA:O	1:CCC:92:ALA:HB3	2.13	0.48
1:EEE:92:ALA:HB3	1:EEE:138:LEU:HB3	1.97	0.47
1:CCC:76:ASN:HB3	3:CCC:403:HOH:O	2.13	0.47
1:AAA:34:GLY:HA2	1:AAA:55:SER:O	2.16	0.45
1:DDD:112:LYS:HE3	3:DDD:356:HOH:O	2.16	0.44
1:AAA:3[B]:SER:H	1:AAA:18:GLN:NE2	2.15	0.44
1:AAA:3[A]:SER:H	1:AAA:18:GLN:NE2	2.15	0.44
1:DDD:78:ILE:N	1:DDD:78:ILE:HD12	2.33	0.44
1:BBB:3:SER:H	1:BBB:18:GLN:NE2	2.16	0.43
1:DDD:9:VAL:HA	3:DDD:337:HOH:O	2.19	0.42
1:BBB:122:ALA:HB3	1:BBB:134:VAL:CG2	2.49	0.41
1:AAA:22:HIS:HE1	3:AAA:286:HOH:O	2.03	0.41
1:CCC:114:VAL:O	1:CCC:134:VAL:HA	2.19	0.41
2:DDD:201:EDO:C2	1:EEE:74:THR:HB	2.51	0.41
1:AAA:122:ALA:HB3	1:AAA:134:VAL:CG2	2.51	0.40
1:CCC:98:SER:O	1:CCC:123:THR:HA	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	138/145 (95%)	137 (99%)	1 (1%)	0	100	100
1	BBB	136/145 (94%)	131 (96%)	5 (4%)	0	100	100
1	CCC	136/145 (94%)	132 (97%)	4 (3%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	DDD	134/145 (92%)	132 (98%)	2 (2%)	0	100	100
1	EEE	136/145 (94%)	135 (99%)	1 (1%)	0	100	100
All	All	680/725 (94%)	667 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	94/99 (95%)	94 (100%)	0	100	100
1	BBB	92/99 (93%)	92 (100%)	0	100	100
1	CCC	92/99 (93%)	92 (100%)	0	100	100
1	DDD	90/99 (91%)	90 (100%)	0	100	100
1	EEE	92/99 (93%)	92 (100%)	0	100	100
All	All	460/495 (93%)	460 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	BBB	201	-	3,3,3	0.12	0	2,2,2	0.37	0
2	EDO	CCC	301	-	3,3,3	0.23	0	2,2,2	0.79	0
2	EDO	DDD	201	-	3,3,3	0.15	0	2,2,2	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	BBB	201	-	-	0/1/1/1	-
2	EDO	CCC	301	-	-	0/1/1/1	-
2	EDO	DDD	201	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	BBB	201	EDO	4	0
2	CCC	301	EDO	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	DDD	201	EDO	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	139/145 (95%)	0.09	2 (1%) 75 78	22, 28, 44, 60	0
1	BBB	138/145 (95%)	-0.02	3 (2%) 62 66	23, 30, 43, 56	0
1	CCC	138/145 (95%)	0.04	5 (3%) 42 46	24, 29, 42, 53	0
1	DDD	136/145 (93%)	0.02	6 (4%) 34 37	30, 39, 53, 57	0
1	EEE	138/145 (95%)	0.35	9 (6%) 18 20	32, 42, 61, 67	0
All	All	689/725 (95%)	0.10	25 (3%) 42 46	22, 34, 52, 67	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	DDD	11	ALA	4.3
1	DDD	9	VAL	4.2
1	CCC	138	LEU	4.2
1	DDD	10	ASP	4.2
1	BBB	9	VAL	4.0
1	CCC	12	SER	3.9
1	AAA	139	GLU	3.5
1	EEE	138	LEU	3.5
1	DDD	104	SER	3.4
1	CCC	92	ALA	3.3
1	EEE	79	GLY	3.1
1	AAA	92	ALA	3.1
1	EEE	1	MET	3.0
1	EEE	78	ILE	3.0
1	DDD	12	SER	2.8
1	BBB	10	ASP	2.8
1	CCC	9	VAL	2.7
1	EEE	108	PRO	2.7
1	EEE	127	ILE	2.6
1	CCC	11	ALA	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	EEE	103	ALA	2.5
1	EEE	107	GLU	2.5
1	BBB	11	ALA	2.4
1	EEE	109	THR	2.3
1	DDD	103	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EDO	BBB	201	4/4	0.83	0.22	48,49,50,54	0
2	EDO	CCC	301	4/4	0.85	0.23	34,39,43,46	0
2	EDO	DDD	201	4/4	0.85	0.32	51,51,54,54	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.