



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 02:21 am BST

PDB ID : 1XP6
Title : HUMAN ESTROGEN RECEPTOR ALPHA LIGAND-BINDING DOMAIN
IN COMPLEX WITH COMPOUND 16
Authors : Fitzgerald, P.M.D.; Sharma, N.
Deposited on : 2004-10-08
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	248	71% 23% . . .

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Estrogen receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	1951	1251	332	349	19	0	0	0

- Molecule 2 is (2S,3R)-2-(4-{2-[(3S,4S)-3,4-DIMETHYLPYRROLIDIN-1-YL]ETHOXY}PHENYL)-3-(4-HYDROXYPHENYL)-2,3-DIHYDRO-1,4-BENZOXATHIIN-6-OL (three-letter code: AIU) (formula: C₂₈H₃₁NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	34	28	1	4	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	157	Total	O	0	0
			157	157		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Estrogen receptor



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	58.57Å 58.57Å 276.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 1.70 10.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	95.3 (10.00-1.70) 89.7 (10.00-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.70Å)	Xtrriage
Refinement program	SHELXL 97-1	Depositor
R, R_{free}	0.177 , 0.267 0.230 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	29.7	Xtrriage
Anisotropy	0.150	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 83.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2142	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AIU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1989	1.02	5/2690 (0.2%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	459	TYR	CB-CG-CD2	6.41	124.84	121.00
1	A	548	ARG	NE-CZ-NH2	6.04	123.32	120.30
1	A	548	ARG	NE-CZ-NH1	-5.99	117.30	120.30
1	A	436	ARG	CD-NE-CZ	5.39	131.14	123.60
1	A	462	LEU	C-N-CA	5.23	134.77	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1951	0	1997	61	0
2	A	34	0	29	3	0
3	A	157	0	0	5	0
All	All	2142	0	2026	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 15.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:411:ASP:H	1:A:414:GLN:HE21	1.13	0.94
1:A:311:THR:HG23	1:A:314:GLN:H	1.31	0.92
1:A:311:THR:HG21	3:A:1130:HOH:O	1.85	0.75
1:A:403:LEU:HD13	1:A:409:LEU:HD13	1.67	0.75
1:A:308:LEU:HD23	1:A:477:ARG:HD2	1.73	0.70
1:A:307:ALA:HB2	1:A:365:PRO:HB2	1.73	0.69
1:A:311:THR:HG23	1:A:314:GLN:N	2.07	0.69
1:A:474:HIS:HA	1:A:477:ARG:NH1	2.07	0.69
1:A:486:LEU:O	1:A:490:MET:HG3	1.93	0.68
1:A:437:MET:HB2	3:A:1092:HOH:O	1.94	0.68
1:A:308:LEU:H	1:A:308:LEU:HD12	1.62	0.64
1:A:315:MET:SD	1:A:365:PRO:HG2	2.39	0.62
1:A:381:CYS:HB3	3:A:1096:HOH:O	1.99	0.62
1:A:331:TYR:HB3	1:A:345:LEU:HD21	1.82	0.62
1:A:411:ASP:H	1:A:414:GLN:NE2	1.92	0.61
1:A:410:LEU:HA	1:A:414:GLN:NE2	2.16	0.61
1:A:529:LYS:HG2	1:A:530:CYS:N	2.16	0.60
1:A:308:LEU:HD23	1:A:477:ARG:HB3	1.85	0.59
1:A:308:LEU:CD2	1:A:477:ARG:HD2	2.33	0.58
1:A:307:ALA:HB2	1:A:365:PRO:CB	2.35	0.56
1:A:498:GLN:HA	1:A:501:HIS:CE1	2.40	0.56
1:A:354:LEU:HD22	2:A:600:AIU:H332	1.86	0.56
1:A:435:PHE:CE2	1:A:510:ILE:HG21	2.41	0.55
1:A:530:CYS:O	1:A:531:LYS:HB2	2.07	0.54
1:A:547:HIS:HD2	1:A:549:LEU:H	1.55	0.53
1:A:458:VAL:HG13	1:A:459:TYR:CD1	2.45	0.52
1:A:308:LEU:HD21	1:A:477:ARG:HH11	1.75	0.50
1:A:533:VAL:O	1:A:534:VAL:O	2.30	0.50
1:A:374:ASP:OD2	1:A:471:GLU:OE1	2.30	0.50
1:A:539:LEU:HG	1:A:543:MET:CE	2.42	0.50
1:A:379:LEU:HD12	1:A:544:LEU:HD11	1.93	0.49
1:A:310:LEU:HG	1:A:314:GLN:HB3	1.95	0.48
1:A:435:PHE:HE2	1:A:510:ILE:HG21	1.77	0.48
1:A:458:VAL:HG13	1:A:459:TYR:HD1	1.77	0.47
1:A:547:HIS:CD2	1:A:549:LEU:H	2.31	0.47
1:A:307:ALA:HB1	1:A:315:MET:HE1	1.96	0.47
1:A:456:SER:HA	1:A:515:ARG:NH2	2.30	0.47
1:A:326:ILE:HD11	3:A:1098:HOH:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:366:GLY:O	1:A:474:HIS:HE1	1.98	0.46
1:A:389:ILE:O	1:A:392:VAL:HG22	2.15	0.46
1:A:530:CYS:HB2	2:A:600:AIU:H262	1.98	0.46
1:A:307:ALA:HA	1:A:310:LEU:HB2	1.98	0.46
1:A:481:LYS:HB2	1:A:481:LYS:HE3	1.55	0.46
1:A:311:THR:OG1	1:A:312:ALA:N	2.49	0.45
1:A:389:ILE:HA	1:A:392:VAL:HG22	1.98	0.45
1:A:435:PHE:CE2	1:A:510:ILE:HD13	2.51	0.45
1:A:458:VAL:O	1:A:458:VAL:HG22	2.15	0.45
1:A:308:LEU:CD2	1:A:477:ARG:HH11	2.30	0.45
1:A:307:ALA:O	1:A:481:LYS:HG2	2.16	0.45
1:A:396:MET:HB2	1:A:396:MET:HE2	1.32	0.45
1:A:435:PHE:HE2	1:A:510:ILE:HD13	1.82	0.44
1:A:523:GLU:HG2	3:A:1111:HOH:O	2.17	0.44
1:A:311:THR:HG22	1:A:314:GLN:CD	2.38	0.44
1:A:535:PRO:HD2	1:A:538:ASP:OD2	2.19	0.43
1:A:307:ALA:HB1	1:A:315:MET:CE	2.50	0.42
1:A:535:PRO:HG2	1:A:538:ASP:OD2	2.21	0.41
1:A:531:LYS:HB3	2:A:600:AIU:H291	2.02	0.41
1:A:495:LEU:HB3	1:A:499:GLN:HB2	2.03	0.41
1:A:358:ILE:HD13	1:A:544:LEU:HG	2.03	0.41
1:A:328:TYR:CE2	1:A:406:PRO:HB2	2.56	0.40
1:A:372:LEU:HD21	1:A:545:ASP:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	243/248 (98%)	230 (95%)	9 (4%)	4 (2%)	9 1

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	530	CYS
1	A	534	VAL
1	A	462	LEU
1	A	531	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	219/223 (98%)	202 (92%)	17 (8%)	12 3

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	308	LEU
1	A	311	THR
1	A	313	ASP
1	A	338	SER
1	A	358	ILE
1	A	394	ARG
1	A	396	MET
1	A	437	MET
1	A	462	LEU
1	A	463	SER
1	A	481	LYS
1	A	492	LYS
1	A	512	SER
1	A	527	SER
1	A	529	LYS
1	A	531	LYS
1	A	532	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	414	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	474	HIS
1	A	547	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AIU	A	600	-	36,38,38	2.89	1 (2%)	47,54,54	1.70	11 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AIU	A	600	-	-	3/14/38/38	0/4/5/5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	AIU	C10-S11	-16.30	1.53	1.76

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	600	AIU	C14-C13-C12	-4.01	117.16	121.20
2	A	600	AIU	C18-C12-C13	3.79	123.03	118.29
2	A	600	AIU	C10-S11-C1	3.21	109.10	100.03
2	A	600	AIU	C9-C10-C4	-3.18	116.68	119.36
2	A	600	AIU	C12-C1-S11	2.94	117.82	106.41
2	A	600	AIU	C24-C19-C2	2.87	125.97	120.64
2	A	600	AIU	C27-N28-C29	-2.80	105.74	113.19
2	A	600	AIU	O25-C26-C27	2.66	114.23	107.68
2	A	600	AIU	C5-C6-C7	-2.54	117.08	119.88
2	A	600	AIU	C7-C9-C10	2.27	121.82	118.63
2	A	600	AIU	C18-C12-C1	-2.03	114.15	120.81

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	600	AIU	C24-C19-C2-O3
2	A	600	AIU	C20-C19-C2-O3
2	A	600	AIU	S11-C1-C12-C13


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	AIU	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

Unable to reproduce the depositor's R factor - this section is therefore empty.