



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 10, 2023 – 12:11 PM JST

PDB ID : 7XSG  
Title : Crystal structure of ClAgl29B  
Authors : Shishiuchi, R.; Kang, H.; Tagami, T.; Okuyama, M.  
Deposited on : 2022-05-14  
Resolution : 1.61 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.3  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.3


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	587	 16% 87% 5% 7%
1	B	587	 5% 89% 7% 7%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 10033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-L-fucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	545	4481	2882	727	855	17	0	8	0
1	B	545	4504	2893	731	863	17	0	12	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP K1KV82
A	1	GLY	-	expression tag	UNP K1KV82
A	2	SER	-	expression tag	UNP K1KV82
A	3	SER	-	expression tag	UNP K1KV82
A	4	HIS	-	expression tag	UNP K1KV82
A	5	HIS	-	expression tag	UNP K1KV82
A	6	HIS	-	expression tag	UNP K1KV82
A	7	HIS	-	expression tag	UNP K1KV82
A	8	HIS	-	expression tag	UNP K1KV82
A	9	HIS	-	expression tag	UNP K1KV82
A	10	SER	-	expression tag	UNP K1KV82
A	11	SER	-	expression tag	UNP K1KV82
A	12	GLY	-	expression tag	UNP K1KV82
A	13	LEU	-	expression tag	UNP K1KV82
A	14	VAL	-	expression tag	UNP K1KV82
A	15	PRO	-	expression tag	UNP K1KV82
A	16	ARG	-	expression tag	UNP K1KV82
A	17	GLY	-	expression tag	UNP K1KV82
A	18	SER	-	expression tag	UNP K1KV82
A	19	HIS	-	expression tag	UNP K1KV82
A	20	MET	-	expression tag	UNP K1KV82
B	0	MET	-	initiating methionine	UNP K1KV82
B	1	GLY	-	expression tag	UNP K1KV82
B	2	SER	-	expression tag	UNP K1KV82
B	3	SER	-	expression tag	UNP K1KV82

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	4	HIS	-	expression tag	UNP K1KV82
B	5	HIS	-	expression tag	UNP K1KV82
B	6	HIS	-	expression tag	UNP K1KV82
B	7	HIS	-	expression tag	UNP K1KV82
B	8	HIS	-	expression tag	UNP K1KV82
B	9	HIS	-	expression tag	UNP K1KV82
B	10	SER	-	expression tag	UNP K1KV82
B	11	SER	-	expression tag	UNP K1KV82
B	12	GLY	-	expression tag	UNP K1KV82
B	13	LEU	-	expression tag	UNP K1KV82
B	14	VAL	-	expression tag	UNP K1KV82
B	15	PRO	-	expression tag	UNP K1KV82
B	16	ARG	-	expression tag	UNP K1KV82
B	17	GLY	-	expression tag	UNP K1KV82
B	18	SER	-	expression tag	UNP K1KV82
B	19	HIS	-	expression tag	UNP K1KV82
B	20	MET	-	expression tag	UNP K1KV82

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0
2	B	1	Total Na 1 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 7 4 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	393	Total O 393 393	0	0
6	B	615	Total O 615 615	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-L-fucosidase



- Molecule 1: Alpha-L-fucosidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.13Å 123.41Å 166.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.42 – 1.61 45.38 – 1.61	Depositor EDS
% Data completeness (in resolution range)	99.5 (45.42-1.61) 99.5 (45.38-1.61)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.50 (at 1.61Å)	Xtrriage
Refinement program	REFMAC 5.8.0349	Depositor
R, $R_{free}$	0.172 , 0.188 0.185 , 0.200	Depositor DCC
$R_{free}$ test set	9719 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.5	Xtrriage
Anisotropy	0.478	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 47.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10033	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, GOL, CIT, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/4604	0.71	1/6237 (0.0%)
1	B	0.45	0/4627	0.71	1/6268 (0.0%)
All	All	0.43	0/9231	0.71	2/12505 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	419	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	A	149	TYR	CB-CG-CD1	5.16	124.10	121.00

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	162[A]	ARG	Sidechain
1	A	44	ARG	Sidechain
1	B	162[A]	ARG	Sidechain
1	B	44	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4481	0	4316	16	0
1	B	4504	0	4330	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	12	0	16	1	0
3	B	6	0	8	0	0
4	A	13	0	5	0	0
5	B	7	0	10	0	0
6	A	393	0	0	2	0
6	B	615	0	0	3	0
All	All	10033	0	8685	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 27 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:ASN:HD21	1:A:265:ASN:H	1.19	0.86
1:A:541:LYS:HA	1:A:549:LEU:HD13	1.83	0.61
1:A:547:ILE:HD12	1:A:548:ASN:O	2.02	0.59
1:A:73:ARG:HG3	6:A:701:HOH:O	2.05	0.56
1:B:55:TYR:HE1	1:B:63:GLU:OE1	1.89	0.56

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	549/587 (94%)	534 (97%)	15 (3%)	0	100	100
1	B	553/587 (94%)	538 (97%)	15 (3%)	0	100	100
All	All	1102/1174 (94%)	1072 (97%)	30 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	488/517 (94%)	482 (99%)	6 (1%)	71	54
1	B	492/517 (95%)	488 (99%)	4 (1%)	81	70
All	All	980/1034 (95%)	970 (99%)	10 (1%)	76	61

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	112	PHE
1	B	196	THR
1	B	325	TRP
1	A	547	ILE
1	A	559	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	275	GLN
1	B	491	HIS
1	B	586	ASN
1	B	518	HIS
1	B	186	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	A	602	-	5,5,5	0.09	0	5,5,5	0.17	0
4	CIT	A	603	-	12,12,12	1.36	1 (8%)	17,17,17	1.41	2 (11%)
3	GOL	B	602	-	5,5,5	0.17	0	5,5,5	0.44	0
3	GOL	A	604	-	5,5,5	0.02	0	5,5,5	0.25	0
5	PEG	B	603	-	6,6,6	0.33	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	602	-	-	0/4/4/4	-
4	CIT	A	603	-	-	0/16/16/16	-
3	GOL	B	602	-	-	0/4/4/4	-
3	GOL	A	604	-	-	4/4/4/4	-
5	PEG	B	603	-	-	1/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	603	CIT	C3-C6	2.41	1.55	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	603	CIT	O5-C6-C3	-3.77	116.91	122.25
4	A	603	CIT	O6-C6-C3	2.62	117.59	113.05

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	604	GOL	O1-C1-C2-C3
3	A	604	GOL	O1-C1-C2-O2
3	A	604	GOL	O2-C2-C3-O3
5	B	603	PEG	O2-C3-C4-O4
3	A	604	GOL	C1-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	604	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	545/587 (92%)	0.82	96 (17%) <b>1</b> <b>1</b>	10, 20, 51, 100	0
1	B	545/587 (92%)	0.13	29 (5%) <b>26</b> <b>24</b>	8, 12, 34, 67	0
All	All	1090/1174 (92%)	0.47	125 (11%) <b>4</b> <b>4</b>	8, 15, 47, 100	0

The worst 5 of 125 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	555	SER	8.5
1	A	549	LEU	7.6
1	A	522	TRP	7.5
1	A	573	LEU	7.2
1	A	567	PHE	7.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	PEG	B	603	7/7	0.62	0.21	36,41,47,48	0
3	GOL	A	604	6/6	0.80	0.22	37,43,46,48	0
4	CIT	A	603	13/13	0.81	0.20	22,25,28,29	0
3	GOL	B	602	6/6	0.88	0.13	19,21,25,27	0
2	NA	A	601	1/1	0.91	0.28	33,33,33,33	0
3	GOL	A	602	6/6	0.96	0.07	19,21,22,23	0
2	NA	B	601	1/1	0.99	0.08	19,19,19,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.