Full wwPDB X-ray Structure Validation Report (i)

Jun 17, 2024 – 05:23 AM EDT

PDB ID	:	2XXI
Title	:	Crystal structure of 1-((4-(3-(trifluoromethyl)-6,7-dihydropyrano(4,3- c(pyrazol-1(4H)-yl)phenyl)methyl)-2-pyrrolidinone in complex with the ligand binding domain of the Rat GluA2 receptor and glutamate at 1.6A resolution.
Authors	:	Ward, S.E.; Harries, M.; Aldegheri, L.; Austin, N.E.; Ballantine, S.; Ballini, E.; Bradley, D.M.; Bax, B.D.; Clarke, B.P.; Harris, A.J.; Harrison, S.A.; Melarange, R.A.; Mookherjee, C.; Mosley, J.; DalNegro, G.; Oliosi, B.; Smith, K.J.; Thewlis, K.M.; Woollard, P.M.; Yusaf, S.P.
Deposited on	:	2010-11-10
Resolution	:	1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	3% 94% 6%
1	B	263	5% 95% 5%
1	C	263	4% 91% 7% ••

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	JAC	A	1302[B]	-	-	X	-
3	JAC	B	1302	-	-	X	-

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6955 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTAMATE RECEPTOR 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	263	2105	1335	348	405	17	0	11	1
1	B	263	2110	1339	351	404	16	0	10	1
1	C	260	2100	1339	348	397	16	0	12	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P19491
A	2	ALA	-	expression tag	UNP P19491
A	118	GLY	-	linker	UNP P19491
A	119	THR	-	linker	UNP P19491
A	242	SER	ASN	engineered mutation	UNP P19491
B	1	GLY	-	expression tag	UNP P19491
B	2	ALA	-	expression tag	UNP P19491
B	118	GLY	-	linker	UNP P19491
B	119	THR	-	linker	UNP P19491
B	242	SER	ASN	engineered mutation	UNP P19491
C	1	GLY	-	expression tag	UNP P19491
C	2	ALA	-	expression tag	UNP P19491
C	118	GLY	-	linker	UNP P19491
C	119	THR	-	linker	UNP P19491
C	242	SER	ASN	engineered mutation	UNP P19491

- Molecule 2 is GLUTAMIC ACID (three-letter code: GLU) (formula: $C_5H_9NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	10	5	1	4	0	0
			Total	C	N	O		
2	B	1	10	5	1	4	0	0
			Total	C	N	O		
2	C	1	10	5	1	4	0	0

- Molecule 3 is 1-({4-[3-(TRIFLUOROMETHYL)-6,7-DIHYDROPYRANO[4,3-C]PYRAZOL -1(4H)-YL]PHENYL}METHYL)-2-PYRROLIDINONE (three-letter code: JAC) (formula: $C_{18}H_{18}F_3N_3O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
3	A	1	52	36	6	6	4	0	1
			Total	C	F	N	O		
3	B	1	26	18	3	3	2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
4	A	1	5	4	1	0	0

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
5	A	1	1	1	0	0
			Total	Zn		
5	B	2	2	2	0	0
			Total	Zn		
5	C	2	2	2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	176	176	176	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	B	188	188	188	0	0
			Total	O		
6	C	158	158	158	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GLUTAMATE RECEPTOR 2

Chain A:



• Molecule 1: GLUTAMATE RECEPTOR 2

Chain B:





• Molecule 1: GLUTAMATE RECEPTOR 2

Chain C:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	114.43Å 163.36Å 47.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	93.66 – 1.60 37.15 – 1.60	Depositor EDS
% Data completeness (in resolution range)	97.9 (93.66-1.60) 97.8 (37.15-1.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.65 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.190 , 0.208 0.199 , 0.219	Depositor DCC
R_{free} test set	4595 reflections (4.00%)	wwPDB-VP
Wilson B-factor (Å^2)	14.5	Xtriage
Anisotropy	0.099	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 48.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	6955	wwPDB-VP
Average B, all atoms (Å^2)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.29% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4, JAC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.45	0/2142	0.58	1/2883 (0.0%)
1	B	0.47	0/2147	0.60	0/2890
1	C	0.43	0/2136	0.57	0/2872
All	All	0.45	0/6425	0.59	1/8645 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	148	ARG	NE-CZ-NH1	5.15	122.88	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2105	0	2093	15	0
1	B	2110	0	2114	8	0
1	C	2100	0	2117	17	0
2	A	10	0	5	1	0
2	B	10	0	5	1	0
2	C	10	0	5	1	0
3	A	52	0	36	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	26	0	18	9	0
4	A	5	0	0	0	0
5	A	1	0	0	0	0
5	B	2	0	0	0	0
5	C	2	0	0	1	0
6	A	176	0	0	14	0
6	B	188	0	0	13	0
6	C	158	0	0	11	0
All	All	6955	0	6393	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1302[B]:JAC:C1	6:C:2072:HOH:O	1.95	1.14
3:B:1302:JAC:C1	6:B:2080:HOH:O	2.06	1.03
3:B:1302:JAC:C12	6:B:2083:HOH:O	2.17	0.93
1:A:244[B]:GLN:CD	6:A:2158:HOH:O	2.07	0.90
1:A:244[B]:GLN:HG3	6:A:2158:HOH:O	1.73	0.88
3:A:1302[A]:JAC:C1	6:A:2074:HOH:O	2.21	0.87
1:A:244[B]:GLN:CG	6:A:2158:HOH:O	2.23	0.86
3:A:1302[B]:JAC:C12	6:C:2076:HOH:O	2.23	0.86
1:C:241:LEU:HD22	1:C:246:LEU:HD13	1.56	0.85
3:A:1302[A]:JAC:F1	6:A:2074:HOH:O	1.87	0.83
3:B:1302:JAC:H12A	6:B:2083:HOH:O	1.76	0.81
3:A:1302[B]:JAC:F1	6:C:2072:HOH:O	1.86	0.80
1:C:156:ASP:OD2	6:C:2103:HOH:O	2.02	0.77
3:A:1302[A]:JAC:C12	6:A:2076:HOH:O	2.33	0.77
5:C:1302:ZN:ZN	6:C:2155:HOH:O	1.38	0.73
1:B:183:LYS:HG3	6:B:2134:HOH:O	1.91	0.71
3:A:1302[A]:JAC:H12A	6:A:2076:HOH:O	1.88	0.71
3:A:1302[B]:JAC:F3	6:C:2072:HOH:O	1.87	0.70
3:B:1302:JAC:F3	6:B:2080:HOH:O	1.93	0.70
3:A:1302[B]:JAC:C2	6:C:2072:HOH:O	2.30	0.69
3:A:1302[B]:JAC:H12A	6:C:2076:HOH:O	1.91	0.69
3:B:1302:JAC:F2	6:B:2080:HOH:O	1.93	0.68
1:A:108[A]:SER:OG	6:A:2076:HOH:O	2.14	0.65
1:B:156:ASP:OD1	6:B:2116:HOH:O	2.15	0.64
3:A:1302[B]:JAC:H12	6:C:2076:HOH:O	1.93	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:125[A]:GLU:HG3	1:C:154:VAL:HG21	1.85	0.59
3:B:1302:JAC:C6	6:B:2170:HOH:O	2.52	0.58
1:C:21:LYS:H	1:C:21:LYS:HZ3	1.50	0.57
1:C:108[C]:SER:OG	6:C:2076:HOH:O	2.16	0.57
3:A:1302[A]:JAC:C6	6:A:2151:HOH:O	2.52	0.57
1:A:42:GLU:HG3	1:A:246:LEU:HD21	1.85	0.57
3:A:1302[A]:JAC:H6A	6:A:2151:HOH:O	2.05	0.56
1:B:106:PHE:CD1	1:B:107[A]:MET:HG3	2.40	0.56
3:B:1302:JAC:H6A	6:B:2170:HOH:O	2.04	0.56
1:B:23:HIS:CD2	1:B:26:LEU:HD12	2.41	0.55
3:B:1302:JAC:C2	6:B:2080:HOH:O	2.45	0.54
1:C:151[A]:LYS:HA	1:C:151[A]:LYS:HE2	1.88	0.54
1:C:21:LYS:H	1:C:21:LYS:NZ	2.10	0.50
1:A:81:GLY:HA2	6:A:2062:HOH:O	2.11	0.49
1:A:152:ILE:HD12	1:A:152:ILE:N	2.28	0.48
1:B:81:GLY:HA2	6:B:2067:HOH:O	2.13	0.48
1:C:42:GLU:CD	1:C:246:LEU:HD11	2.35	0.47
1:C:199:TYR:CZ	1:C:203:ARG:HD2	2.49	0.47
1:A:106:PHE:CD1	1:A:107[B]:MET:HG3	2.50	0.46
1:C:131:THR:HG22	1:C:131:THR:O	2.16	0.46
1:C:100:ILE:HD12	1:C:223:ALA:HB1	1.97	0.45
1:B:100:ILE:HD12	1:B:223:ALA:HB1	1.99	0.45
1:C:81:GLY:HA2	6:C:2058:HOH:O	2.16	0.45
1:C:179:ALA:O	1:C:183:LYS:HD2	2.17	0.45
1:A:115:ILE:HD11	1:A:211:VAL:HG11	1.99	0.44
1:A:241:LEU:HD22	1:A:246:LEU:HD23	1.99	0.44
3:B:1302:JAC:H12	6:B:2083:HOH:O	1.96	0.44
3:A:1302[B]:JAC:H6A	6:A:2022:HOH:O	2.18	0.44
1:A:91:THR:HG1	2:A:301:GLU:N	2.16	0.43
1:A:109:LEU:HD22	1:A:220:TYR:CE1	2.53	0.43
1:C:179:ALA:O	1:C:183:LYS:CD	2.67	0.43
1:A:244[B]:GLN:OE1	1:C:151[B]:LYS:HE2	2.19	0.42
1:A:115:ILE:CD1	1:A:211:VAL:HG11	2.50	0.42
1:B:145:GLU:HG2	6:B:2108:HOH:O	2.20	0.42
1:A:140[B]:SER:O	1:A:140[B]:SER:OG	2.23	0.42
1:C:42:GLU:HG3	1:C:246:LEU:HD11	2.01	0.41
3:A:1302[A]:JAC:H6A	6:A:2074:HOH:O	2.21	0.41
1:B:91:THR:HG1	2:B:301:GLU:N	2.19	0.41
3:A:1302[B]:JAC:C6	6:A:2022:HOH:O	2.68	0.41
1:C:91:THR:HG1	2:C:301:GLU:N	2.18	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/263 (103%)	269 (99%)	3 (1%)	0	100	100
1	B	271/263 (103%)	269 (99%)	2 (1%)	0	100	100
1	C	270/263 (103%)	268 (99%)	2 (1%)	0	100	100
All	All	813/789 (103%)	806 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/219 (102%)	223 (100%)	1 (0%)	91	84
1	B	226/219 (103%)	225 (100%)	1 (0%)	91	84
1	C	224/219 (102%)	220 (98%)	4 (2%)	59	36
All	All	674/657 (103%)	668 (99%)	6 (1%)	78	65

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	LEU
1	B	109	LEU
1	C	21	LYS
1	C	109	LEU
1	C	132	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	203	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 5 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	JAC	A	1302[B]	-	26,29,29	1.11	4 (15%)	29,43,43	1.43	6 (20%)
2	GLU	A	301	-	8,9,9	0.97	0	8,11,11	0.90	0
3	JAC	A	1302[A]	-	26,29,29	1.11	3 (11%)	29,43,43	1.60	7 (24%)
4	SO4	A	1304	5	4,4,4	0.41	0	6,6,6	0.66	0
2	GLU	C	301	-	8,9,9	1.05	1 (12%)	8,11,11	0.88	0
2	GLU	B	301	-	8,9,9	1.17	0	8,11,11	0.80	0
3	JAC	B	1302	-	26,29,29	1.11	4 (15%)	29,43,43	1.48	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	JAC	A	1302[B]	-	-	2/10/31/31	0/4/4/4
2	GLU	A	301	-	-	1/9/9/9	-
3	JAC	A	1302[A]	-	-	2/10/31/31	0/4/4/4
2	GLU	C	301	-	-	0/9/9/9	-
2	GLU	B	301	-	-	1/9/9/9	-
3	JAC	B	1302	-	-	3/10/31/31	0/4/4/4

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1302	JAC	C18-C8	-2.27	1.38	1.41
3	A	1302[A]	JAC	C18-C8	-2.24	1.38	1.41
2	C	301	GLU	OXT-C	-2.13	1.23	1.30
3	A	1302[B]	JAC	C18-C8	-2.12	1.38	1.41
3	B	1302	JAC	C9-C8	-2.12	1.38	1.41
3	A	1302[B]	JAC	C2-N1	-2.11	1.31	1.34
3	B	1302	JAC	C2-N1	-2.10	1.31	1.34
3	A	1302[A]	JAC	C9-C8	-2.07	1.39	1.41
3	B	1302	JAC	F2-C1	2.07	1.40	1.33
3	A	1302[A]	JAC	C2-N1	-2.03	1.31	1.34
3	A	1302[B]	JAC	F3-C1	2.03	1.40	1.33
3	A	1302[B]	JAC	C9-C8	-2.03	1.39	1.41

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	1302[A]	JAC	C1-C2-N1	3.30	123.63	119.69
3	B	1302	JAC	C10-C9-C8	-3.25	119.08	121.79
3	A	1302[A]	JAC	C10-C9-C8	-2.95	119.33	121.79
3	B	1302	JAC	C1-C2-N1	2.91	123.16	119.69
3	A	1302[B]	JAC	C1-C2-N1	2.85	123.10	119.69
3	A	1302[A]	JAC	C11-C12-N3	-2.81	108.73	113.15
3	B	1302	JAC	C9-C8-C18	2.79	120.47	116.27
3	A	1302[A]	JAC	C9-C8-C18	2.77	120.43	116.27
3	A	1302[B]	JAC	F2-C1-C2	-2.65	108.14	112.36
3	A	1302[B]	JAC	C10-C9-C8	-2.64	119.59	121.79
3	A	1302[B]	JAC	C9-C8-C18	2.58	120.14	116.27
3	A	1302[A]	JAC	F2-C1-C2	-2.56	108.30	112.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	1302[A]	JAC	C13-N3-C16	-2.47	110.58	113.76
3	B	1302	JAC	C13-N3-C16	-2.43	110.62	113.76
3	B	1302	JAC	C17-C18-C8	-2.20	119.95	121.79
3	A	1302[B]	JAC	C17-C18-C8	-2.18	119.98	121.79
3	A	1302[B]	JAC	F3-C1-C2	-2.12	109.00	112.36
3	A	1302[A]	JAC	C17-C18-C8	-2.08	120.06	121.79
3	B	1302	JAC	F1-C1-C2	-2.03	109.14	112.36

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1302[A]	JAC	F1-C1-C2-C7
3	A	1302[B]	JAC	F1-C1-C2-C7
3	B	1302	JAC	F3-C1-C2-C7
3	A	1302[A]	JAC	F3-C1-C2-C7
3	A	1302[B]	JAC	F3-C1-C2-C7
3	B	1302	JAC	F1-C1-C2-C7
3	B	1302	JAC	F2-C1-C2-C7
2	A	301	GLU	O-C-CA-CB
2	B	301	GLU	O-C-CA-CB

There are no ring outliers.

6 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1302[B]	JAC	9	0
2	A	301	GLU	1	0
3	A	1302[A]	JAC	7	0
2	C	301	GLU	1	0
2	B	301	GLU	1	0
3	B	1302	JAC	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand JAC A 1302 (B)



Ligand JAC A 1302 (A)



Ligand JAC B 1302



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	263/263 (100%)	0.21	9 (3%)	45	42	9, 16, 28, 36	0
1	B	263/263 (100%)	0.19	13 (4%)	29	27	8, 14, 27, 35	0
1	C	260/263 (98%)	0.23	11 (4%)	36	33	10, 17, 31, 38	0
All	All	786/789 (99%)	0.21	33 (4%)	36	33	8, 16, 29, 38	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	263	SER	14.9
1	B	263	SER	7.5
1	A	67	ASP	7.0
1	B	1	GLY	5.9
1	A	2	ALA	5.8
1	A	66	ALA	4.7
1	B	2	ALA	4.6
1	B	67	ASP	4.2
1	C	227	GLY	4.0
1	A	3	ASN	3.8
1	B	262	GLY	3.7
1	B	3	ASN	3.7
1	C	256	TYR	3.5
1	B	27	GLU	3.4
1	A	262	GLY	3.3
1	B	227	GLY	3.2
1	C	25	MET	3.1
1	C	131	THR	3.0
1	A	1	GLY	3.0
1	C	166	GLU	2.9
1	C	3	ASN	2.9
1	C	132	GLU	2.9
1	C	129	LYS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	256	TYR	2.6
1	C	258	LYS	2.5
1	B	66	ALA	2.5
1	A	131	THR	2.5
1	B	140[A]	SER	2.2
1	B	129	LYS	2.1
1	A	68	THR	2.1
1	C	229	SER	2.1
1	B	258	LYS	2.1
1	C	262	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	JAC	A	1302[A]	26/26	0.93	0.18	13,14,18,19	26
3	JAC	A	1302[B]	26/26	0.93	0.18	11,13,16,18	26
4	SO4	A	1304	5/5	0.94	0.14	16,18,25,27	0
2	GLU	A	301	10/10	0.96	0.10	9,10,12,12	0
2	GLU	B	301	10/10	0.96	0.08	8,9,10,10	0
3	JAC	B	1302	26/26	0.96	0.15	7,10,15,19	26
2	GLU	C	301	10/10	0.96	0.09	10,10,12,14	0
5	ZN	C	1302	1/1	0.98	0.04	20,20,20,20	0
5	ZN	C	1303	1/1	0.98	0.12	36,36,36,36	0
5	ZN	B	1304	1/1	0.99	0.05	15,15,15,15	0
5	ZN	A	1305	1/1	0.99	0.04	15,15,15,15	0
5	ZN	B	1303	1/1	0.99	0.04	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around JAC A 1302 (A): $2mF_o$-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)	
	

Electron density around JAC A 1302 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around JAC B 1302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.