

wwPDB X-ray Structure Validation Summary Report (i)

Jun 14, 2020 - 07:42 pm BST

PDB ID	:	2Y02
Title	:	TURKEY BETA1 ADRENERGIC RECEPTOR WITH STABILISING MU-
		TATIONS AND BOUND AGONIST CARMOTEROL
Authors	:	Warne, A.; Moukhametzianov, R.; Baker, J.G.; Nehme, R.; Edwards, P.C.;
		Leslie, A.G.W.; Schertler, G.F.X.; Tate, C.G.
Deposited on	:	2010-11-30
$\operatorname{Resolution}$:	2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.


The following versions of software and data (see references (1)) were used in the production of this report:

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455(2.60-2.60)
Sidechain outliers	138945	3455(2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			9%		
1	A	315	82%	9%	9%
			8%		
1	В	315	85%	9%	• 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	2CV	А	504	-	-	-	Х
3	2CV	А	505	-	-	-	Х



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NA	А	701	-	-	-	Х



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	А	286	Total 2269	C 1498	11	O 379	S 20	0	0	0
1	В	298	Total 2371	C 1560	± 1	O 395	S 20	0	0	0

• Molecule 1 is a protein called BETA-1 ADRENERGIC RECEPTOR.

Chain	Residue	Modelled	Actual	Comment	Reference
А	31	MET	-	expression tag	UNP P07700
A	32	GLY	-	expression tag	UNP P07700
А	68	SER	ARG	engineered mutation	UNP P07700
A	90	VAL	MET	engineered mutation	UNP P07700
А	116	LEU	CYS	engineered mutation	UNP P07700
А	227	ALA	TYR	engineered mutation	UNP P07700
A	?	-	CYS	deletion	UNP P07700
А	?	-	GLU	deletion	UNP P07700
A	?	-	GLY	deletion	UNP P07700
А	?	-	ARG	deletion	UNP P07700
А	?	-	PHE	deletion	UNP P07700
А	?	-	TYR	deletion	UNP P07700
А	?	-	GLY	deletion	UNP P07700
A	?	-	SER	deletion	UNP P07700
А	?	-	GLN	deletion	UNP P07700
А	?	-	GLU	deletion	UNP P07700
А	?	-	GLN	deletion	UNP P07700
А	?	-	PRO	deletion	UNP P07700
A	?	-	GLN	deletion	UNP P07700
А	?	-	PRO	deletion	UNP P07700
A	?	-	PRO	deletion	UNP P07700
А	?	-	PRO	deletion	UNP P07700
А	?	-	LEU	deletion	UNP P07700
А	?	-	PRO	deletion	UNP P07700
А	?	-	GLN	deletion	UNP P07700

There are 86 discrepancies between the modelled and reference sequences:




Chain	Residue	Modelled	Actual	Comment	Reference
А	?	-	HIS	deletion	UNP P07700
А	?	-	GLN	deletion	UNP P07700
А	?	-	PRO	deletion	UNP P07700
А	?	-	ILE	deletion	UNP P07700
А	?	-	LEU	deletion	UNP P07700
А	?	-	GLY	deletion	UNP P07700
А	?	-	ASN	deletion	UNP P07700
А	?	-	GLY	deletion	UNP P07700
А	?	-	ARG	deletion	UNP P07700
А	282	LEU	ALA	engineered mutation	UNP P07700
А	327	ALA	PHE	engineered mutation	UNP P07700
А	338	MET	PHE	engineered mutation	UNP P07700
А	358	ALA	CYS	engineered mutation	UNP P07700
А	369	HIS	-	expression tag	UNP P07700
А	370	HIS	-	expression tag	UNP P07700
А	371	HIS	-	expression tag	UNP P07700
А	372	HIS	-	expression tag	UNP P07700
А	373	HIS	-	expression tag	UNP P07700
В	31	MET	-	expression tag	UNP P07700
В	32	GLY	-	expression tag	UNP P07700
В	68	SER	ARG	engineered mutation	UNP P07700
В	90	VAL	MET	engineered mutation	UNP P07700
В	116	LEU	CYS	engineered mutation	UNP P07700
В	227	ALA	TYR	engineered mutation	UNP P07700
В	?	-	CYS	deletion	UNP P07700
В	?	-	GLU	deletion	UNP P07700
В	?	-	GLY	deletion	UNP P07700
В	?	-	ARG	deletion	UNP P07700
В	?	-	PHE	deletion	UNP P07700
В	?	-	TYR	deletion	UNP P07700
В	?	-	GLY	deletion	UNP P07700
В	?	-	SER	deletion	UNP P07700
В	?	-	GLN	deletion	UNP P07700
В	?	-	GLU	deletion	UNP P07700
В	?	-	GLN	deletion	UNP P07700
В	?		PRO	deletion	UNP P07700
В	?	-	GLN	deletion	UNP P07700
В	?	_	PRO	deletion	UNP P07700
В	?	-	PRO	deletion	UNP P07700
В	?	_	PRO	deletion	UNP P07700
В	?	-	LEU	deletion	UNP P07700
В	?	-	PRO	deletion	UNP P07700



Chain	Residue	Modelled	Actual	Comment	Reference
В	?	-	GLN	deletion	UNP P07700
В	?	-	HIS	deletion	UNP P07700
В	?	-	GLN	deletion	UNP P07700
В	?	-	PRO	deletion	UNP P07700
В	?	-	ILE	deletion	UNP P07700
В	?	-	LEU	deletion	UNP P07700
В	?	-	GLY	deletion	UNP P07700
В	?	-	ASN	deletion	UNP P07700
В	?	-	GLY	deletion	UNP P07700
В	?	-	ARG	deletion	UNP P07700
В	282	LEU	ALA	engineered mutation	UNP P07700
В	327	ALA	PHE	engineered mutation	UNP P07700
В	338	MET	PHE	engineered mutation	UNP P07700
В	358	ALA	CYS	engineered mutation	UNP P07700
В	369	HIS	-	expression tag	UNP P07700
В	370	HIS	-	expression tag	UNP P07700
В	371	HIS	-	expression tag	UNP P07700
В	372	HIS	-	expression tag	UNP P07700
В	373	HIS	-	expression tag	UNP P07700

• Molecule 2 is CHOLESTEROL HEMISUCCINATE (three-letter code: Y01) (formula: $C_{31}H_{50}O_4$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
2	Λ	1	Total C O	0	0	
			35 31 4	0	0	
2	Λ	1	Total C O	0	0	
	А	L	35 31 4	0	0	

Continued from previous page...

	ol	Chain	Residues	Atoms	ZeroOcc	AltConf
2		В	1	Total C O 35 31 4	0	0
2		В	1	Total C O 35 31 4	0	0

 Molecule 3 is HEGA-10 (three-letter code: 2CV) (formula: $\rm C_{18}H_{37}NO_7).$




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C N O 26 18 1 7	0	0
3	А	1	Total C 5 5	0	0
3	А	1	Total C N O 20 15 1 4	0	0
3	А	1	Total C N O 26 18 1 7	0	0
3	А	1	Total C N O 26 18 1 7	0	0
3	В	1	Total C N O 20 15 1 4	0	0
3	В	1	Total C N O 16 13 1 2	0	0
3	В	1	Total C N O 16 13 1 2	0	0
3	В	1	$\begin{array}{cccc} \mathrm{Total} & \mathrm{C} & \mathrm{N} & \mathrm{O} \\ 8 & 6 & 1 & 1 \end{array}$	0	0



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf		
3	В	1	Total 26	C 18	N 1	0 7	0	0

• Molecule 4 is CARMOTEROL (three-letter code: WHJ) (formula: $C_{21}H_{24}N_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	А	1	Total 27			0	0
4	В	1	Total 27			0	0

• Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total Na 1 1	0	0


• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	21	Total O 21 21	0	0
6	В	19	Total O 19 19	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: BETA-1 ADRENERGIC RECEPTOR



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	89.60Å 61.70 Å 101.40 Å	Depositor
a, b, c, α , β , γ	90.00° 109.30° 90.00°	Depositor
Resolution (Å)	95.73 - 2.60	Depositor
Resolution (A)	42.28 - 2.60	EDS
% Data completeness	96.7 (95.73-2.60)	Depositor
(in resolution range)	96.7(42.28-2.60)	EDS
R _{merge}	0.13	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.89 (at 2.61 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
D D .	0.229 , 0.268	Depositor
R, R_{free}	0.229 , (Not available)	DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor $(Å^2)$	38.0	Xtriage
Anisotropy	0.256	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.35, 66.3	EDS
L-test for $twinning^2$	$ \langle L \rangle = 0.48, \langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5064	wwPDB-VP
Average B, all atoms $(Å^2)$	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, 2CV, Y01, WHJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain		lengths	Bond	angles
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.52	0/2323	0.58	0/3167
1	В	0.53	0/2426	0.59	0/3303
All	All	0.53	0/4749	0.58	0/6470

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2269	0	2349	14	0
1	В	2371	0	2471	17	0
2	А	70	0	98	6	0
2	В	70	0	98	4	0
3	А	103	0	146	6	0
3	В	86	0	122	6	0
4	А	27	0	23	1	0
4	В	27	0	23	0	0
5	А	1	0	0	0	0
6	А	21	0	0	0	0
6	В	19	0	0	1	0
All	All	5064	0	5330	44	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:501:2CV:H271	3:B:505:2CV:H352	1.47	0.95
3:B:501:2CV:H601	3:B:501:2CV:H272	1.56	0.84
1:A:175:LEU:HD13	1:A:179:MET:HG3	1.65	0.78
3:B:501:2CV:C60	3:B:501:2CV:H272	2.17	0.75
1:B:175:LEU:HB2	6:B:2012:HOH:O	1.86	0.75

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{n} tiles
1	А	282/315~(90%)	273~(97%)	9~(3%)	0	100	100
1	В	296/315~(94%)	289~(98%)	7 (2%)	0	100	100
All	All	578/630~(92%)	562~(97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentile
1	А	247/273~(90%)	239~(97%)	8 (3%)	39 65
1	В	258/273~(94%)	$251 \ (97\%)$	7 (3%)	44 71
All	All	505/546~(92%)	490~(97%)	15 (3%)	41 67

5 of 15 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	А	235	LYS
1	А	329	ASN
1	В	236	GLU
1	А	229	ARG
1	В	202	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	\mathbf{Res}	Type
1	В	180	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 17 ligands modelled in this entry, 1 is monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



2Y02

Mal	Trees	Chain	Dec	Link	Bo	ond leng	ths	B	ond ang	gles
Mol	Type	Chain	\mathbf{Res}	LINK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	2CV	В	501	-	19, 19, 25	0.58	0	20,21,30	1.08	1(5%)
3	2CV	А	501	-	25,25,25	0.49	0	29,30,30	0.83	1 (3%)
3	2CV	В	505	-	25,25,25	0.73	0	29,30,30	0.69	1(3%)
2	Y01	В	402	-	35,38,38	1.30	2(5%)	54,57,57	1.50	9 (16%)
3	2CV	А	504	-	25,25,25	0.90	1 (4%)	29,30,30	0.74	0
3	2CV	В	502	-	15, 15, 25	0.81	1 (6%)	15,16,30	0.54	0
3	2CV	А	503	-	19, 19, 25	0.59	0	20,21,30	0.48	0
3	2CV	В	503	-	15, 15, 25	0.75	0	$15,\!16,\!30$	0.50	0
4	WHJ	В	601	-	28,29,29	1.43	4 (14%)	37,40,40	1.42	5 (13%)
4	WHJ	А	601	-	28,29,29	1.39	3 (10%)	37,40,40	1.63	8 (21%)
2	Y01	А	402	-	35,38,38	1.38	3 (8%)	54,57,57	1.23	6(11%)
3	2CV	А	505	-	25,25,25	0.65	0	29,30,30	0.73	1 (3%)
3	2CV	А	502	-	4,4,25	0.48	0	$3,\!3,\!30$	0.36	0
3	2CV	В	504	-	7,7,25	0.47	0	7,7,30	0.37	0
2	Y01	А	401	-	35,38,38	1.22	3 (8%)	54,57,57	1.96	12 (22%)
2	Y01	В	401	-	35,38,38	1.27	2 (5%)	54,57,57	1.79	10 (18%)

expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2CV	В	501	-	-	8/22/22/34	-
3	2CV	А	501	-	-	12/34/34/34	-
3	2CV	В	505	-	-	16/34/34/34	-
2	Y01	В	402	-	-	6/17/77/77	0/4/4/4
3	2CV	А	504	-	-	12/34/34/34	-
3	2CV	В	502	-	-	4/16/16/34	-
3	2CV	А	503	-	-	9/22/22/34	-
3	2CV	В	503	-	-	3/16/16/34	-
4	WHJ	В	601	-	-	4/15/15/15	0/3/3/3
4	WHJ	А	601	-	-	6/15/15/15	0/3/3/3
2	Y01	А	402	-	_	5/17/77/77	0/4/4/4



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	$2\mathrm{CV}$	А	505	-	-	14/34/34/34	-
3	$2\mathrm{CV}$	А	502	-	-	1/2/2/34	-
3	$2\mathrm{CV}$	В	504	-	-	$\frac{3/5/5/34}{3}$	-
2	Y01	А	401	-	-	9/17/77/77	0/4/4/4
2	Y01	В	401	-	-	10/17/77/77	0/4/4/4

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	В	402	Y01	CAK-CAI	-5.17	1.39	1.50
2	А	402	Y01	CAK-CAI	-4.76	1.39	1.50
2	А	401	Y01	CAK-CAI	-4.72	1.40	1.50
2	В	401	Y01	CAK-CAI	-4.37	1.40	1.50
4	В	601	WHJ	C3-C2	4.27	1.50	1.43

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	401	Y01	OAW-CBC-CAV	6.84	122.12	108.12
2	В	401	Y01	OAW-CBC-CAR	6.58	124.20	108.33
2	А	401	Y01	CBC-CAV-CAZ	-4.49	104.54	111.52
4	В	601	WHJ	C11-N2-C12	4.48	121.04	114.97
2	В	401	Y01	OAW-CAY-CAM	4.40	120.99	111.50

There are no chirality outliers.

5 of 122 torsion outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms
3	В	501	2CV	N33-C35-C60-O63
3	В	501	2CV	N33-C36-C37-C40
3	В	501	2CV	N33-C36-C37-O47
3	А	501	2CV	C36-C37-C40-C41
3	А	501	2CV	C36-C37-C40-O49

There are no ring outliers.


9 monomers are involved in 20 short contacts:

Mol	Chain	\mathbf{Res}	Type	Clashes	Symm-Clashes
3	В	501	2CV	3	0
3	А	501	$2\mathrm{CV}$	3	0




Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	505	2CV	3	0
2	В	402	Y01	1	0
4	А	601	WHJ	1	0
2	А	402	Y01	3	0
3	А	505	2CV	3	0
2	А	401	Y01	3	0
2	B	401	Y01	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# RSRZ {>}2$	$OWAB(Å^2)$	Q<0.9
1	А	286/315~(90%)	0.66	29 (10%) 7 4	11, 33, 75, 91	0
1	В	298/315~(94%)	0.52	25 (8%) 11 7	13, 32, 60, 80	0
All	All	584/630~(92%)	0.59	54 (9%) 9 6	11, 33, 67, 91	0

The worst 5 of 54 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	283	MET	8.0
1	А	280	VAL	6.7
1	А	281	MET	6.3
1	А	279	ARG	6.3
1	А	282	LEU	5.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




2Y02

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} extsf{-}\mathbf{factors}(\mathbf{A}^2)$	Q<0.9
3	$2\mathrm{CV}$	А	504	26/26	0.50	0.51	$40,\!72,\!90,\!91$	0
3	$2\mathrm{CV}$	А	505	26/26	0.57	0.44	40,85,98,99	0
3	$2\mathrm{CV}$	В	505	26/26	0.67	0.35	$30,\!77,\!88,\!90$	0
3	$2\mathrm{CV}$	А	503	20/26	0.72	0.39	$50,\!73,\!85,\!86$	0
3	$2\mathrm{CV}$	В	502	16/26	0.72	0.32	$20,\!52,\!73,\!78$	0
3	$2\mathrm{CV}$	А	501	26/26	0.73	0.29	$16,\!50,\!62,\!63$	0
3	$2\mathrm{CV}$	В	503	16/26	0.74	0.34	48,66,89,90	0
5	NA	А	701	1/1	0.75	0.49	$48,\!48,\!48,\!48$	0
3	$2\mathrm{CV}$	В	501	20/26	0.79	0.34	$17,\!51,\!62,\!64$	0
3	$2\mathrm{CV}$	А	502	5/26	0.81	0.28	$34,\!37,\!41,\!41$	0
2	Y01	А	402	35/35	0.82	0.28	$29,\!49,\!95,\!97$	0
3	$2\mathrm{CV}$	В	504	8/26	0.85	0.55	$58,\!64,\!67,\!69$	0
2	Y01	А	401	35/35	0.85	0.30	$15,\!27,\!47,\!52$	0
2	Y01	В	402	35/35	0.88	0.22	$35,\!41,\!74,\!75$	0
4	WHJ	А	601	27/27	0.91	0.26	21,24,67,70	0
2	Y01	В	401	35/35	0.91	0.26	$13,\!22,\!47,\!50$	0
4	WHJ	В	601	27/27	0.92	0.26	$16,\!25,\!49,\!54$	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









































6.5 Other polymers (i)

There are no such residues in this entry.

