



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 12:25 am BST

PDB ID : 5Y28
Title : Crystal structure of *H. pylori* HtrA with PDZ2 deletion
Authors : Zhang, Z.; Huang, Q.; Tao, X.
Deposited on : 2017-07-24
Resolution : 3.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1447 (3.10-3.06)
Clashscore	141614	1546 (3.10-3.06)
Ramachandran outliers	138981	1487 (3.10-3.06)
Sidechain outliers	138945	1486 (3.10-3.06)
RSRZ outliers	127900	1416 (3.10-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	357	 3% 64% 10% 25%
1	B	357	 50% 6% 43%
1	C	357	 3% 65% 12% 22%
2	G	4	 50% 50%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 5193 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Periplasmic serine endoprotease DegP-like.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	268	Total 1872	C 1166	N 323	O 380	S 3	0	0	0
1	B	202	Total 1393	C 871	N 238	O 281	S 3	0	0	0
1	C	278	Total 1899	C 1187	N 321	O 388	S 3	0	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	11	GLY	-	expression tag	UNP G2J5T2
A	12	PRO	-	expression tag	UNP G2J5T2
A	13	GLY	-	expression tag	UNP G2J5T2
A	14	TYR	-	expression tag	UNP G2J5T2
A	15	GLN	-	expression tag	UNP G2J5T2
A	16	ASP	-	expression tag	UNP G2J5T2
A	17	PRO	-	expression tag	UNP G2J5T2
B	11	GLY	-	expression tag	UNP G2J5T2
B	12	PRO	-	expression tag	UNP G2J5T2
B	13	GLY	-	expression tag	UNP G2J5T2
B	14	TYR	-	expression tag	UNP G2J5T2
B	15	GLN	-	expression tag	UNP G2J5T2
B	16	ASP	-	expression tag	UNP G2J5T2
B	17	PRO	-	expression tag	UNP G2J5T2
C	11	GLY	-	expression tag	UNP G2J5T2
C	12	PRO	-	expression tag	UNP G2J5T2
C	13	GLY	-	expression tag	UNP G2J5T2
C	14	TYR	-	expression tag	UNP G2J5T2
C	15	GLN	-	expression tag	UNP G2J5T2
C	16	ASP	-	expression tag	UNP G2J5T2
C	17	PRO	-	expression tag	UNP G2J5T2

- Molecule 2 is a protein called UNK-UNK-UNK-UNK.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	G	4	20	12	4	4	0	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	4	2	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O 1 1	0	0
4	B	1	Total O 1 1	0	0
4	C	3	Total O 3 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Periplasmic serine endoprotease DegP-like



- Molecule 1: Periplasmic serine endoprotease DegP-like



- Molecule 1: Periplasmic serine endoprotease DegP-like





- Molecule 2: UNK-UNK-UNK-UNK

Chain G: 50% 50%

A1
A2
A3
A4

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.91Å 91.50Å 120.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.88 – 3.09 43.88 – 3.09	Depositor EDS
% Data completeness (in resolution range)	99.5 (43.88-3.09) 99.5 (43.88-3.09)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.66 (at 3.06Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.257 , 0.298 0.258 , 0.299	Depositor DCC
R_{free} test set	959 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	87.4	Xtrriage
Anisotropy	0.477	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 67.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.002 for k,h,-l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5193	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/1888	0.46	0/2564
1	B	0.25	0/1405	0.45	0/1907
1	C	0.25	0/1914	0.45	0/2602
2	G	0.23	0/19	0.41	0/25
All	All	0.24	0/5226	0.45	0/7098

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1872	0	1774	23	0
1	B	1393	0	1289	17	0
1	C	1899	0	1748	23	0
2	G	20	0	22	4	0
3	A	4	0	6	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	3	0	0	0	0
All	All	5193	0	4839	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:GLN:N	1:B:23:GLN:OE1	2.15	0.73
1:C:141:GLY:HA3	1:C:265:LEU:HD22	1.72	0.69
1:C:212:THR:HG22	1:C:214:ALA:H	1.56	0.69
1:C:268:THR:HG22	1:C:269:GLY:H	1.61	0.66
1:C:226:ILE:HD12	1:C:230:GLY:HA2	1.78	0.64
1:A:211:GLN:HB3	1:A:249:ILE:HD11	1.79	0.63
1:A:130:GLY:O	1:A:132:ASN:N	2.34	0.60
1:A:319:ILE:HG22	1:A:320:THR:HG23	1.84	0.60
1:A:341:LEU:HD23	2:G:4:ALA:HB3	1.84	0.59
1:A:35:PRO:HG2	1:C:32:VAL:HG21	1.85	0.58
1:A:41:ILE:HG22	1:B:229:ARG:HD3	1.84	0.58
1:A:114:ASN:HB2	1:A:117:VAL:HG23	1.86	0.57
1:B:114:ASN:ND2	1:B:221:SER:OG	2.38	0.56
1:B:178:ILE:HG23	1:B:188:VAL:HG12	1.88	0.56
1:A:161:ILE:O	1:B:23:GLN:NE2	2.38	0.56
1:B:46:ASP:HA	1:B:49:LYS:HD3	1.88	0.55
1:C:114:ASN:HB2	1:C:117:VAL:HG23	1.89	0.54
1:A:189:THR:HG21	1:A:215:SER:HB3	1.91	0.52
1:B:114:ASN:HB2	1:B:117:VAL:HG23	1.90	0.52
1:B:174:LEU:HD11	1:B:190:GLN:HE21	1.74	0.52
1:B:173:ASP:OD1	1:C:31:ARG:NH2	2.44	0.51
1:C:327:VAL:HG11	1:C:333:LEU:HB2	1.94	0.49
1:C:274:GLY:HA3	1:C:362:THR:HB	1.94	0.49
1:C:304:LYS:HA	1:C:309:LYS:HD3	1.94	0.49
1:A:340:MET:HA	2:G:3:ALA:HB1	1.96	0.47
1:A:226:ILE:HD12	1:A:230:GLY:HA2	1.95	0.47
1:A:259:LYS:HG3	1:B:22:ILE:HG21	1.96	0.47
1:C:113:THR:HG22	1:C:114:ASN:H	1.81	0.46
1:B:31:ARG:HG2	1:C:43:SER:HB3	1.98	0.46
1:A:99:LEU:HD12	1:A:117:VAL:O	2.15	0.46
1:C:347:THR:HG22	1:C:359:PHE:HD1	1.80	0.46
1:B:25:MET:HG2	1:B:26:PRO:HD2	1.98	0.45
1:A:274:GLY:HA2	2:G:3:ALA:HB3	1.98	0.45
1:A:352:ARG:HA	1:A:352:ARG:NE	2.32	0.45
1:C:322:VAL:HG22	1:C:348:LEU:HD22	1.98	0.44
1:A:275:TYR:HB3	2:G:3:ALA:H	1.83	0.44
1:A:124:ILE:HG22	1:A:137:ALA:HB3	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:33:SER:HG	1:C:47:SER:HG	1.67	0.43
1:C:148:LEU:HD11	1:C:258:VAL:HG22	2.00	0.43
1:C:44:TYR:O	1:C:48:ILE:HG12	2.18	0.43
1:B:58:ILE:HG23	1:B:124:ILE:HG23	2.01	0.43
1:B:190:GLN:HB2	1:C:192:ILE:HD13	2.01	0.42
1:A:32:VAL:HG23	1:B:40:THR:HG21	2.02	0.42
1:C:225:LEU:HB2	1:C:236:ASN:OD1	2.19	0.42
1:A:216:ILE:HG23	1:A:236:ASN:HB3	2.01	0.42
1:A:353:ASP:OD1	1:A:354:LYS:N	2.52	0.42
1:C:48:ILE:HD12	1:C:226:ILE:HD11	2.02	0.42
1:B:164:SER:O	1:B:255:SER:OG	2.34	0.41
1:A:352:ARG:HB3	1:A:353:ASP:H	1.69	0.41
1:A:48:ILE:HG22	1:A:226:ILE:HD11	2.02	0.41
1:C:209:PHE:HA	1:C:254:PRO:HD3	2.03	0.41
1:A:41:ILE:HD11	1:C:31:ARG:CZ	2.51	0.40
1:C:59:SER:HA	1:C:98:ALA:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	258/357 (72%)	228 (88%)	21 (8%)	9 (4%)	3 18
1	B	192/357 (54%)	174 (91%)	15 (8%)	3 (2%)	9 35
1	C	266/357 (74%)	235 (88%)	22 (8%)	9 (3%)	3 19
2	G	2/4 (50%)	1 (50%)	1 (50%)	0	100 100
All	All	718/1075 (67%)	638 (89%)	59 (8%)	21 (3%)	4 22

All (21) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	131	SER
1	A	340	MET
1	C	134	GLU
1	C	278	VAL
1	C	341	LEU
1	C	342	PRO
1	A	342	PRO
1	A	350	VAL
1	C	240	ILE
1	A	317	ASP
1	A	349	LYS
1	B	124	ILE
1	B	272	GLU
1	C	270	LYS
1	C	277	GLY
1	A	360	THR
1	B	134	GLU
1	A	327	VAL
1	C	35	PRO
1	C	98	ALA
1	A	319	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/300 (64%)	186 (97%)	5 (3%)	46	72
1	B	140/300 (47%)	138 (99%)	2 (1%)	67	84
1	C	184/300 (61%)	183 (100%)	1 (0%)	88	94
All	All	515/900 (57%)	507 (98%)	8 (2%)	62	83

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	LEU
1	A	190	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	326	LYS
1	A	345	ARG
1	A	352	ARG
1	B	115	ASN
1	B	229	ARG
1	C	27	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	114	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	401	-	3,3,3	0.46	0	2,2,2	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	401	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/357 (75%)	0.11	10 (3%) 41 21	44, 77, 127, 143	0
1	B	202/357 (56%)	-0.09	1 (0%) 91 80	48, 76, 112, 132	0
1	C	278/357 (77%)	0.06	9 (3%) 47 25	48, 90, 114, 131	0
2	G	4/4 (100%)	1.20	0 100 100	99, 99, 103, 104	0
All	All	752/1075 (69%)	0.04	20 (2%) 54 29	44, 82, 118, 143	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	GLY	4.4
1	C	141	GLY	3.4
1	C	130	GLY	3.1
1	A	280	LEU	2.9
1	A	278	VAL	2.8
1	A	350	VAL	2.7
1	A	276	LEU	2.7
1	A	281	GLN	2.6
1	C	140	VAL	2.5
1	C	298	VAL	2.5
1	A	323	ASN	2.4
1	C	299	VAL	2.4
1	C	265	LEU	2.4
1	C	318	LEU	2.3
1	C	350	VAL	2.3
1	A	124	ILE	2.2
1	C	357	ARG	2.1
1	B	209	PHE	2.1
1	A	330	THR	2.0
1	A	331	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	A	401	4/4	0.90	0.29	78,82,83,87	0

6.5 Other polymers [i](#)

There are no such residues in this entry.