



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 26, 2020 – 08:11 AM BST

PDB ID : 6YCQ
Title : Crystal structure of the DNA binding domain of Arabidopsis thaliana Auxin Response Factor 1 (AtARF1) in complex with High Affinity DNA
Authors : Crespo, I.; Weijers, D.; Boer, D.R.
Deposited on : 2020-03-18
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	 7% 83% 12% ••
1	B	362	 7% 80% 14% 5%
2	C	21	 76% 24%
3	D	21	 76% 24%

2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 7154 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Auxin response factor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	347	2810	1769	506	520	15	0	4	0
1	B	343	2789	1755	504	516	14	0	5	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	ASN	-	expression tag	UNP Q8L7G0
A	357	SER	-	expression tag	UNP Q8L7G0
A	358	TYR	-	expression tag	UNP Q8L7G0
A	359	SER	-	expression tag	UNP Q8L7G0
A	360	GLN	-	expression tag	UNP Q8L7G0
A	361	SER	-	expression tag	UNP Q8L7G0
A	362	MET	-	expression tag	UNP Q8L7G0
B	356	ASN	-	expression tag	UNP Q8L7G0
B	357	SER	-	expression tag	UNP Q8L7G0
B	358	TYR	-	expression tag	UNP Q8L7G0
B	359	SER	-	expression tag	UNP Q8L7G0
B	360	GLN	-	expression tag	UNP Q8L7G0
B	361	SER	-	expression tag	UNP Q8L7G0
B	362	MET	-	expression tag	UNP Q8L7G0

- Molecule 2 is a DNA chain called 21-7A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	21	426	204	75	127	20	0	0	0


- Molecule 3 is a DNA chain called 21-7B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	D	21	429	204	84	121	20	0	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	5	Total	Cl	0	0
			5	5		
4	A	6	Total	Cl	0	0
			6	6		
4	C	1	Total	Cl	0	0
			1	1		

- Molecule 5 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			3	1	2		
5	A	1	Total	C	O	0	0
			3	1	2		
5	A	1	Total	C	O	0	0
			3	1	2		
5	B	1	Total	C	O	0	0
			3	1	2		
5	B	1	Total	C	O	0	0
			3	1	2		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	8	4	1	3	0	0
6	B	1	8	4	1	3	0	0
6	C	1	8	4	1	3	0	0

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
7	B	1	1	1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
8	A	295	295	295	0	0
8	B	239	239	239	0	0
8	C	62	62	62	0	0
8	D	52	52	52	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Auxin response factor 1




- Molecule 1: Auxin response factor 1



- Molecule 2: 21-7A



- Molecule 3: 21-7B



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.30Å 102.78Å 127.04Å 90.00° 98.04° 90.00°	Depositor
Resolution (Å)	47.57 – 1.65 47.57 – 1.65	Depositor EDS
% Data completeness (in resolution range)	57.0 (47.57-1.65) 54.1 (47.57-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.51 (at 1.65Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.172 , 0.199 0.172 , 0.199	Depositor DCC
R_{free} test set	3634 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.027 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7154	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, TRS, FMT, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/2889	0.60	3/3918 (0.1%)
1	B	0.42	0/2866	0.59	0/3887
2	C	0.82	0/476	1.04	0/733
3	D	0.84	0/482	0.95	0/742
All	All	0.50	0/6713	0.68	3/9280 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	267	ARG	NE-CZ-NH2	-6.53	117.03	120.30
1	A	267	ARG	CD-NE-CZ	5.83	131.76	123.60
1	A	267	ARG	NE-CZ-NH1	5.23	122.91	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	62	GLN	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2810	0	2742	36	0
1	B	2789	0	2723	42	0
2	C	426	0	239	4	0
3	D	429	0	236	5	0
4	A	6	0	0	0	0
4	B	5	0	0	0	0
4	C	1	0	0	0	0
5	A	9	0	3	1	0
5	B	6	0	2	0	0
6	A	8	0	12	0	0
6	B	8	0	12	0	0
6	C	8	0	12	0	0
7	B	1	0	0	0	0
8	A	295	0	0	11	1
8	B	239	0	0	10	1
8	C	62	0	0	3	0
8	D	52	0	0	2	0
All	All	7154	0	5981	84	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:16:DC:OP1	8:D:101:HOH:O	1.86	0.93
1:B:325:ASP:OD2	8:B:501:HOH:O	1.93	0.85
1:B:165:VAL:O	8:B:502:HOH:O	1.99	0.81
1:A:339:SER:OG	8:A:501:HOH:O	2.01	0.77
2:C:21:DA:OP2	8:C:201:HOH:O	2.03	0.76
1:A:16:SER:OG	8:A:502:HOH:O	2.03	0.75
1:A:295:LYS:HG3	1:A:307:ARG:HG3	1.70	0.73
2:C:15:DC:OP2	8:C:202:HOH:O	2.06	0.71
1:B:53:GLU:OE1	8:B:503:HOH:O	2.10	0.70
3:D:15:DC:OP2	8:D:102:HOH:O	2.09	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21[B]:ARG:NH2	8:A:508:HOH:O	2.26	0.68
1:B:299:GLU:OE1	8:B:504:HOH:O	2.11	0.67
1:B:63:GLN:O	8:B:505:HOH:O	2.12	0.67
1:A:94:GLN:OE1	8:A:503:HOH:O	2.12	0.67
1:A:212:ARG:HD2	1:A:216:GLU:OE2	1.96	0.66
1:B:317:GLU:HG3	1:B:332:LYS:HB2	1.78	0.65
1:B:87:GLU:HG3	1:B:88:THR:HG23	1.79	0.64
1:B:143:ARG:NH2	1:B:147:ASP:OD2	2.31	0.63
1:B:57:HIS:NE2	1:B:230:THR:HG23	2.18	0.59
1:A:60:LEU:HD22	1:A:243:HIS:CG	2.39	0.57
1:A:60:LEU:HG	1:A:61:GLU:O	2.04	0.57
3:D:16:DC:H2'	3:D:17:DG:C8	2.39	0.57
1:A:137:GLY:O	8:A:504:HOH:O	2.16	0.56
1:A:306:LYS:NZ	8:A:511:HOH:O	2.39	0.56
1:A:10:LYS:HB2	1:A:15:LEU:HD21	1.88	0.56
1:B:167:THR:HG22	1:B:173:GLU:HG2	1.88	0.56
1:A:90:GLU:HG3	1:A:267:ARG:HD2	1.89	0.55
1:B:143:ARG:C	1:B:143:ARG:HD3	2.27	0.54
1:A:60:LEU:HD12	1:A:61:GLU:H	1.74	0.53
1:A:21[A]:ARG:NH2	8:A:514:HOH:O	2.41	0.52
1:B:318:ASN:O	8:B:506:HOH:O	2.19	0.52
1:B:143:ARG:HG3	8:C:226:HOH:O	2.10	0.52
1:B:325:ASP:OD1	1:B:325:ASP:N	2.38	0.52
1:B:239[A]:SER:HA	1:B:242:MET:HG3	1.92	0.52
1:A:295:LYS:CG	1:A:307:ARG:HG3	2.41	0.51
1:B:13:GLY:N	8:B:514:HOH:O	2.43	0.50
1:B:63:GLN:HB3	8:B:505:HOH:O	2.11	0.50
1:A:60:LEU:HD22	1:A:243:HIS:ND1	2.27	0.50
1:B:117:GLU:CD	1:B:117:GLU:H	2.16	0.49
1:B:60:LEU:HD11	1:B:64:MET:HE3	1.95	0.49
1:A:94:GLN:HG3	1:A:263:PHE:CE1	2.48	0.48
1:A:66:SER:HB3	1:A:322:VAL:HG11	1.94	0.48
1:A:301:GLU:OE1	1:A:302:GLU:HB2	2.14	0.47
1:B:89:ASP:OD1	1:B:170:HIS:HE1	1.97	0.47
1:A:267:ARG:HB2	5:A:409:FMT:C	2.45	0.47
2:C:16:DC:H2'	2:C:17:DG:C8	2.49	0.47
1:B:239[B]:SER:HA	1:B:242:MET:HG3	1.96	0.46
1:A:177:ARG:NH1	8:A:521:HOH:O	2.45	0.46
1:B:24:TRP:CD1	1:B:82:ARG:HB3	2.50	0.46
1:B:209:ILE:HD12	1:B:209:ILE:N	2.31	0.46
1:A:66:SER:OG	1:A:66:SER:O	2.27	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:69:LEU:HD11	1:B:250:ALA:HB3	1.98	0.46
1:B:229:GLN:H	1:B:229:GLN:CD	2.18	0.46
1:B:143:ARG:O	1:B:143:ARG:HD3	2.16	0.46
1:B:275:VAL:HG21	1:B:350:TRP:CE2	2.51	0.45
1:A:24:TRP:CD1	1:A:82:ARG:HB3	2.51	0.45
1:B:56:MET:HE2	1:B:233:PRO:HG3	1.97	0.45
1:A:252:HIS:NE2	1:B:65:PRO:HD3	2.32	0.45
1:A:236:VAL:HG11	1:B:56:MET:HE3	1.99	0.45
1:A:22:GLU:HG3	1:A:121:VAL:HG21	1.98	0.45
1:A:293:ARG:HB2	1:A:355:LEU:HD11	1.98	0.45
3:D:9:DC:H2"	3:D:10:DA:C8	2.52	0.45
1:A:228:GLN:HB3	8:A:528:HOH:O	2.18	0.44
1:A:66:SER:CB	1:A:322:VAL:HG11	2.48	0.44
1:B:158:GLN:HA	1:B:159:PRO:HA	1.77	0.44
1:B:322:VAL:HG23	1:B:323:TRP:CE3	2.53	0.43
1:A:214:GLU:O	1:A:215:ASN:HB2	2.18	0.43
1:B:13:GLY:CA	8:B:514:HOH:O	2.66	0.43
1:A:46:GLU:OE1	8:A:505:HOH:O	2.22	0.43
1:A:295:LYS:HD2	1:A:353:GLU:HB2	2.01	0.42
2:C:13:DG:C2	3:D:10:DA:C2	3.07	0.42
1:B:293:ARG:HG2	1:B:293:ARG:HH11	1.83	0.42
1:B:167:THR:HA	1:B:172:SER:O	2.19	0.42
1:B:64:MET:HG2	1:B:244:ILE:HG13	2.00	0.42
1:B:139:PHE:HB2	1:B:193:TRP:CE2	2.55	0.42
1:A:108:SER:HB3	8:A:693:HOH:O	2.18	0.42
1:A:232:ILE:HD12	1:A:233:PRO:HD2	2.02	0.41
1:A:288:LEU:HD12	1:A:349:PRO:HB3	2.02	0.41
1:B:307:ARG:NH1	1:B:309[A]:SER:OG	2.53	0.41
1:A:293:ARG:HH11	1:A:293:ARG:HG2	1.86	0.41
1:B:14:VAL:N	8:B:514:HOH:O	2.39	0.41
1:B:15:LEU:HA	1:B:15:LEU:HD23	1.81	0.41
1:B:144[B]:ARG:NH2	1:B:144[B]:ARG:HG3	2.36	0.40
1:B:331:LEU:O	1:B:346:ARG:HA	2.22	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:675:HOH:O	8:A:747:HOH:O[2_546]	1.91	0.29
8:B:737:HOH:O	8:B:738:HOH:O[1_655]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	349/362 (96%)	340 (97%)	8 (2%)	1 (0%)	41	22
1	B	346/362 (96%)	336 (97%)	10 (3%)	0	100	100
All	All	695/724 (96%)	676 (97%)	18 (3%)	1 (0%)	51	31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	320	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/322 (97%)	309 (99%)	4 (1%)	69	50
1	B	311/322 (97%)	308 (99%)	3 (1%)	76	62
All	All	624/644 (97%)	617 (99%)	7 (1%)	73	57

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	SER
1	A	94	GLN
1	A	186	ARG
1	A	319	LYS
1	B	61	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	143	ARG
1	B	186	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	170	HIS
1	B	215	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 13 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	FMT	B	406	-	0,2,2	0.00	-	0,1,1	0.00	-
5	FMT	A	408	-	0,2,2	0.00	-	0,1,1	0.00	-
6	TRS	A	410	-	7,7,7	0.29	0	9,9,9	0.66	0
5	FMT	A	409	-	0,2,2	0.00	-	0,1,1	0.00	-
6	TRS	B	409	-	7,7,7	0.25	0	9,9,9	0.51	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	TRS	C	102	-	7,7,7	0.31	0	9,9,9	0.46	0
5	FMT	B	407	-	0,2,2	0.00	-	0,1,1	0.00	-
5	FMT	A	407	-	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	TRS	B	409	-	-	0/9/9/9	-
6	TRS	C	102	-	-	3/9/9/9	-
6	TRS	A	410	-	-	0/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	102	TRS	N-C-C1-O1
6	C	102	TRS	C3-C-C1-O1
6	C	102	TRS	C2-C-C1-O1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	409	FMT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	347/362 (95%)	-0.02	26 (7%) 14 14	18, 28, 77, 108	0
1	B	343/362 (94%)	0.08	27 (7%) 12 12	19, 32, 82, 111	0
2	C	21/21 (100%)	-0.46	0 100 100	28, 36, 57, 63	0
3	D	21/21 (100%)	-0.35	0 100 100	26, 42, 60, 77	0
All	All	732/766 (95%)	0.01	53 (7%) 15 14	18, 31, 79, 111	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	322	VAL	6.6
1	B	15	LEU	6.4
1	A	9	GLY	6.3
1	B	215	ASN	6.1
1	B	321	SER	5.2
1	A	63	GLN	5.2
1	A	10	LYS	5.0
1	B	324	HIS	5.0
1	B	14	VAL	4.7
1	A	11	PRO	4.5
1	A	14	VAL	4.4
1	A	229	GLN	4.2
1	A	12	GLY	3.7
1	A	319	LYS	3.6
1	A	321	SER	3.6
1	B	227	ARG	3.5
1	B	302	GLU	3.5
1	A	62	GLN	3.3
1	B	118	LYS	3.3
1	A	324	HIS	3.3
1	B	226	MET	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	319	LYS	3.2
1	A	322	VAL	3.2
1	B	300	GLY	3.1
1	B	64	MET	3.1
1	A	232	ILE	3.1
1	A	13	GLY	2.9
1	A	226	MET	2.9
1	A	325	ASP	2.8
1	B	65	PRO	2.7
1	B	13	GLY	2.7
1	B	63	GLN	2.7
1	A	15	LEU	2.6
1	A	302	GLU	2.6
1	B	66	SER	2.4
1	A	60	LEU	2.4
1	A	118	LYS	2.4
1	B	68	ASN	2.3
1	B	231	ASN	2.3
1	B	232	ILE	2.3
1	B	228	GLN	2.3
1	B	230	THR	2.2
1	A	301	GLU	2.2
1	B	120	THR	2.2
1	A	66	SER	2.2
1	A	231	ASN	2.1
1	A	320	SER	2.1
1	B	117	GLU	2.1
1	A	227	ARG	2.1
1	B	62	GLN	2.1
1	B	61	GLU	2.0
1	B	59	GLY	2.0
1	A	117	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	FMT	A	409	3/3	0.45	0.21	69,69,76,76	0
4	CL	A	403	1/1	0.74	0.11	76,76,76,76	0
5	FMT	B	407	3/3	0.80	0.18	45,45,47,56	0
5	FMT	A	407	3/3	0.86	0.09	48,48,55,55	0
6	TRS	C	102	8/8	0.90	0.15	39,49,56,59	0
6	TRS	B	409	8/8	0.90	0.08	30,40,46,47	0
5	FMT	B	406	3/3	0.91	0.09	48,48,52,55	0
5	FMT	A	408	3/3	0.93	0.14	49,49,52,54	0
4	CL	B	404	1/1	0.93	0.07	68,68,68,68	0
4	CL	B	401	1/1	0.94	0.12	63,63,63,63	0
4	CL	B	405	1/1	0.94	0.05	69,69,69,69	0
4	CL	B	402	1/1	0.95	0.08	40,40,40,40	0
6	TRS	A	410	8/8	0.95	0.11	30,40,42,43	0
4	CL	A	405	1/1	0.96	0.05	54,54,54,54	0
4	CL	B	403	1/1	0.96	0.17	56,56,56,56	0
4	CL	C	101	1/1	0.96	0.09	42,42,42,42	0
4	CL	A	402	1/1	0.98	0.07	58,58,58,58	0
4	CL	A	401	1/1	0.98	0.05	53,53,53,53	0
4	CL	A	406	1/1	0.98	0.09	39,39,39,39	0
7	NA	B	408	1/1	0.99	0.04	34,34,34,34	0
4	CL	A	404	1/1	0.99	0.11	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.