



## wwPDB EM Validation Summary Report ⓘ

Dec 17, 2022 – 01:21 pm GMT

PDB ID : 6YG8  
EMDB ID : EMD-10799  
Title : Cryo-EM structure of a BcsB pentamer in the context of an assembled Bcs macrocomplex  
Authors : Zouhir, S.; Krasteva, P.V.  
Deposited on : 2020-03-27  
Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.3


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	779	
1	B	779	
1	C	779	
1	D	779	
1	E	779	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 24330 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bacterial cellulose secretion regulator BcsB.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	643	Total 4996	C 3169	N 853	O 950	S 24	0	0
1	B	637	Total 4944	C 3136	N 842	O 943	S 23	0	0
1	C	637	Total 4944	C 3136	N 842	O 943	S 23	0	0
1	D	639	Total 4960	C 3146	N 844	O 947	S 23	0	0
1	E	576	Total 4486	C 2855	N 763	O 846	S 22	0	0


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bacterial cellulose secretion regulator BcsB




- Molecule 1: Bacterial cellulose secretion regulator BcsB






• Molecule 1: Bacterial cellulose secretion regulator BcsB



• Molecule 1: Bacterial cellulose secretion regulator BcsB



LEU TRP TYR ALA LEU ALA ASN HIS PRO CYS ILE LEU LEU LEU VAL ALA LEU LEU ALA ILE ILE VAL ILE LEU LEU LEU ALA TRP VAL LEU TRP ARG ARG ARG ARG ARG LEU ASN PRO ASP ASN GLU

● Molecule 1: Bacterial cellulose secretion regulator BcsB



MET LYS ARG LYS VAL LEU PHE TRP ILE CYS ALA VAL VAL MET ALA MET GLY MET SER PHE PHE PRO SER ILE PHE MET THR GLN ALA THR PRO ALA THR GLN PRO LEU ILE ASN ALA GLU PRO ALA VAL VAL VAL THR GLN ASN PRO GLN VAL GLN VAL MET PRO GLY VAL GLN VAL ASP

ALA PRO VAL P239 D240 GLN R67 F77 A78 Q79 I80 A81 F82 P83 S258 Q263 S86 M87 I98 S104 L115 L118 P119 S120 P121 S122 L123 L130 A167 M168 R169 L172 Y178 E183 T189 L190 W191 L192 D193 R196 K210 N211 D212 F220 R224

M233 P239 D240 L243 Q244 I249 S252 S258 Q263 N264 L272 P273 D274 R286 H293 H306 L315 L325 I332 N336 R340 Y344 R365 T371 Y403 LEU MET ARG SER THR GLY I410 D413 I414 Y418 K424

D430 L438 L443 SER SER LYS GLN ALA ASN ARG ARG LEU LEU LEU LEU ARG LEU LEU ILE PRO VAL GLN GLN GLY LEU LEU ASP I471 L476 G477 T479 M480 D485 M492 S496 V497 D498 N499 F503 Q507 S516 T517 Y523 F526 F535 A536 R537

A538 G539 M545 L548 S549 Q550 M555 P556 P559 M564 L568 M569 N570 V571 G572 F573 I574 G577 T578 L585 I601 I604 G605 G608 R675 S676 V677 A678 A679 D683 S684 F705 F708 A709 V710 ILE ARG GLU SER GLY ILE TRP THR GLU SER ILE TRP VAL LYS THR MET

ARG GLN THR PRO PHE LEU TRP TYR ALA VAL ASP GLU ASP ARG ALA ALA GLU THR ARG SER THR THR SER GLY ALA VAL I665 Q668 R675 S676 V677 A678 A679 D683 S684 F705 F708 A709 V710 ILE ARG GLU SER GLY ILE TRP THR GLU SER ILE TRP VAL LYS THR MET

PRO TRP PHE GLU ARG LEU TRP TYR ALA VAL ASP GLU ASP ARG ALA ALA GLU THR ARG SER THR THR SER GLY ALA VAL I665 Q668 R675 S676 V677 A678 A679 D683 S684 F705 F708 A709 V710 ILE ARG GLU SER GLY ILE TRP THR GLU SER ILE TRP VAL LYS THR MET

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	576455	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION; Gctf through the cryoSPARC v2 interface.	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	1.2	Depositor
Minimum defocus (nm)	750	Depositor
Maximum defocus (nm)	2750	Depositor
Magnification	130000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	55.833	Depositor
Minimum map value	-31.929	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.989	Depositor
Recommended contour level	4.5	Depositor
Map size ( $\text{\AA}$ )	431.51678, 431.51678, 431.51678	wwPDB
Map dimensions	410, 410, 410	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.05248, 1.05248, 1.05248	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/5111	0.46	0/6958
1	B	0.31	0/5057	0.45	0/6884
1	C	0.30	0/5057	0.45	0/6884
1	D	0.28	0/5073	0.45	0/6906
1	E	0.28	0/4590	0.45	0/6247
All	All	0.30	0/24888	0.45	0/33879

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4996	0	4951	78	0
1	B	4944	0	4892	79	0
1	C	4944	0	4892	59	0
1	D	4960	0	4907	70	0
1	E	4486	0	4431	67	0
All	All	24330	0	24073	325	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 325 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:426:SER:HB3	1:D:489:MET:HB3	1.66	0.77
1:D:420:MET:HB2	1:D:463:LEU:HD13	1.67	0.75
1:C:428:ARG:NH2	1:E:507:GLN:O	2.20	0.74
1:A:300:VAL:HG12	1:A:345:VAL:HG22	1.72	0.72
1:E:258:SER:O	1:E:263:GLN:NE2	2.23	0.72

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	639/779 (82%)	612 (96%)	27 (4%)	0	100	100
1	B	631/779 (81%)	606 (96%)	25 (4%)	0	100	100
1	C	631/779 (81%)	610 (97%)	21 (3%)	0	100	100
1	D	633/779 (81%)	599 (95%)	34 (5%)	0	100	100
1	E	566/779 (73%)	527 (93%)	39 (7%)	0	100	100
All	All	3100/3895 (80%)	2954 (95%)	146 (5%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	557/669 (83%)	557 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	551/669 (82%)	551 (100%)	0	100	100
1	C	551/669 (82%)	551 (100%)	0	100	100
1	D	553/669 (83%)	553 (100%)	0	100	100
1	E	499/669 (75%)	499 (100%)	0	100	100
All	All	2711/3345 (81%)	2711 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	569	ASN
1	E	334	GLN
1	E	537	ASN
1	A	696	ASN
1	A	79	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-10799. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 224



Y Index: 162




Z Index: 249


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)

### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 4.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate [\(i\)](#)




The volume at the recommended contour level is 245 nm<sup>3</sup>; this corresponds to an approximate mass of 222 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum [i](#)



\*Reported resolution corresponds to spatial frequency of 0.333 Å<sup>-1</sup>


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-10799 and PDB model 6YG8. Per-residue inclusion information can be found in section 3 on page 4.

### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 4.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (4.5).













## 9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 96% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (4.5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9648	 0.5680
A	 0.9694	 0.5800
B	 0.9743	 0.5800
C	 0.9724	 0.5730
D	 0.9485	 0.5580
E	 0.9587	 0.5450

