Full wwPDB X-ray Structure Validation Report (i)

Jun 24, 2024 – 07:16 PM EDT

PDB ID	:	5YJ2
Title	:	Crystal structure of Bacillus sp. TB-90 urate oxidase without dehydration
Authors	:	Hibi, T.; Itoh, T.; Nishiya, Y.
Deposited on	:	2017-10-07
Resolution	:	1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5722 (1.74-1.70)
Clashscore	141614	6152 (1.74-1.70)
Ramachandran outliers	138981	6051 (1.74-1.70)
Sidechain outliers	138945	6051 (1.74-1.70)
RSRZ outliers	127900	5629 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	2%, 92%, 5%
1	B	312	%, 94%
1	C	312	94%
1	D	312	2%, 93%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 20470 atoms, of which 9414 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uric acid degradation bifunctional protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	303	4746	1547	2341	400	454	4	0	1	0
1	B	303	4810	1576	2367	398	465	4	0	4	0
1	C	303	4776	1562	2351	397	462	4	0	2	0
1	D	303	4714	1545	2319	394	452	4	0	1	0

- Molecule 2 is 8-AZAXANTHINE (three-letter code: AZA) (formula: $C_4H_3N_5O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	14	4	3	5	2	0	0
2	B	1	14	4	3	5	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	C	1	14	4	3	5	2	0	0
			Total	C	H	N	O		
2	D	1	14	4	3	5	2	0	0

- Molecule 3 is OXYGEN MOLECULE (three-letter code: OXY) (formula: O_2) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	1	2	2	0	0
			Total	O		
3	B	1	2	2	0	0
			Total	O		
3	C	1	2	2	0	0
			Total	O		
3	D	1	2	2	0	0

- Molecule 4 is 2-METHOXYETHANOL (three-letter code: MXE) (formula: $C_3H_8O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	13	3	8	2	0	0
			Total	C	H	O		
4	B	1	13	3	8	2	0	0
			Total	C	H	O		
4	C	1	13	3	8	2	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	B	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	C	1	5	4	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	314	314	314	0	0
			Total	O		
6	B	336	336	336	0	0
			Total	O		
6	C	346	346	346	0	0
			Total	O		
6	D	315	315	315	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uric acid degradation bifunctional protein

Chain A:





• Molecule 1: Uric acid degradation bifunctional protein

Chain B:





• Molecule 1: Uric acid degradation bifunctional protein

Chain C:





• Molecule 1: Uric acid degradation bifunctional protein

Chain D:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	133.07Å 143.90Å 70.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.11 – 1.71 34.11 – 1.71	Depositor EDS
% Data completeness (in resolution range)	99.0 (34.11-1.71) 88.6 (34.11-1.71)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	0.80 (at 1.71Å)	Xtriage
Refinement program	PHENIX (1.12_2829)	Depositor
R, R_{free}	0.163 , 0.190 0.163 , 0.190	Depositor DCC
R_{free} test set	14403 reflections (9.94%)	wwPDB-VP
Wilson B-factor (Å^2)	15.4	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.42 , 53.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	20470	wwPDB-VP
Average B, all atoms (Å^2)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.89 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.9091e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: OXY, OCS, MXE, SO4, AZA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.32	0/2451	0.54	0/3326
1	B	0.32	0/2490	0.55	0/3379
1	C	0.32	0/2472	0.56	0/3354
1	D	0.32	0/2441	0.53	0/3312
All	All	0.32	0/9854	0.54	0/13371

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2405	2341	2338	7	0
1	B	2443	2367	2366	7	0
1	C	2425	2351	2347	6	0
1	D	2395	2319	2317	6	0
2	A	11	3	3	1	0
2	B	11	3	3	1	0
2	C	11	3	3	1	0
2	D	11	3	3	1	0
3	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	2	0	0	1	0
3	C	2	0	0	0	0
3	D	2	0	0	1	0
4	A	5	8	8	1	0
4	B	5	8	8	2	0
4	C	5	8	8	0	0
5	B	5	0	0	0	0
5	C	5	0	0	0	0
6	A	314	0	0	1	0
6	B	336	0	0	2	0
6	C	346	0	0	3	0
6	D	315	0	0	0	0
All	All	11056	9414	9404	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:201:ARG:HH12	2:B:401:AZA:HN3	1.45	0.65
1:D:201:ARG:HH12	2:D:401:AZA:HN3	1.44	0.64
1:C:201:ARG:HH12	2:C:402:AZA:HN3	1.47	0.62
1:A:201:ARG:HH12	2:A:401:AZA:HN3	1.47	0.61
1:C:214:LYS:NZ	6:C:505:HOH:O	2.35	0.59
1:B:275:GLN:HE21	4:B:404:MXE:H31	1.67	0.59
1:C:234:ARG:NH1	6:C:504:HOH:O	2.34	0.57
1:B:234:ARG:NH2	6:B:502:HOH:O	2.39	0.56
1:B:304:GLN:NE2	3:B:402:OXY:O2	2.43	0.51
1:C:234:ARG:NH2	6:C:509:HOH:O	2.44	0.49
1:A:275:GLN:OE1	4:A:403:MXE:H31	2.14	0.48
1:B:104:LEU:O	1:B:156:ARG:NH2	2.47	0.47
1:B:275:GLN:NE2	4:B:404:MXE:H31	2.30	0.47
1:D:114[B]:SER:OG	1:D:153:ASN:OD1	2.30	0.45
1:A:157:ASN:OD1	1:A:163:ASN:OD1	2.35	0.44
1:B:167:GLN:HG2	6:B:541:HOH:O	2.19	0.42
1:C:299:PRO:N	1:C:300:PRO:CD	2.82	0.42
1:A:178:LYS:NZ	1:A:244:THR:O	2.52	0.42
1:A:12:GLY:HA3	1:A:51:ILE:O	2.20	0.41
6:A:599:HOH:O	1:D:128:VAL:HG21	2.20	0.41
1:A:239:SER:O	1:A:243:GLU:HG3	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:PRO:N	1:A:300:PRO:CD	2.83	0.41
1:C:307:THR:HG1	1:D:8:VAL:N	2.19	0.41
1:D:304:GLN:NE2	3:D:402:OXY:O2	2.51	0.40
1:D:12:GLY:HA3	1:D:51:ILE:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/312 (96%)	296 (98%)	5 (2%)	0	100	100
1	B	304/312 (97%)	300 (99%)	4 (1%)	0	100	100
1	C	302/312 (97%)	297 (98%)	5 (2%)	0	100	100
1	D	301/312 (96%)	297 (99%)	4 (1%)	0	100	100
All	All	1208/1248 (97%)	1190 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/278 (92%)	254 (99%)	3 (1%)	71	58
1	B	258/278 (93%)	255 (99%)	3 (1%)	71	58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	257/278 (92%)	255 (99%)	2 (1%)	81	73
1	D	252/278 (91%)	249 (99%)	3 (1%)	71	58
All	All	1024/1112 (92%)	1013 (99%)	11 (1%)	73	62

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	LYS
1	A	177	ILE
1	A	241	PHE
1	B	77	LYS
1	B	177	ILE
1	B	241	PHE
1	C	177	ILE
1	C	241	PHE
1	D	77	LYS
1	D	177	ILE
1	D	241	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	275	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	OCS	D	305	1	7,8,9	0.89	0	6,11,13	1.82	2 (33%)
1	OCS	B	305	1	7,8,9	0.92	0	6,11,13	1.92	2 (33%)
1	OCS	C	305	1	7,8,9	0.91	0	6,11,13	1.66	2 (33%)
1	OCS	A	305	1	7,8,9	0.96	0	6,11,13	1.88	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	D	305	1	-	0/4/7/9	-
1	OCS	B	305	1	-	0/4/7/9	-
1	OCS	C	305	1	-	0/4/7/9	-
1	OCS	A	305	1	-	0/4/7/9	-

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	305	OCS	OD3-SG-CB	3.47	111.07	106.94
1	A	305	OCS	OD2-SG-CB	2.98	110.48	105.74
1	D	305	OCS	OD1-SG-CB	2.93	110.42	106.94
1	C	305	OCS	OD1-SG-CB	2.76	110.22	106.94
1	A	305	OCS	OD1-SG-CB	2.59	110.02	106.94
1	D	305	OCS	OD2-SG-CB	2.52	109.76	105.74
1	B	305	OCS	OD1-SG-CB	2.23	109.59	106.94
1	C	305	OCS	OD2-SG-CB	2.10	109.09	105.74

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	AZA	A	401	-	9,12,12	1.18	0	4,17,17	3.94	3 (75%)
5	SO4	B	403	-	4,4,4	0.14	0	6,6,6	0.18	0
4	MXE	A	403	-	4,4,4	0.44	0	3,3,3	0.30	0
2	AZA	B	401	-	9,12,12	1.19	0	4,17,17	4.38	3 (75%)
2	AZA	C	402	-	9,12,12	1.28	0	4,17,17	4.15	3 (75%)
3	OXY	D	402	-	1,1,1	0.39	0	-		
3	OXY	A	402	-	1,1,1	0.47	0	-		
5	SO4	C	401	-	4,4,4	0.17	0	6,6,6	0.11	0
3	OXY	B	402	-	1,1,1	0.36	0	-		
4	MXE	C	404	-	4,4,4	0.43	0	3,3,3	0.48	0
2	AZA	D	401	-	9,12,12	1.19	0	4,17,17	4.13	3 (75%)
4	MXE	B	404	-	4,4,4	0.47	0	3,3,3	0.34	0
3	OXY	C	403	-	1,1,1	0.33	0	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AZA	A	401	-	-	-	0/2/2/2
4	MXE	A	403	-	-	2/2/2/2	-
2	AZA	B	401	-	-	-	0/2/2/2
2	AZA	C	402	-	-	-	0/2/2/2
4	MXE	C	404	-	-	0/2/2/2	-
2	AZA	D	401	-	-	-	0/2/2/2
4	MXE	B	404	-	-	2/2/2/2	-

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	401	AZA	C2-N1-C6	7.57	121.53	115.14
2	D	401	AZA	C2-N1-C6	7.25	121.26	115.14
2	C	402	AZA	C2-N1-C6	7.10	121.14	115.14
2	A	401	AZA	C2-N1-C6	6.77	120.86	115.14
2	B	401	AZA	C5-C6-N1	-3.21	119.04	123.43
2	D	401	AZA	C5-C6-N1	-3.05	119.26	123.43
2	C	402	AZA	C5-C6-N1	-3.01	119.31	123.43
2	C	402	AZA	N9-N8-N7	-2.89	107.50	111.25
2	B	401	AZA	N9-N8-N7	-2.82	107.59	111.25
2	A	401	AZA	C5-C6-N1	-2.81	119.58	123.43
2	A	401	AZA	N9-N8-N7	-2.78	107.64	111.25
2	D	401	AZA	N9-N8-N7	-2.33	108.22	111.25

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	403	MXE	C1-C2-O2-C3
4	A	403	MXE	O1-C1-C2-O2
4	B	404	MXE	C1-C2-O2-C3
4	B	404	MXE	O1-C1-C2-O2

There are no ring outliers.

8 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	AZA	1	0
4	A	403	MXE	1	0
2	B	401	AZA	1	0
2	C	402	AZA	1	0
3	D	402	OXY	1	0
3	B	402	OXY	1	0
2	D	401	AZA	1	0
4	B	404	MXE	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand AZA A 401



Ligand AZA B 401



Bond lengths



Bond angles



Torsions



Rings

Ligand AZA C 402



Bond lengths



Bond angles



Torsions



Rings

Ligand OXY D 402



Bond lengths



Bond angles



Torsions



Rings

Ligand OXY A 402	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand OXY B 402	
 Bond lengths	Bond angles
 Torsions	Rings

Ligand AZA D 401



Ligand OXY C 403



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	302/312 (96%)	-0.26	7 (2%)	60	65	12, 18, 32, 61	0
1	B	302/312 (96%)	-0.25	3 (0%)	82	85	11, 17, 33, 57	0
1	C	302/312 (96%)	-0.26	1 (0%)	94	95	12, 17, 30, 50	0
1	D	302/312 (96%)	-0.24	5 (1%)	70	74	12, 18, 35, 49	0
All	All	1208/1248 (96%)	-0.25	16 (1%)	77	81	11, 17, 33, 61	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	287	PRO	3.7
1	D	130	ASN	3.5
1	A	130	ASN	3.5
1	B	291	GLY	3.3
1	B	287	PRO	3.3
1	A	131	GLY	3.2
1	D	222	THR	2.7
1	B	289	SER	2.6
1	D	288	GLU	2.5
1	A	129	LYS	2.4
1	A	223	ASN	2.3
1	A	288	GLU	2.2
1	D	215	ASN	2.2
1	A	222	THR	2.2
1	D	287	PRO	2.1
1	C	10	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	OCS	A	305	9/10	0.86	0.16	20,30,54,60	0
1	OCS	B	305	9/10	0.89	0.14	16,21,50,51	0
1	OCS	D	305	9/10	0.90	0.12	18,25,55,57	0
1	OCS	C	305	9/10	0.94	0.12	15,24,55,57	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	MXE	C	404	5/5	0.65	0.20	35,43,51,51	0
4	MXE	B	404	5/5	0.73	0.19	38,46,55,56	0
4	MXE	A	403	5/5	0.74	0.20	36,44,55,55	0
5	SO4	C	401	5/5	0.92	0.14	41,48,52,53	0
5	SO4	B	403	5/5	0.94	0.13	37,41,45,50	0
3	OXY	A	402	2/2	0.94	0.14	9,9,9,11	0
3	OXY	D	402	2/2	0.95	0.18	6,6,6,14	0
3	OXY	B	402	2/2	0.95	0.16	6,6,6,15	0
3	OXY	C	403	2/2	0.95	0.16	5,5,5,16	0
2	AZA	B	401	11/11	0.96	0.06	11,13,15,16	0
2	AZA	A	401	11/11	0.96	0.08	13,14,17,18	0
2	AZA	D	401	11/11	0.97	0.07	13,14,16,18	0
2	AZA	C	402	11/11	0.98	0.08	12,14,15,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OXY A 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OXY D 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OXY B 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OXY C 403:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AZA B 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AZA A 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around AZA D 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AZA C 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.