

Full wwPDB X-ray Structure Validation Report (i)

Aug 16, 2023 – 06:58 AM EDT

PDB ID : 1YR2

Title: Structural and Mechanistic Analysis of Two Prolyl Endopeptidases: Role of

Inter-Domain Dynamics in Catalysis and Specificity

Authors: Shan, L.; Mathews, I.I.; Khosla, C.

Deposited on : 2005-02-02

Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain			
1	A	741	86%	5%	8%	



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5929 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called prolyl oligopeptidase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	680	Total	C	N	0	S	0	1	0
			5256	3360	917	973	О			

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	667	PHE	LEU	see remark 999	UNP Q9ZNM8
A	724	TRP	-	cloning artifact	UNP Q9ZNM8
A	725	SER	-	cloning artifact	UNP Q9ZNM8
A	726	SER	_	cloning artifact	UNP Q9ZNM8
A	727	VAL	-	cloning artifact	UNP Q9ZNM8
A	728	ASP	_	cloning artifact	UNP Q9ZNM8
A	729	LYS	-	cloning artifact	UNP Q9ZNM8
A	730	LEU	-	cloning artifact	UNP Q9ZNM8
A	731	ALA	-	cloning artifact	UNP Q9ZNM8
A	732	ALA	-	cloning artifact	UNP Q9ZNM8
A	733	ALA	-	cloning artifact	UNP Q9ZNM8
A	734	LEU	-	cloning artifact	UNP Q9ZNM8
A	735	GLU	-	cloning artifact	UNP Q9ZNM8
A	736	HIS	-	expression tag	UNP Q9ZNM8
A	737	HIS	-	expression tag	UNP Q9ZNM8
A	738	HIS	-	expression tag	UNP Q9ZNM8
A	739	HIS	-	expression tag	UNP Q9ZNM8
A	740	HIS	-	expression tag	UNP Q9ZNM8
A	741	HIS	-	expression tag	UNP Q9ZNM8

• Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	660	Total O 661 661	0	1




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: prolyl oligopeptidase





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	53.34Å 91.22Å 79.79Å	Depositor	
a, b, c, α , β , γ	90.00° 91.00° 90.00°	Depositor	
Resolution (Å)	30.00 - 1.80	Depositor	
% Data completeness	98.6 (30.00-1.80)	Depositor	
(in resolution range)	30.0 (80.00 1.00)		
R_{merge}	(Not available)	Depositor	
R_{sym}	0.07	Depositor	
Refinement program	REFMAC 5.1.24	Depositor	
R, R_{free}	0.161 , 0.186	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	5929	wwPDB-VP	
Average B, all atoms (Å ²)	18.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	nd lengths	Во	ond angles
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.78	1/5414 (0.0%)	0.89	$15/7385 \ (0.2\%)$

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	${f Z}$	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
1	A	180	SER	CB-OG	-5.23	1.35	1.42

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^{o})$
1	A	345	ARG	NE-CZ-NH2	-6.41	117.09	120.30
1	A	71	ASP	CB-CG-OD2	6.09	123.78	118.30
1	A	141	ASP	CB-CG-OD2	6.03	123.73	118.30
1	A	107	ASP	CB-CG-OD2	5.95	123.66	118.30
1	A	166	ASP	CB-CG-OD2	5.80	123.52	118.30
1	A	728	ASP	CB-CG-OD2	5.67	123.41	118.30
1	A	324	ASP	CB-CG-OD1	5.52	123.27	118.30
1	A	345	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	618	ASP	CB-CG-OD1	5.44	123.20	118.30
1	A	347	ASP	CB-CG-OD2	5.28	123.05	118.30
1	A	356	ASP	CB-CG-OD2	5.24	123.02	118.30
1	A	590	ASP	CB-CG-OD2	5.20	122.98	118.30
1	A	183	ASP	CB-CG-OD2	5.18	122.96	118.30
1	A	542	ASP	CB-CG-OD2	5.09	122.88	118.30
1	A	458	ASP	CB-CG-OD2	5.02	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5256	0	5108	13	0
2	A	12	0	16	0	0
3	A	661	0	0	5	0
All	All	5929	0	5124	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:239:LEU:N	3:A:1397:HOH:O	2.33	0.61
1:A:280:ARG:NH1	3:A:1035:HOH:O	2.39	0.54
1:A:610:PHE:O	1:A:611:THR:C	2.46	0.53
1:A:572:GLU:CG	3:A:1165:HOH:O	2.58	0.52
1:A:249:ARG:NE	3:A:1400:HOH:O	2.34	0.49
1:A:572:GLU:HG3	3:A:1165:HOH:O	2.12	0.48
1:A:331:TRP:CZ2	1:A:393:LEU:HB3	2.51	0.45
1:A:102:MET:SD	1:A:706:THR:HG21	2.59	0.42
1:A:436:ALA:O	1:A:444:THR:HA	2.19	0.42
1:A:280:ARG:HH11	1:A:280:ARG:CG	2.32	0.42
1:A:280:ARG:HH11	1:A:280:ARG:HB3	1.85	0.42
1:A:206:LEU:HD21	1:A:248:HIS:CG	2.54	0.42
1:A:286:SER:HB2	1:A:295:THR:OG1	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	673/741 (91%)	651 (97%)	20 (3%)	2 (0%)	41 27	

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	290	THR
1	A	611	THR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric		Percentiles	
1	A	541/584 (93%)	532 (98%)	9 (2%)	60 51	

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	LYS
1	A	91	GLN
1	A	177	LEU
1	A	204	ASP
1	A	229	GLU
1	A	280	ARG
1	A	291	ASP
1	A	312	LEU
1	A	619	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

\mathbf{M}	۵1	Type	Chain	Res Link		В	ond leng	$_{ m gths}$	В	ond ang	gles
101	OI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	2	GOL	A	750	-	5,5,5	0.41	0	5,5,5	0.33	0
2	2	GOL	A	755	-	5,5,5	0.77	0	5,5,5	1.07	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	750	-	-	2/4/4/4	-
2	GOL	A	755	-	-	1/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	755	GOL	O2-C2-C1	-2.14	99.68	109.12

There are no chirality outliers.

All (3) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	750	GOL	O1-C1-C2-C3
2	A	750	GOL	O1-C1-C2-O2
2	A	755	GOL	C1-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

