

# Full wwPDB X-ray Structure Validation Report (i)

### Feb 22, 2021 – 03:08 PM GMT

PDB ID : 6YW9

Title: Arabidopsis aspartate transcarbamoylase mutant F161A complex with PALA

Authors: Ramon Maiques, S.; Del Cano Ochoa, F.; Bellin, L.; Mohlmann, T.

Deposited on : 2020-04-29

Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

 $\begin{array}{ccc} EDS & : & 2.17.1.dev1 \\ buster\text{-report} & : & 1.1.7 \ (2018) \end{array}$ 

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove) roteins) : Engh & Huber (2001)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.17.1.dev1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\mathring{\rm A})}) \end{array}$
$R_{free}$	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
4		000	2%	
1	A	332	89%	5% • 5%
			3%	
1	В	332	86%	8% 6%
			4%	
1	С	332	86%	8% • 6%



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 16087 atoms, of which 7691 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called PYRB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	Λ	215	Total	С	Н	N	О	S	0	17	0
1	Λ	315	5128	1617	2577	428	492	14	U	11	U
1	D	311	Total	С	Н	N	О	S	0	17	0
1	Ъ	911	5065	1598	2540	427	487	13	0	11	
1	С	311	Total	С	Н	N	О	S	0	16	0
1		911	5066	1597	2549	424	482	14	U	10	U

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	59	MET	-	initiating methionine	UNP A0A178VJE3
A	60	GLY	-	expression tag	UNP A0A178VJE3
A	61	SER	-	expression tag	UNP A0A178VJE3
A	62	SER	-	expression tag	UNP A0A178VJE3
A	63	HIS	-	expression tag	UNP A0A178VJE3
A	64	HIS	=	expression tag	UNP A0A178VJE3
A	65	HIS	-	expression tag	UNP A0A178VJE3
A	66	HIS	-	expression tag	UNP A0A178VJE3
A	67	HIS	-	expression tag	UNP A0A178VJE3
A	68	HIS	-	expression tag	UNP A0A178VJE3
A	69	SER	-	expression tag	UNP A0A178VJE3
A	70	SER	-	expression tag	UNP A0A178VJE3
A	71	GLY	-	expression tag	UNP A0A178VJE3
A	72	LEU	-	expression tag	UNP A0A178VJE3
A	73	GLU	-	expression tag	UNP A0A178VJE3
A	74	VAL	-	expression tag	UNP A0A178VJE3
A	75	LEU	-	expression tag	UNP A0A178VJE3
A	76	PHE	=	expression tag	UNP A0A178VJE3
A	77	GLN	-	expression tag	UNP A0A178VJE3
A	78	GLY	-	expression tag	UNP A0A178VJE3
A	79	PRO	=	expression tag	UNP A0A178VJE3
A	80	HIS	-	expression tag	UNP A0A178VJE3
A	81	MET	-	expression tag	UNP A0A178VJE3





Continued from previous page...


Chain	Residue	Modelled    Woodelled   Modelled   Modelled	Actual	Comment	Reference
A	161	ALA	PHE	engineered mutation	UNP A0A178VJE3
В	59	MET	-	initiating methionine	UNP A0A178VJE3
В	60	GLY	-	expression tag	UNP A0A178VJE3
В	61	SER	-	expression tag	UNP A0A178VJE3
В	62	SER	-	expression tag	UNP A0A178VJE3
В	63	HIS	-	expression tag	UNP A0A178VJE3
В	64	HIS	-	expression tag	UNP A0A178VJE3
В	65	HIS	-	expression tag	UNP A0A178VJE3
В	66	HIS	-	expression tag	UNP A0A178VJE3
В	67	HIS	-	expression tag	UNP A0A178VJE3
В	68	HIS	-	expression tag	UNP A0A178VJE3
В	69	SER	-	expression tag	UNP A0A178VJE3
В	70	SER	-	expression tag	UNP A0A178VJE3
В	71	GLY	-	expression tag	UNP A0A178VJE3
В	72	LEU	-	expression tag	UNP A0A178VJE3
В	73	GLU	-	expression tag	UNP A0A178VJE3
В	74	VAL	-	expression tag	UNP A0A178VJE3
В	75	LEU	=	expression tag	UNP A0A178VJE3
В	76	PHE	-	expression tag	UNP A0A178VJE3
В	77	GLN	-	expression tag	UNP A0A178VJE3
В	78	GLY	-	expression tag	UNP A0A178VJE3
В	79	PRO	-	expression tag	UNP A0A178VJE3
В	80	HIS	-	expression tag	UNP A0A178VJE3
В	81	MET	ı	expression tag	UNP A0A178VJE3
В	161	ALA	PHE	engineered mutation	UNP A0A178VJE3
С	59	MET	-	initiating methionine	UNP A0A178VJE3
С	60	GLY	-	expression tag	UNP A0A178VJE3
С	61	SER	-	expression tag	UNP A0A178VJE3
С	62	SER	-	expression tag	UNP A0A178VJE3
С	63	HIS	-	expression tag	UNP A0A178VJE3
С	64	HIS	-	expression tag	UNP A0A178VJE3
С	65	HIS	-	expression tag	UNP A0A178VJE3
С	66	HIS	-	expression tag	UNP A0A178VJE3
С	67	HIS	-	expression tag	UNP A0A178VJE3
С	68	HIS	-	expression tag	UNP A0A178VJE3
С	69	SER	-	expression tag	UNP A0A178VJE3
С	70	SER	-	expression tag	UNP A0A178VJE3
С	71	GLY	=	expression tag	UNP A0A178VJE3
С	72	LEU	ı	expression tag	UNP A0A178VJE3
С	73	GLU	-	expression tag	UNP A0A178VJE3
С	74	VAL	-	expression tag	UNP A0A178VJE3
С	75	LEU	-	expression tag	UNP A0A178VJE3



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	76	PHE	-	expression tag	UNP A0A178VJE3
С	77	GLN	-	expression tag	UNP A0A178VJE3
С	78	GLY	-	expression tag	UNP A0A178VJE3
С	79	PRO	-	expression tag	UNP A0A178VJE3
С	80	HIS	-	expression tag	UNP A0A178VJE3
С	81	MET	ı	expression tag	UNP A0A178VJE3
С	161	ALA	PHE	engineered mutation	UNP A0A178VJE3


• Molecule 2 is N-(PHOSPHONACETYL)-L-ASPARTIC ACID (three-letter code: PAL) (formula:  $C_6H_{10}NO_8P$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C H N O P 22 6 6 1 8 1	0	0
2	В	1	Total C H N O P 22 6 6 1 8 1	0	0
2	С	1	Total C H N O P 22 6 6 1 8 1	0	0

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	Λ	1	Total	С	Н	О	0	0
3	A	1	13	3	7	3	0	0


### • Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	267	Total O 267 267	0	0
4	В	275	Total O 275 275	0	0
4	С	207	Total O 207 207	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	80.27Å 98.28Å 138.78Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.30 - 1.68	Depositor
resolution (A)	46.30 - 1.59	EDS
% Data completeness	100.0 (46.30-1.68)	Depositor
(in resolution range)	92.9 (46.30-1.59)	EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.55 (at 1.59Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
D D.	0.151 , 0.174	Depositor
$R, R_{free}$	0.151 , $0.174$	DCC
$R_{free}$ test set	6278 reflections $(4.29%)$	wwPDB-VP
Wilson B-factor $(\mathring{A}^2)$	21.8	Xtriage
Anisotropy	0.191	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.41 , 48.8	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	16087	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	nd lengths	Bond angles		
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.51	0/2648	0.69	1/3561~(0.0%)	
1	В	0.49	0/2620	0.69	$1/3526 \ (0.0\%)$	
1	С	0.52	1/2602~(0.0%)	0.74	$4/3500 \ (0.1\%)$	
All	All	0.51	$1/7870 \ (0.0\%)$	0.71	6/10587 (0.1%)	

#### All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	${ m Observed}({ m \AA})$	$\operatorname{Ideal}( ext{\AA})$
1	С	107	GLU	CD-OE1	-9.60	1.15	1.25

#### All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
1	С	107	GLU	OE1-CD-OE2	-18.62	100.96	123.30
1	В	352	ARG	NE-CZ-NH2	-6.20	117.20	120.30
1	A	352	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	С	352	ARG	NE-CZ-NH2	-5.83	117.38	120.30
1	С	107	GLU	CG-CD-OE2	5.63	129.56	118.30
1	С	107	GLU	CG-CD-OE1	5.58	129.46	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2551	2577	2585	17	0
1	В	2525	2540	2559	16	0
1	С	2517	2549	2578	19	0
2	A	16	6	6	0	0
2	В	16	6	6	0	0
2	С	16	6	6	0	0
3	A	6	7	7	0	0
4	A	267	0	0	3	0
4	В	275	0	0	5	0
4	С	207	0	0	2	0
All	All	8396	7691	7747	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:342:LYS:NZ	4:A:501:HOH:O	1.85	1.09
1:B:334:LYS:NZ	4:B:501:HOH:O	2.15	0.71
1:A:172:ASP:O	1:A:176:THR:HG23	1.92	0.69
4:B:503:HOH:O	1:C:154[A]:THR:HG22	1.92	0.69
1:C:111:ILE:HG13	4:C:502:HOH:O	1.92	0.68
4:B:507:HOH:O	1:C:154[A]:THR:HG23	1.94	0.68
4:A:509:HOH:O	1:B:154[A]:THR:HG23	1.95	0.65
1:A:176:THR:HG21	1:C:136:ARG:NH2	2.13	0.64
1:A:176:THR:HG21	1:C:136:ARG:HH21	1.65	0.62
1:B:80:HIS:N	4:B:504:HOH:O	2.32	0.62
1:A:125:LEU:CD2	1:A:151[B]:GLU:HB2	2.31	0.60
1:A:154[A]:THR:HG23	4:C:501:HOH:O	2.01	0.60
1:C:170:LEU:HD11	1:C:186:MET:SD	2.41	0.59
1:B:170:LEU:HD11	1:B:186:MET:SD	2.42	0.59
1:A:93:PHE:CZ	1:A:220[B]:LEU:HD11	2.39	0.57
1:C:323:TYR:CZ	1:C:327:ARG:HD2	2.41	0.55
1:B:80:HIS:N	4:B:505:HOH:O	2.41	0.53
1:B:136:ARG:HH21	1:C:176:THR:HG21	1.74	0.52
1:A:109[B]:GLU:HG2	1:A:377:PHE:CG	2.45	0.51
1:A:109[B]:GLU:HG2	1:A:377:PHE:CD2	2.45	0.51
1:B:334:LYS:HG3	1:B:359:ASP:HB3	1.93	0.50
1:B:125:LEU:HD23	1:B:151[B]:GLU:HB2	1.94	0.50
1:C:142[B]:GLU:HG3	1:C:152:VAL:HG21	1.94	0.49
1:C:211:GLY:O	1:C:248:ARG:HD3	2.15	0.47



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash		
Atom-1	Atom-2	${f distance} ({f \AA})$	overlap (Å)       0.46       0.46       0.45       0.44       0.43       0.43       0.43       0.42       0.42       0.42       0.42       0.41       0.41       0.41       0.40		
1:B:93:PHE:O	1:B:259:LYS:NZ	2.45	0.46		
1:C:93:PHE:CZ	1:C:220[A]:LEU:HD11	2.51	0.46		
1:A:136:ARG:HG3	1:B:176:THR:HG21	1.98	0.46		
1:A:170:LEU:HD11	1:A:186:MET:SD	2.56	0.45		
1:A:142[B]:GLU:HG3	1:A:152:VAL:HG21	2.00	0.44		
1:B:265:ILE:O	1:B:290:TRP:HA	2.19	0.43		
1:A:342:LYS:CD	4:A:516:HOH:O	2.66	0.43		
1:B:211:GLY:O	1:B:248:ARG:HD3	2.19	0.43		
1:C:203:ILE:HB	1:C:204:PRO:HD2	2.00	0.43		
1:A:125:LEU:HD23	1:A:151[B]:GLU:HB2	1.99	0.42		
1:C:112:GLU:HG2	1:C:147[B]:ARG:NH1	2.34	0.42		
1:A:175:ARG:O	1:A:178:GLU:HG2	2.20	0.42		
1:B:142[B]:GLU:HG3	1:B:152:VAL:HG21	2.01	0.42		
1:A:342:LYS:HA	1:A:342:LYS:HD2	1.87	0.41		
1:C:342:LYS:HA	1:C:342:LYS:HD3	1.82	0.41		
1:C:313:ARG:HB2	1:C:323:TYR:CE1	2.55	0.41		
1:B:350:LEU:HB3	1:B:351:PRO:HA	2.03	0.41		
1:C:175[A]:ARG:HE	1:C:175[A]:ARG:HB2	1.68	0.40		
1:C:323:TYR:CE2	1:C:327:ARG:HD2	2.55	0.40		
1:B:203:ILE:HB	1:B:204:PRO:HD2	2.02	0.40		
1:A:136:ARG:HG3	1:B:176:THR:CG2	2.51	0.40		

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Favoured Allowed Out		Outliers	Perce	entiles
1	A	$331/332 \ (100\%)$	322 (97%)	9 (3%)	0	100	100
1	В	$328/332 \ (99\%)$	320 (98%)	8 (2%)	0	100	100
1	С	325/332~(98%)	316 (97%)	9 (3%)	0	100	100
All	All	984/996 (99%)	958 (97%)	26 (3%)	0	100	100



There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	$280/279 \; (100\%)$	277 (99%)	3 (1%)	73	61	
1	В	276/279 (99%)	271 (98%)	5 (2%)	59	40	
1	С	275/279 (99%)	269 (98%)	6 (2%)	52	32	
All	All	831/837 (99%)	817 (98%)	14 (2%)	67	43	

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	342	LYS
1	A	352	ARG
1	A	368	TYR
1	В	115	SER
1	В	119[A]	GLU
1	В	119[B]	GLU
1	В	352	ARG
1	В	368	TYR
1	С	80	HIS
1	С	175[A]	ARG
1	С	175[B]	ARG
1	С	285[A]	SER
1	С	285[B]	SER
1	С	352	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	В	ond leng	$\operatorname{gths}$	В	ond ang	les
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	$\mid \# Z  > 2 \mid$
3	GOL	A	402	-	5,5,5	1.63	1 (20%)	5,5,5	1.50	1 (20%)
2	PAL	В	401	-	9,15,15	1.88	3 (33%)	11,21,21	1.44	2 (18%)
2	PAL	С	401	-	9,15,15	2.06	3 (33%)	11,21,21	1.49	2 (18%)
2	PAL	A	401	-	9,15,15	2.04	1 (11%)	11,21,21	1.24	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	402	_	-	3/4/4/4	-
2	PAL	В	401	_	-	0/11/17/17	-
2	PAL	С	401	-	-	0/11/17/17	-
2	PAL	A	401	-	-	0/11/17/17	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\text{\AA})$
2	Α	401	PAL	C1-N2	4.59	1.43	1.34



$\alpha \cdots$	· ·	•	
Continued	trom	nromanne	naae
-	110111	picolous	payc

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	${ m Observed}({ m \AA})$	$\operatorname{Ideal}( ext{\AA})$
2	В	401	PAL	C1-N2	4.20	1.43	1.34
2	С	401	PAL	C1-N2	4.16	1.42	1.34
3	A	402	GOL	O2-C2	-2.89	1.34	1.43
2	С	401	PAL	P-C1P	2.60	1.83	1.79
2	В	401	PAL	P-O3P	-2.50	1.49	1.54
2	С	401	PAL	C2-N2	-2.24	1.43	1.46
2	В	401	PAL	O1-C1	-2.06	1.19	1.23

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^o)$
2	С	401	PAL	O1P-P-C1P	-2.98	104.15	110.94
2	A	401	PAL	O2P-P-C1P	2.60	112.30	106.84
3	A	402	GOL	C3-C2-C1	-2.32	102.69	111.70
2	A	401	PAL	O1P-P-C1P	-2.30	105.70	110.94
2	В	401	PAL	C3-C2-N2	-2.23	104.96	109.01
2	С	401	PAL	C3-C2-N2	-2.22	104.97	109.01
2	В	401	PAL	O3P-P-C1P	2.10	111.24	106.84

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	402	GOL	O1-C1-C2-C3
3	A	402	GOL	O1-C1-C2-O2
3	A	402	GOL	O2-C2-C3-O3


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	$egin{array}{c c} Analysed & <& RSRZ> & \#RSRZ>2 & \end{array}$		$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9		
1	A	315/332~(94%)	-0.15	7 (2%) 62	2 65	16, 22, 42, 67	0
1	В	311/332 (93%)	-0.10	11 (3%) 4	4 47	16, 21, 43, 71	0
1	С	311/332 (93%)	0.15	12 (3%) 3	9 42	19, 27, 46, 67	0
All	All	937/996 (94%)	-0.03	30 (3%) 4	7 50	16, 23, 44, 71	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	В	116	SER	5.2	
1	A	76	PHE	4.9	
1	В	115	SER	4.8	
1	С	116	SER	3.8	
1	В	320	LEU	3.7	
1	A	115	SER	3.6	
1	В	318	GLU	3.4	
1	С	81[A]	MET	2.9	
1	С	80	HIS	2.9	
1	С	339	VAL	2.7	
1	С	115	SER	2.7	
1	В	317	GLY	2.7	
1	В	319	ARG	2.6	
1	С	288	VAL	2.6	
1	A	154[A]	THR	2.6	
1	A	113	LYS	2.6	
1	С	284	THR	2.5	
1	В	321	ASP	2.5	
1	В	322	LEU	2.5	
1	С	283	LEU	2.5	
1	A	116	SER	2.5	
1	В	117	GLN	2.4	
1	С	343	LYS	2.3	



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Res	Type	RSRZ
1	В	114	SER	2.1
1	В	80	HIS	2.1
1	С	117	GLN	2.1
1	С	113	LYS	2.1
1	A	114	SER	2.1
1	С	139	LEU	2.0
1	A	153	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	GOL	A	402	6/6	0.94	0.12	24,31,39,47	0
2	PAL	С	401	16/16	0.97	0.12	18,21,25,27	0
2	PAL	A	401	16/16	0.98	0.09	16,19,23,23	0
2	PAL	В	401	16/16	0.98	0.10	14,18,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














## 6.5 Other polymers (i)

There are no such residues in this entry.

