



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 17, 2023 – 04:39 PM EDT

PDB ID : 4Z3T
Title : Meningococcal Factor H binding protein mutant L130R/G133D
Authors : Konar, M.; Beernink, P.T.
Deposited on : 2015-03-31
Resolution : 1.62 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4693 (1.64-1.60)
Clashscore	141614	5002 (1.64-1.60)
Ramachandran outliers	138981	4888 (1.64-1.60)
Sidechain outliers	138945	4887 (1.64-1.60)
RSRZ outliers	127900	4609 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	 91% 7%
1	B	267	 83% 6% 12% 3%

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4230 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Factor H binding protein variant A10_001.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	249	1984	1237	351	396	0	12	0
1	B	236	1867	1164	333	370	0	8	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	MET	-	initiating methionine	UNP Q6VS32
A	-3	GLY	-	expression tag	UNP Q6VS32
A	-2	PRO	-	expression tag	UNP Q6VS32
A	-1	ASP	-	expression tag	UNP Q6VS32
A	0	SER	-	expression tag	UNP Q6VS32
A	1	ASP	-	expression tag	UNP Q6VS32
A	2	ARG	-	expression tag	UNP Q6VS32
A	3	LEU	-	expression tag	UNP Q6VS32
A	4	GLN	-	expression tag	UNP Q6VS32
A	5	GLN	-	expression tag	UNP Q6VS32
A	6	ARG	-	expression tag	UNP Q6VS32
A	7	ARG	-	expression tag	UNP Q6VS32
A	130	ARG	LEU	engineered mutation	UNP Q6VS32
A	133	ASP	GLY	engineered mutation	UNP Q6VS32
A	255	LEU	-	expression tag	UNP Q6VS32
A	256	GLU	-	expression tag	UNP Q6VS32
A	257	HIS	-	expression tag	UNP Q6VS32
A	258	HIS	-	expression tag	UNP Q6VS32
A	259	HIS	-	expression tag	UNP Q6VS32
A	260	HIS	-	expression tag	UNP Q6VS32
A	261	HIS	-	expression tag	UNP Q6VS32
A	262	HIS	-	expression tag	UNP Q6VS32
B	-4	MET	-	initiating methionine	UNP Q6VS32
B	-3	GLY	-	expression tag	UNP Q6VS32
B	-2	PRO	-	expression tag	UNP Q6VS32

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	ASP	-	expression tag	UNP Q6VS32
B	0	SER	-	expression tag	UNP Q6VS32
B	1	ASP	-	expression tag	UNP Q6VS32
B	2	ARG	-	expression tag	UNP Q6VS32
B	3	LEU	-	expression tag	UNP Q6VS32
B	4	GLN	-	expression tag	UNP Q6VS32
B	5	GLN	-	expression tag	UNP Q6VS32
B	6	ARG	-	expression tag	UNP Q6VS32
B	7	ARG	-	expression tag	UNP Q6VS32
B	130	ARG	LEU	engineered mutation	UNP Q6VS32
B	133	ASP	GLY	engineered mutation	UNP Q6VS32
B	255	LEU	-	expression tag	UNP Q6VS32
B	256	GLU	-	expression tag	UNP Q6VS32
B	257	HIS	-	expression tag	UNP Q6VS32
B	258	HIS	-	expression tag	UNP Q6VS32
B	259	HIS	-	expression tag	UNP Q6VS32
B	260	HIS	-	expression tag	UNP Q6VS32
B	261	HIS	-	expression tag	UNP Q6VS32
B	262	HIS	-	expression tag	UNP Q6VS32

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mg 2 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	226	Total O 226 226	0	0
3	B	151	Total O 151 151	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Factor H binding protein variant A10_001

Chain A:  91% 7%



- Molecule 1: Factor H binding protein variant A10_001

Chain B:  83% 6% 12% 3%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.55Å 35.67Å 90.41Å 90.00° 104.11° 90.00°	Depositor
Resolution (Å)	66.03 – 1.62 66.03 – 1.62	Depositor EDS
% Data completeness (in resolution range)	99.3 (66.03-1.62) 99.2 (66.03-1.62)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 1.62Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.194 , 0.231 0.195 , 0.231	Depositor DCC
R_{free} test set	3005 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	15.7	Xtrriage
Anisotropy	0.064	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4230	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/2049	0.56	0/2752
1	B	0.32	0/1918	0.53	0/2575
All	All	0.34	0/3967	0.54	0/5327

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1984	0	1988	6	0
1	B	1867	0	1863	11	0
2	A	2	0	0	0	0
3	A	226	0	0	3	0
3	B	151	0	0	3	0
All	All	4230	0	3851	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 17 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:242:ARG:NH2	3:B:301:HOH:O	2.11	0.84
1:B:94:GLY:HA2	1:B:112[B]:GLU:HG2	1.62	0.82
1:A:37[B]:ASP:OD1	3:A:401:HOH:O	2.12	0.66
1:B:212[A]:ARG:NH2	1:B:217:GLU:OE2	2.29	0.65
1:A:166[A]:HIS:ND1	3:A:404:HOH:O	2.31	0.60

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/267 (97%)	250 (96%)	9 (4%)	0	100	100
1	B	240/267 (90%)	234 (98%)	6 (2%)	0	100	100
All	All	499/534 (93%)	484 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/220 (98%)	213 (99%)	2 (1%)	78	64
1	B	200/220 (91%)	199 (100%)	1 (0%)	88	80
All	All	415/440 (94%)	412 (99%)	3 (1%)	88	72

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	166[A]	HIS
1	A	166[B]	HIS
1	B	175	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/267 (93%)	-0.09	3 (1%) 79 78	8, 15, 31, 49	0
1	B	236/267 (88%)	-0.01	7 (2%) 50 47	11, 19, 37, 55	0
All	All	485/534 (90%)	-0.05	10 (2%) 63 62	8, 17, 35, 55	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	23	PRO	4.0
1	A	255	LEU	3.9
1	B	86	GLY	3.0
1	B	85	ASP	2.5
1	B	43	ASN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	A	302	1/1	0.92	0.17	28,28,28,28	0
2	MG	A	301	1/1	0.99	0.08	12,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.