

Full wwPDB X-ray Structure Validation Report (i)

Sep 18, 2021 - 08:00 am BST

PDB ID	:	6Z5Y
Title	:	Structure of a novel LPMO from Phytophthora infestans
Authors	:	Urresti, S.; Davies, G.J.
Deposited on		
Resolution	:	1.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Wiethe	$(\# \mathbf{Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
Clashscore	141614	1253 (1.08-0.96)
Ramachandran outliers	138981	1178 (1.08-0.96)
Sidechain outliers	138945	1180 (1.08-0.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain	
1	А	180	89%	8% ••
1	В	180	86%	13% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PGE	А	202	-	-	Х	-



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3216 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	177	Total	С	Ν	Ο	S	0	1.4	0
		111	1415	904	230	268	13	0	14	0
1	р	177	Total	С	Ν	Ο	S	0	19	0
	D	111	1441	919	235	275	12	0	19	0

• Molecule 1 is a protein called Lytic Polysaccharide Monooxygenase.

Chain	Residue	Modelled	Actual	Comment	Reference
A	173	TRP	-	expression tag	UNP D0N2F7
A	174	SER	-	expression tag	UNP D0N2F7
A	175	HIS	-	expression tag	UNP D0N2F7
A	176	PRO	-	expression tag	UNP D0N2F7
A	177	GLN	-	expression tag	UNP D0N2F7
A	178	PHE	-	expression tag	UNP D0N2F7
A	179	GLU	-	expression tag	UNP D0N2F7
A	180	LYS	-	expression tag	UNP D0N2F7
В	173	TRP	-	expression tag	UNP D0N2F7
В	174	SER	-	expression tag	UNP D0N2F7
В	175	HIS	-	expression tag	UNP D0N2F7
В	176	PRO	-	expression tag	UNP D0N2F7
В	177	GLN	-	expression tag	UNP D0N2F7
В	178	PHE	-	expression tag	UNP D0N2F7
В	179	GLU	-	expression tag	UNP D0N2F7
В	180	LYS	-	expression tag	UNP D0N2F7

There are 16 discrepancies between the modelled and reference sequences:

• Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Cu 1 1	0	0


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	TotalCu11	0	0

• Molecule 3 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	А	1	Total C 7 4	O 3	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	183	Total O 183 183	0	0
4	В	168	Total O 168 168	0	0





3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Lytic Polysaccharide Monooxygenase







4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants	36.29Å 45.46 Å 53.69 Å	Depositor
a, b, c, α , β , γ	83.91° 71.85° 83.78°	Depositor
Resolution (Å)	28.79 - 1.01	Depositor
% Data completeness	92.3 (28.79-1.01)	Depositor
(in resolution range)	52.5 (20.75-1.01)	Depositor
R_{merge}	0.05	Depositor
R _{sym}	(Not available)	Depositor
Refinement program	REFMAC $5.8.0258$	Depositor
R, R_{free}	0.143 , 0.164	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3216	wwPDB-VP
Average B, all atoms $(Å^2)$	13.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain		nd lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.67	2/1489~(0.1%)	0.89	2/2026~(0.1%)	
1	В	0.72	1/1509~(0.1%)	0.89	1/2057~(0.0%)	
All	All	0.69	3/2998~(0.1%)	0.89	3/4083~(0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	В	34	SER	CA-CB	-5.82	1.44	1.52
1	А	14[A]	SER	C-O	5.02	1.32	1.23
1	А	14[B]	SER	C-O	5.02	1.32	1.23

All (3) bond length outliers are listed below:

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\mathbf{Ideal}(^{o})$
1	А	24[A]	GLU	CB-CA-C	-5.26	99.88	110.40
1	А	24[B]	GLU	CB-CA-C	-5.26	99.88	110.40
1	В	54	LYS	CB-CA-C	5.16	120.73	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:



	Mol	Chain	Res	Type	Group
ſ	1	В	41	LEU	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1415	0	1346	19	0
1	В	1441	0	1357	19	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0
3	А	7	0	9	6	0
4	А	183	0	0	10	0
4	В	168	0	0	1	0
All	All	3216	0	2712	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:47[B]:LEU:O	1:B:51:ASN:ND2	1.97	0.95
1:A:27:TRP:HE1	3:A:202:PGE:C5	1.85	0.90
1:B:29[B]:GLY:HA3	1:B:47[B]:LEU:HD11	1.56	0.86
1:A:6[A]:LYS:CE	4:A:302:HOH:O	2.20	0.83
1:A:6[A]:LYS:NZ	4:A:301:HOH:O	1.69	0.82
1:A:27:TRP:HE1	3:A:202:PGE:H5	1.45	0.81
1:B:28:LYS:O	1:B:47[B]:LEU:HD22	1.79	0.81
3:A:202:PGE:H4	4:A:307:HOH:O	1.86	0.74
1:A:6[A]:LYS:NZ	4:A:302:HOH:O	1.83	0.74
1:A:6[A]:LYS:HE2	4:A:302:HOH:O	1.81	0.73
1:A:27:TRP:HE1	3:A:202:PGE:H52	1.54	0.72
1:A:24[B]:GLU:H	1:A:24[B]:GLU:CD	1.94	0.71
1:B:104:LEU:HD13	1:B:138[A]:CYS:SG	2.35	0.66
1:B:28:LYS:O	1:B:47[B]:LEU:CD2	2.43	0.66
1:B:30[A]:GLY:HA2	1:B:33:GLU:OE1	1.96	0.65
1:A:12[A]:LYS:HE3	1:A:73:THR:O	1.97	0.65

Continued on next page...



Continued from previ	1.0	Interatomic	Clash
Atom-1	Atom-2	distance $(Å)$	overlap (Å)
1:A:24[B]:GLU:OE2	4:A:303:HOH:O	2.17	0.58
1:A:26:GLN:OE1	4:A:304:HOH:O	2.18	0.57
1:B:29[B]:GLY:CA	1:B:47[B]:LEU:HD11	2.34	0.55
1:A:27:TRP:NE1	3:A:202:PGE:H52	2.20	0.55
1:B:115[B]:CYS:SG	1:B:130:PHE:HE1	2.30	0.54
1:B:63:ASN:OD1	1:B:65[B]:VAL:HG13	2.07	0.53
1:A:59:LEU:HD11	3:A:202:PGE:H32	1.92	0.52
1:B:138[B]:CYS:SG	1:B:142:GLY:HA2	2.50	0.51
1:B:14[B]:SER:HB2	4:B:372:HOH:O	2.12	0.50
1:B:29[B]:GLY:HA3	1:B:47[B]:LEU:HD21	1.94	0.48
1:B:138[B]:CYS:SG	1:B:142:GLY:C	2.93	0.47
1:A:12[B]:LYS:HE3	1:A:73:THR:O	2.15	0.47
1:A:115[B]:CYS:SG	1:A:130:PHE:HE1	2.38	0.47
1:B:48[B]:ALA:O	1:B:51:ASN:N	2.43	0.46
1:B:142:GLY:O	1:B:143:CYS:HB3	2.16	0.45
1:A:138[B]:CYS:SG	1:A:143:CYS:N	2.90	0.45
1:A:6[A]:LYS:HE2	4:A:447:HOH:O	2.18	0.44
1:B:48[A]:ALA:HB3	1:B:49[A]:PRO:HD3	1.99	0.44
1:B:30[A]:GLY:CA	1:B:33:GLU:OE1	2.63	0.43
1:B:39:GLY:O	1:B:42[B]:ALA:HB3	2.18	0.43
1:A:63:ASN:ND2	4:A:307:HOH:O	2.52	0.42
1:B:29[B]:GLY:HA3	1:B:47[B]:LEU:CD1	2.40	0.41
1:A:12[A]:LYS:HE2	4:A:424:HOH:O	2.19	0.41

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	189/180~(105%)	184 (97%)	5(3%)	0	100	100
1	В	194/180~(108%)	188 (97%)	6 (3%)	0	100	100

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	383/360~(106%)	372~(97%)	11 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	151/149~(101%)	148~(98%)	3 (2%)	55 21
1	В	151/149~(101%)	150~(99%)	1 (1%)	84 58
All	All	302/298~(101%)	298~(99%)	4 (1%)	65 35

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	90	PHE
1	А	111	GLN
1	А	131	LYS
1	В	111	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type	
1	В	177	GLN	

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Chain R		Res	Link	Bond lengths			Bond angles			
	туре	Cham	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PGE	А	202	-	$6,\!6,\!9$	0.35	0	$5,\!5,\!8$	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	А	202	-	-	1/4/4/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	202	PGE	O2-C3-C4-O3

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	202	PGE	6	0



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

