



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 24, 2021 – 06:14 pm GMT

PDB ID : 6Z8S
Title : Copper transporter OprC
Authors : Bhamidimarri, S.P.; van den Berg, B.
Deposited on : 2020-06-02
Resolution : 2.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5509 (2.40-2.36)
Clashscore	141614	6082 (2.40-2.36)
Ramachandran outliers	138981	5973 (2.40-2.36)
Sidechain outliers	138945	5975 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	723	
1	B	723	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10273 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative copper transport outer membrane porin OprC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	655	5081	3188	899	977	17	0	0	0
1	B	655	5083	3190	899	976	18	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	325	HIS	MET	engineered mutation	UNP G3XD89
B	325	HIS	MET	engineered mutation	UNP G3XD89

- Molecule 2 is COPPER (I) ION (three-letter code: CU1) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cu	0	0
			1	1		
2	B	1	Total	Cu	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	72	Total	O	0	0
			72	72		
3	B	35	Total	O	0	0
			35	35		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative copper transport outer membrane porin OprC



- Molecule 1: Putative copper transport outer membrane porin OprC



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	156.10Å 195.69Å 166.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.02 – 2.37 97.84 – 2.37	Depositor EDS
% Data completeness (in resolution range)	99.7 (61.02-2.37) 99.9 (97.84-2.37)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 2.37Å)	Xtriage
Refinement program	PHENIX 1.18_3855	Depositor
R, R_{free}	0.219 , 0.252 0.219 , 0.255	Depositor DCC
R_{free} test set	5166 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	62.8	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10273	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/5208	0.65	2/7060 (0.0%)
1	B	0.44	0/5210	0.64	2/7062 (0.0%)
All	All	0.45	0/10418	0.64	4/14122 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	555	ARG	CG-CD-NE	8.74	130.15	111.80
1	A	236	ASP	CB-CG-OD1	6.00	123.70	118.30
1	B	555	ARG	NE-CZ-NH2	5.31	122.95	120.30
1	A	348	ASP	CB-CG-OD1	5.23	123.01	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5081	0	4854	33	0
1	B	5083	0	4858	59	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	72	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	35	0	0	0	0
All	All	10273	0	9712	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:671:THR:HG22	1:B:673:ASN:H	1.26	0.96
1:B:151:THR:HA	1:B:154:ILE:HD12	1.59	0.85
1:A:117:ASN:OD1	1:A:251:LYS:NZ	2.14	0.80
1:A:555:ARG:HH21	1:A:556:GLU:HB2	1.48	0.78
1:B:94:PRO:O	1:B:209:ARG:NH2	2.19	0.76
1:A:95:ALA:O	1:A:97:ASP:N	2.22	0.72
1:A:72:THR:HG22	1:A:91:GLN:HB3	1.73	0.70
1:A:555:ARG:NH2	1:A:556:GLU:HB2	2.07	0.70
1:A:385:TYR:OH	1:A:390:ASP:OD1	2.08	0.67
1:B:82:ILE:HD11	1:B:166:LYS:HE2	1.78	0.66
1:B:286:MET:HE2	1:B:325:HIS:CD2	2.31	0.66
1:B:649:VAL:O	1:B:694:LYS:HE2	1.98	0.63
1:A:555:ARG:NE	1:A:556:GLU:H	1.98	0.62
1:B:193:GLU:N	1:B:193:GLU:OE1	2.34	0.60
1:B:264:GLU:H	1:B:264:GLU:CD	2.07	0.58
1:B:72:THR:HG22	1:B:91:GLN:HB2	1.87	0.57
1:B:300:PHE:HE1	1:B:302:LYS:HB2	1.70	0.57
1:B:625:ASP:HB2	1:B:669:ARG:HB2	1.87	0.56
1:A:334:ASP:HB3	1:A:337:SER:HB3	1.87	0.56
1:A:189:GLU:OE2	1:A:197:ARG:NH1	2.39	0.56
1:B:612:PRO:HB3	1:B:638:GLN:HB2	1.87	0.55
1:B:68:PRO:HB3	1:B:675:LYS:HD2	1.89	0.55
1:B:270:LEU:HD12	1:B:298:LEU:HD13	1.89	0.55
1:A:197:ARG:NH2	1:A:215:ASP:OD1	2.40	0.54
1:A:416:THR:HG23	1:A:426:ILE:HG13	1.89	0.54
1:B:448:MET:HG2	1:B:450:HIS:HB2	1.90	0.54
1:B:95:ALA:HA	1:B:209:ARG:NH2	2.22	0.54
1:A:255:ASP:OD2	1:A:273:GLY:HA3	2.09	0.53
1:B:385:TYR:OH	1:B:390:ASP:OD2	2.22	0.53
1:A:464:THR:HG23	1:B:547:GLN:OE1	2.09	0.53
1:B:79:PRO:O	1:B:81:THR:HG23	2.09	0.52
1:B:214:LEU:HD12	1:B:215:ASP:H	1.75	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:484:TYR:HH	1:B:540:SER:HG	1.51	0.52
1:A:79:PRO:O	1:A:81:THR:HG23	2.09	0.52
1:A:248:ARG:HD3	1:A:280:ARG:NH2	2.26	0.51
1:B:262:PRO:HD2	1:B:266:THR:HB	1.93	0.51
1:B:209:ARG:HA	1:B:233:GLN:O	2.11	0.50
1:B:286:MET:CE	1:B:325:HIS:CD2	2.95	0.50
1:B:209:ARG:H	1:B:234:SER:HA	1.77	0.49
1:A:96:SER:O	1:A:251:LYS:HE2	2.12	0.49
1:B:72:THR:HG22	1:B:91:GLN:CB	2.42	0.49
1:B:286:MET:HE3	1:B:326:ASP:CA	2.43	0.49
1:A:268:ILE:HG12	1:A:300:PHE:HD1	1.77	0.48
1:A:531:ASN:HD22	1:A:536:GLN:HE21	1.61	0.48
1:B:671:THR:HG22	1:B:673:ASN:N	2.10	0.48
1:B:484:TYR:OH	1:B:540:SER:OG	2.25	0.48
1:B:363:ASP:OD1	1:B:364:ASP:N	2.38	0.48
1:B:286:MET:CE	1:B:325:HIS:HD2	2.27	0.47
1:B:101:TYR:CD1	1:B:156:PRO:HG2	2.49	0.47
1:A:388:MET:HA	1:A:388:MET:CE	2.43	0.47
1:A:129:ARG:HD3	1:A:519:GLU:OE2	2.14	0.47
1:A:612:PRO:HB3	1:A:638:GLN:HB2	1.96	0.46
1:A:138:MET:CE	1:A:140:LEU:HD21	2.45	0.46
1:B:361:ARG:NH2	1:B:366:LYS:HE3	2.31	0.46
1:B:368:VAL:O	1:B:413:GLY:HA2	2.15	0.46
1:B:629:GLY:O	1:B:664:LEU:HD12	2.14	0.46
1:B:81:THR:HA	1:B:164:VAL:O	2.16	0.46
1:B:209:ARG:NH1	1:B:211:ASP:OD1	2.49	0.46
1:B:95:ALA:N	1:B:100:ASP:OD2	2.49	0.45
1:B:697:ASP:O	1:B:698:ALA:HB3	2.16	0.45
1:B:286:MET:HE3	1:B:326:ASP:HA	1.98	0.45
1:B:478:ASP:N	1:B:478:ASP:OD1	2.49	0.45
1:A:593:LEU:HD11	1:A:613:LEU:HD11	1.98	0.45
1:B:497:TRP:O	1:B:501:SER:HB2	2.17	0.44
1:B:524:LEU:O	1:B:542:TYR:HA	2.18	0.44
1:B:479:SER:HB2	1:B:481:THR:OG1	2.18	0.44
1:A:621:TYR:OH	1:A:623:GLU:OE2	2.25	0.44
1:A:307:ASP:OD2	1:A:307:ASP:N	2.42	0.44
1:A:329:ARG:NH2	3:A:913:HOH:O	2.51	0.44
1:B:214:LEU:HD12	1:B:215:ASP:N	2.32	0.44
1:A:151:THR:HA	1:A:154:ILE:HD12	1.99	0.43
1:B:91:GLN:OE1	1:B:665:ASN:ND2	2.30	0.43
1:B:696:GLY:HA3	1:B:702:PHE:CG	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:361:ARG:HE	1:B:361:ARG:HB2	1.68	0.43
1:B:203:LEU:HB3	1:B:211:ASP:HB2	2.01	0.43
1:B:325:HIS:NE2	1:B:345:SER:OG	2.52	0.43
1:A:591:ALA:HA	1:A:616:ARG:O	2.18	0.42
1:B:593:LEU:HD11	1:B:613:LEU:HD11	2.01	0.42
1:A:497:TRP:O	1:A:501:SER:HB2	2.19	0.42
1:A:72:THR:HG22	1:A:91:GLN:CB	2.45	0.41
1:A:203:LEU:HD12	1:A:718:LYS:HD2	2.02	0.41
1:B:501:SER:N	1:B:502:PRO:HD2	2.35	0.41
1:B:135:ASN:O	1:B:315:GLN:NE2	2.49	0.41
1:B:444:LYS:HE3	1:B:449:GLY:O	2.21	0.41
1:B:203:LEU:O	1:B:210:PHE:HA	2.21	0.41
1:B:246:PRO:HG2	1:B:700:PHE:CG	2.55	0.41
1:B:416:THR:HG23	1:B:426:ILE:HG13	2.04	0.40
1:B:506:PRO:HA	1:B:566:THR:HG22	2.04	0.40
1:A:384:LYS:HE3	1:A:398:PHE:CZ	2.57	0.40
1:A:286:MET:HE3	1:A:325:HIS:HD2	1.86	0.40
1:B:129:ARG:HH11	1:B:129:ARG:HD2	1.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	651/723 (90%)	625 (96%)	24 (4%)	2 (0%)	41	53
1	B	651/723 (90%)	618 (95%)	30 (5%)	3 (0%)	29	39
All	All	1302/1446 (90%)	1243 (96%)	54 (4%)	5 (0%)	34	46

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	96	SER
1	A	247	SER
1	B	209	ARG
1	B	247	SER
1	B	703	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	524/573 (91%)	501 (96%)	23 (4%)	28	42
1	B	524/573 (91%)	500 (95%)	24 (5%)	27	40
All	All	1048/1146 (91%)	1001 (96%)	47 (4%)	27	41

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	LEU
1	A	97	ASP
1	A	119	ASP
1	A	139	MET
1	A	286	MET
1	A	332	THR
1	A	341	MET
1	A	363	ASP
1	A	445	SER
1	A	450	HIS
1	A	464	THR
1	A	515	LYS
1	A	540	SER
1	A	555	ARG
1	A	558	MET
1	A	561	SER
1	A	562	SER
1	A	588	LYS
1	A	616	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	632	TRP
1	A	672	ARG
1	A	675	LYS
1	A	714	THR
1	B	78	SER
1	B	90	ARG
1	B	96	SER
1	B	97	ASP
1	B	155	SER
1	B	196	SER
1	B	198	VAL
1	B	203	LEU
1	B	209	ARG
1	B	270	LEU
1	B	302	LYS
1	B	307	ASP
1	B	329	ARG
1	B	341	MET
1	B	343	MET
1	B	478	ASP
1	B	479	SER
1	B	529	GLN
1	B	558	MET
1	B	561	SER
1	B	562	SER
1	B	632	TRP
1	B	656	SER
1	B	677	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	531	ASN
1	A	536	GLN
1	B	76	GLN
1	B	131	ASN
1	B	450	HIS

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.