



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 22, 2023 – 03:36 pm GMT

PDB ID : 7Z8D
Title : Structure of Photosynthetic Reaction Center From Rhodobacter Sphaeroides strain RV by fixed-target serial synchrotron crystallography (room temperature, 26keV)
Authors : Gabdulkhakov, A.G.; Selikhanov, G.K.; Guenther, S.; Petrova, T.; Meents, A.; Fufina, T.Y.; Vasilieva, L.G.
Deposited on : 2022-03-17
Resolution : 2.14 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.32.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	242	 5% 89% 11%
2	L	281	 7% 90% 10%
3	M	302	 5% 90% 10%

2 Entry composition [i](#)

There are 16 unique types of molecules in this entry. The entry contains 7660 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reaction center protein H chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	242	1875	1198	322	345	10	0	4	0

- Molecule 2 is a protein called Reaction center protein L chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	281	2430	1645	387	390	8	0	22	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	178	THR	SER	engineered mutation	UNP P0C0Y8


- Molecule 3 is a protein called Reaction center protein M chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	M	302	2464	1643	405	405	11	0	7	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
M	8	THR	SER	engineered mutation	UNP P0C0Y9

- Molecule 4 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula: C₁₄H₃₁NO).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	H	1	Total 16	C 14	N 1	O 1	0	0
4	M	1	Total 16	C 14	N 1	O 1	0	0
4	M	1	Total 16	C 14	N 1	O 1	0	0
4	M	1	Total 16	C 14	N 1	O 1	0	0
4	M	1	Total 16	C 14	N 1	O 1	0	0

- Molecule 5 is BACTERIOCHLOROPHYLL A (three-letter code: BCL) (formula: C₅₅H₇₄MgN₄O₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Mg	N			O
5	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
5	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
5	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
5	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		

- Molecule 6 is BACTERIOPHEOPHYTIN A (three-letter code: BPH) (formula: $C_{55}H_{76}N_4O_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	L	1	Total	C	N	O	0	0
			65	55	4	6		
6	M	1	Total	C	N	O	0	0
			65	55	4	6		

- Molecule 7 is DODECANE (three-letter code: D12) (formula: $C_{12}H_{26}$).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	L	1	Total	C	0	0
			12	12		
7	M	1	Total	C	0	0
			12	12		

- Molecule 8 is PENTADECANE (three-letter code: MYS) (formula: $C_{15}H_{32}$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	L	1	Total C 15 15	0	0

- Molecule 9 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: $C_{21}H_{40}O_4$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	L	1	Total C O 25 21 4	0	0

- Molecule 10 is (2R)-2-hydroxy-3-(phosphonoxy)propyl (9E)-octadec-9-enoate (three-letter code: NKP) (formula: $C_{21}H_{41}O_7P$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
10	M	1	Total	C	O	P	0	0
			29	21	7	1		
10	M	1	Total	C	O	P	0	0
			29	21	7	1		

- Molecule 11 is UBIQUINONE-10 (three-letter code: U10) (formula: $C_{59}H_{90}O_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
11	M	1	Total	C	O		0	0
			63	59	4			

- Molecule 12 is FE (III) ION (three-letter code: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	M	1	Total Fe 1 1	0	0

- Molecule 13 is SPEROIDENONE (three-letter code: SPN) (formula: C₄₁H₇₀O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	M	1	Total C O 43 41 2	0	0

- Molecule 14 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
14	M	1	Total	C	O	0	0
			4	2	2		

- Molecule 15 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
15	M	1	Total	O	P	0	0
			5	4	1		


- Molecule 16 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	H	72	Total 72	O 72	0	0
16	L	53	Total 53	O 53	0	0
16	M	54	Total 54	O 54	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Reaction center protein H chain



- Molecule 2: Reaction center protein L chain



- Molecule 3: Reaction center protein M chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	102.50Å 102.50Å 237.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.05 – 2.14 47.05 – 2.14	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.05-2.14) 88.0 (47.05-2.14)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.56 (at 2.14Å)	Xtrriage
Refinement program	REFMAC 5.8.0258, PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.179 , 0.213 0.182 , 0.219	Depositor DCC
R_{free} test set	3550 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	31.0	Xtrriage
Anisotropy	0.015	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 80.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7660	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, FE, SPN, NKP, OLC, LDA, BCL, U10, D12, BPH, EDO, MYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.40	0/1927	0.62	0/2620
2	L	0.38	0/2531	0.52	0/3468
3	M	0.40	0/2556	0.54	0/3488
All	All	0.39	0/7014	0.55	0/9576

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1875	0	1876	17	0
2	L	2430	0	2366	16	0
3	M	2464	0	2381	22	0
4	H	16	0	31	1	0
4	M	64	0	124	3	0
5	L	132	0	148	3	0
5	M	132	0	148	4	0
6	L	65	0	76	1	0
6	M	65	0	76	3	0
7	L	12	0	26	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	M	12	0	26	0	0
8	L	15	0	32	0	0
9	L	25	0	40	1	0
10	M	58	0	78	3	0
11	M	63	0	90	2	0
12	M	1	0	0	0	0
13	M	43	0	70	8	0
14	M	4	0	6	0	0
15	M	5	0	0	0	0
16	H	72	0	0	0	0
16	L	53	0	0	0	0
16	M	54	0	0	0	0
All	All	7660	0	7594	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:M:108:PRO:HG2	3:M:111:GLU:HB2	1.65	0.79
3:M:2:GLU:O	3:M:4:GLN:NE2	2.24	0.66
1:H:197:LYS:HZ1	3:M:1:ALA:C	2.04	0.61
3:M:60:LEU:HD23	6:M:407:BPH:H4C1	1.84	0.59
2:L:229:ILE:HB	9:L:306:OLC:H21	1.87	0.57

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	244/242 (101%)	242 (99%)	2 (1%)	0	100	100
2	L	300/281 (107%)	286 (95%)	14 (5%)	0	100	100
3	M	307/302 (102%)	296 (96%)	9 (3%)	2 (1%)	22	14
All	All	851/825 (103%)	824 (97%)	25 (3%)	2 (0%)	47	45

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	M	2	GLU
3	M	195	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	200/197 (102%)	194 (97%)	6 (3%)	41	39
2	L	238/220 (108%)	226 (95%)	12 (5%)	24	20
3	M	242/236 (102%)	239 (99%)	3 (1%)	71	74
All	All	680/653 (104%)	659 (97%)	21 (3%)	50	38

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	269[A]	LEU
2	L	272[B]	TRP
3	M	216	PHE
3	M	52[A]	LEU
2	L	272[A]	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 1 is monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	BPH	L	303	-	51,70,70	0.92	2 (3%)	52,101,101	1.13	4 (7%)
7	D12	L	304	-	11,11,11	0.30	0	10,10,10	0.45	0
10	NKP	M	402	-	28,28,28	0.35	0	31,32,32	0.43	0
14	EDO	M	414	-	3,3,3	0.41	0	2,2,2	0.80	0
10	NKP	M	401	-	28,28,28	0.37	0	31,32,32	0.52	0
6	BPH	M	407	-	51,70,70	1.00	3 (5%)	52,101,101	1.35	7 (13%)
8	MYS	L	305	-	14,14,14	0.29	0	13,13,13	0.50	0
7	D12	M	413	-	11,11,11	0.26	0	10,10,10	0.66	0
4	LDA	M	403	-	12,15,15	2.06	1 (8%)	14,17,17	0.69	0
5	BCL	M	405	-	58,74,74	1.23	4 (6%)	69,115,115	1.51	13 (18%)
5	BCL	M	406	-	58,74,74	1.32	5 (8%)	69,115,115	1.49	11 (15%)
13	SPN	M	409	-	40,42,42	0.54	0	50,52,52	1.60	10 (20%)
15	PO4	M	415	-	4,4,4	0.88	0	6,6,6	0.37	0
9	OLC	L	306	-	24,24,24	0.93	1 (4%)	25,25,25	0.87	0
4	LDA	M	412	-	12,15,15	2.05	1 (8%)	14,17,17	0.61	0
4	LDA	M	410	-	12,15,15	2.02	1 (8%)	14,17,17	0.41	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BCL	L	302	-	58,74,74	1.24	4 (6%)	69,115,115	1.25	8 (11%)
5	BCL	L	301	-	58,74,74	1.44	5 (8%)	69,115,115	1.42	12 (17%)
4	LDA	M	411	-	12,15,15	2.06	1 (8%)	14,17,17	0.50	0
4	LDA	H	301	-	12,15,15	1.93	1 (8%)	14,17,17	0.65	0
11	U10	M	404	-	63,63,63	2.64	17 (26%)	76,79,79	1.78	19 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BPH	L	303	-	-	4/37/105/105	0/5/6/6
7	D12	L	304	-	-	4/9/9/9	-
10	NKP	M	402	-	-	8/28/28/28	-
14	EDO	M	414	-	-	1/1/1/1	-
10	NKP	M	401	-	-	9/28/28/28	-
6	BPH	M	407	-	-	8/37/105/105	0/5/6/6
8	MYS	L	305	-	-	6/12/12/12	-
7	D12	M	413	-	-	2/9/9/9	-
4	LDA	M	403	-	-	11/13/13/13	-
5	BCL	M	405	-	-	0/37/137/137	-
5	BCL	M	406	-	-	0/37/137/137	-
13	SPN	M	409	-	-	18/50/51/51	-
9	OLC	L	306	-	-	8/24/24/24	-
4	LDA	M	412	-	-	7/13/13/13	-
4	LDA	M	410	-	-	5/13/13/13	-
5	BCL	L	302	-	-	2/37/137/137	-
5	BCL	L	301	-	-	5/37/137/137	-
4	LDA	M	411	-	-	8/13/13/13	-
4	LDA	H	301	-	-	4/13/13/13	-
11	U10	M	404	-	-	16/63/87/87	0/1/1/1

The worst 5 of 46 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	M	403	LDA	O1-N1	-7.06	1.25	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	M	411	LDA	O1-N1	-7.04	1.25	1.42
4	M	412	LDA	O1-N1	-7.03	1.25	1.42
4	M	410	LDA	O1-N1	-6.88	1.26	1.42
4	H	301	LDA	O1-N1	-6.63	1.26	1.42

The worst 5 of 84 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	M	405	BCL	CMB-C2B-C1B	-4.45	121.63	128.46
5	M	406	BCL	C4A-NA-C1A	4.27	108.62	106.71
5	M	406	BCL	CMB-C2B-C1B	-4.10	122.16	128.46
11	M	404	U10	C22-C23-C24	-4.09	117.81	127.66
13	M	409	SPN	CM5-C13-C14	4.08	122.14	115.27

There are no chirality outliers.

5 of 126 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	M	403	LDA	C2-C1-N1-O1
4	M	403	LDA	C2-C1-N1-CM2
4	M	410	LDA	C2-C1-N1-CM2
4	M	411	LDA	C2-C1-N1-O1
4	M	411	LDA	C2-C1-N1-CM1

There are no ring outliers.

14 monomers are involved in 25 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	L	303	BPH	1	0
10	M	402	NKP	2	0
10	M	401	NKP	1	0
6	M	407	BPH	3	0
4	M	403	LDA	1	0
5	M	405	BCL	3	0
5	M	406	BCL	1	0
13	M	409	SPN	8	0
9	L	306	OLC	1	0
4	M	410	LDA	2	0
5	L	302	BCL	1	0
5	L	301	BCL	2	0
4	H	301	LDA	1	0




Continued on next page...




Continued from previous page...




Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	M	404	U10	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	242/242 (100%)	-0.11	11 (4%) 33 40	25, 40, 63, 114	0
2	L	281/281 (100%)	0.02	20 (7%) 16 20	23, 37, 60, 117	0
3	M	302/302 (100%)	0.02	14 (4%) 32 40	22, 36, 63, 111	0
All	All	825/825 (100%)	-0.02	45 (5%) 25 31	22, 38, 63, 117	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	M	1	ALA	13.3
2	L	59	TRP	9.9
1	H	250	SER	8.5
2	L	277[A]	GLY	6.7
2	L	51	TRP	6.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	LDA	M	403	16/16	0.62	0.26	58,71,83,84	0
4	LDA	M	410	16/16	0.69	0.24	47,63,77,79	0
4	LDA	M	412	16/16	0.69	0.40	66,78,108,108	0
4	LDA	M	411	16/16	0.74	0.25	66,74,88,89	0
10	NKP	M	402	29/29	0.75	0.23	45,62,79,87	0
8	MYS	L	305	15/15	0.76	0.23	49,66,77,82	0
13	SPN	M	409	43/43	0.77	0.25	36,57,78,82	0
9	OLC	L	306	25/25	0.80	0.21	32,55,74,82	0
7	D12	L	304	12/12	0.82	0.14	50,57,62,63	0
11	U10	M	404	63/63	0.86	0.20	22,52,77,82	0
4	LDA	H	301	16/16	0.88	0.15	36,43,65,77	0
7	D12	M	413	12/12	0.90	0.17	47,64,80,83	0
14	EDO	M	414	4/4	0.91	0.13	48,55,61,68	0
10	NKP	M	401	29/29	0.92	0.18	37,69,84,88	0
15	PO4	M	415	5/5	0.92	0.20	58,73,86,90	0
5	BCL	L	301	66/66	0.94	0.11	23,33,46,67	0
6	BPH	L	303	65/65	0.95	0.16	21,29,44,47	0
6	BPH	M	407	65/65	0.95	0.13	24,32,102,108	0
5	BCL	M	405	66/66	0.95	0.14	26,35,82,89	0
5	BCL	M	406	66/66	0.95	0.12	17,32,55,91	0
5	BCL	L	302	66/66	0.96	0.14	22,33,58,68	0
12	FE	M	408	1/1	0.99	0.12	24,24,24,24	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NKP M 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around SPN M 409:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC L 306:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around U10 M 404:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NKP M 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BCL L 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BPH L 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BPH M 407:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around BCL M 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers [i](#)

There are no such residues in this entry.