



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2023 – 04:02 AM EDT

PDB ID : 1ZB7  
Title : Crystal Structure of Botulinum Neurotoxin Type G Light Chain  
Authors : Arndt, J.W.; Yu, W.; Bi, F.; Stevens, R.C.  
Deposited on : 2005-04-07  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	455	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3535 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called neurotoxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	414	3324	2137	544	626	17	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	444	ALA	-	cloning artifact	UNP Q60393
A	445	PRO	-	cloning artifact	UNP Q60393
A	446	PRO	-	cloning artifact	UNP Q60393
A	447	THR	-	cloning artifact	UNP Q60393
A	448	PRO	-	cloning artifact	UNP Q60393
A	449	GLY	-	cloning artifact	UNP Q60393
A	450	HIS	-	cloning artifact	UNP Q60393
A	451	HIS	-	cloning artifact	UNP Q60393
A	452	HIS	-	cloning artifact	UNP Q60393
A	453	HIS	-	cloning artifact	UNP Q60393
A	454	HIS	-	cloning artifact	UNP Q60393
A	455	HIS	-	cloning artifact	UNP Q60393

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is CITRATE ANION (three-letter code: FLC) (formula: C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	197	Total	O	0	0
			197	197		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: neurotoxin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	178.87Å 178.87Å 80.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.81 – 2.35 24.80 – 2.35	Depositor EDS
% Data completeness (in resolution range)	94.0 (24.81-2.35) 94.0 (24.80-2.35)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.83 (at 2.36Å)	Xtrriage
Refinement program	REFMAC 5.2.0011	Depositor
R, $R_{free}$	0.174 , 0.222 0.215 , 0.258	Depositor DCC
$R_{free}$ test set	1536 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	43.2	Xtrriage
Anisotropy	0.062	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3535	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.26	13/3408 (0.4%)	1.08	10/4611 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	84	GLU	CD-OE2	7.89	1.34	1.25
1	A	323	TYR	CD1-CE1	-7.45	1.28	1.39
1	A	169	ASP	CB-CG	6.78	1.66	1.51
1	A	306	VAL	CB-CG2	-6.66	1.38	1.52
1	A	36	ALA	CA-CB	6.54	1.66	1.52
1	A	44	TRP	CG-CD1	6.50	1.45	1.36
1	A	335	GLY	N-CA	-6.23	1.36	1.46
1	A	306	VAL	CA-CB	6.08	1.67	1.54
1	A	363	GLU	CG-CD	5.83	1.60	1.51
1	A	325	ASN	CB-CG	5.65	1.64	1.51
1	A	42	ARG	CZ-NH1	5.58	1.40	1.33
1	A	307	SER	CA-CB	-5.54	1.44	1.52
1	A	129	PHE	N-CA	-5.39	1.35	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	42	ARG	NE-CZ-NH1	13.94	127.27	120.30
1	A	42	ARG	NE-CZ-NH2	-12.84	113.88	120.30
1	A	302	ARG	NE-CZ-NH2	-11.34	114.63	120.30
1	A	302	ARG	NE-CZ-NH1	10.25	125.42	120.30
1	A	169	ASP	CB-CG-OD1	8.69	126.12	118.30
1	A	298	ASP	CB-CG-OD1	7.01	124.61	118.30
1	A	332	ASP	CB-CG-OD1	6.13	123.82	118.30
1	A	41	ASP	CB-CG-OD2	-5.99	112.91	118.30
1	A	12	ASP	CB-CG-OD1	5.43	123.19	118.30
1	A	285	ASP	CB-CG-OD2	-5.01	113.79	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	26	ASN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3324	0	3261	44	0
2	A	1	0	0	0	0
3	A	13	0	5	0	0
4	A	197	0	0	6	0
All	All	3535	0	3266	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:ARG:HD2	4:A:620:HOH:O	1.58	1.03
1:A:218:ARG:CD	4:A:620:HOH:O	2.26	0.71
1:A:251:PRO:HG3	1:A:257:PHE:O	1.95	0.67

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:ASN:O	1:A:132:ASN:ND2	2.29	0.65
1:A:413:LYS:NZ	1:A:422:GLU:OE2	2.30	0.64
1:A:177:MET:HG3	1:A:180:HIS:CE1	2.36	0.61
1:A:258:MET:HG2	4:A:683:HOH:O	2.01	0.60
1:A:208:GLU:HA	1:A:208:GLU:OE1	2.02	0.59
1:A:42:ARG:HH22	1:A:155:MET:HE2	1.66	0.59
1:A:406:THR:HB	4:A:680:HOH:O	2.04	0.57
1:A:23:GLU:OE2	1:A:25:PHE:O	2.22	0.56
1:A:176:ILE:HG22	4:A:693:HOH:O	2.05	0.56
1:A:190:ARG:O	1:A:191:MET:HE3	2.06	0.56
1:A:432:ARG:HH11	1:A:434:ALA:HB2	1.71	0.55
1:A:208:GLU:O	1:A:210:LYS:N	2.41	0.54
1:A:230:HIS:ND1	1:A:357:GLU:OE1	2.40	0.54
1:A:42:ARG:HH22	1:A:155:MET:CE	2.23	0.52
1:A:42:ARG:NH2	1:A:155:MET:HE2	2.26	0.50
1:A:172:THR:HG22	1:A:231:GLU:OE1	2.11	0.50
1:A:172:THR:CG2	1:A:231:GLU:OE1	2.60	0.49
1:A:36:ALA:HB2	1:A:46:VAL:HG22	1.95	0.48
1:A:99:ILE:HG12	1:A:229:MET:HG3	1.95	0.48
1:A:183:ILE:HG21	1:A:191:MET:HE1	1.95	0.47
1:A:23:GLU:HB2	1:A:33:TYR:CE1	2.49	0.47
1:A:369:ARG:HD3	1:A:376:TYR:CE1	2.49	0.47
1:A:237:HIS:NE2	1:A:351:LEU:O	2.39	0.47
1:A:432:ARG:NH1	1:A:434:ALA:HB2	2.30	0.47
1:A:221:PHE:HB3	1:A:380:ILE:HD11	1.97	0.46
1:A:122:ALA:HB3	1:A:180:HIS:CG	2.51	0.46
1:A:183:ILE:CG2	1:A:189:ALA:HB3	2.46	0.45
1:A:183:ILE:HD12	1:A:243:LYS:HD2	1.97	0.45
1:A:272:PHE:O	1:A:358:THR:HG23	2.17	0.45
1:A:177:MET:HG3	1:A:180:HIS:HE1	1.81	0.43
1:A:248:PRO:HG2	1:A:250:THR:CG2	2.47	0.43
1:A:377:LEU:HB3	1:A:378:PRO:CD	2.47	0.43
1:A:218:ARG:HA	1:A:218:ARG:HD3	1.61	0.42
1:A:248:PRO:HG2	1:A:250:THR:HG23	2.01	0.42
1:A:116:ILE:HG23	4:A:524:HOH:O	2.20	0.41
1:A:264:VAL:CG1	1:A:269:LEU:HD11	2.51	0.41
1:A:194:ARG:HE	1:A:194:ARG:HB2	1.58	0.41
1:A:264:VAL:HG13	1:A:269:LEU:HD11	2.02	0.41
1:A:177:MET:CB	1:A:180:HIS:CE1	3.04	0.41
1:A:167:LEU:HB3	1:A:199[B]:CYS:SG	2.61	0.40
1:A:258:MET:SD	1:A:258:MET:N	2.93	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	407/455 (90%)	390 (96%)	14 (3%)	3 (1%)	22 23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	209	ASN
1	A	180	HIS
1	A	30	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	368/405 (91%)	349 (95%)	19 (5%)	23 27

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	VAL
1	A	23	GLU
1	A	27	ASP
1	A	51	THR
1	A	52	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	77	THR
1	A	116	ILE
1	A	152	LYS
1	A	172	THR
1	A	174	SER
1	A	205	ASN
1	A	210	LYS
1	A	214	ILE
1	A	218	ARG
1	A	243	LYS
1	A	258	MET
1	A	291	LYS
1	A	346	LYS
1	A	371	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	180	HIS
1	A	207	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FLC	A	501	2	12,12,12	1.12	0	17,17,17	2.32	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FLC	A	501	2	-	6/16/16/16	-

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	FLC	OB2-CBC-CB	6.02	123.51	113.05
3	A	501	FLC	CB-CG-CGC	-3.00	106.55	113.81
3	A	501	FLC	OB1-CBC-CB	-2.71	118.42	122.25
3	A	501	FLC	CA-CB-CBC	-2.67	104.37	110.11
3	A	501	FLC	CG-CB-CA	2.63	116.03	109.16
3	A	501	FLC	CB-CA-CAC	-2.62	107.47	113.81
3	A	501	FLC	OA2-CAC-CA	2.20	121.42	114.35

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	FLC	OHB-CB-CBC-OB1
3	A	501	FLC	CA-CB-CBC-OB2
3	A	501	FLC	CG-CB-CBC-OB1
3	A	501	FLC	CG-CB-CBC-OB2
3	A	501	FLC	CA-CB-CBC-OB1
3	A	501	FLC	OHB-CB-CBC-OB2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	414/455 (90%)	0.50	40 (9%) <b>7</b> <b>12</b>	35, 45, 76, 94	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	178	ASN	8.3
1	A	179	GLY	7.7
1	A	52	TYR	7.3
1	A	250	THR	6.6
1	A	177	MET	6.5
1	A	210	LYS	6.1
1	A	258	MET	6.0
1	A	28	PRO	5.4
1	A	30	PRO	5.2
1	A	257	PHE	5.2
1	A	209	ASN	4.9
1	A	176	ILE	4.8
1	A	175	MET	4.7
1	A	145	PRO	4.6
1	A	180	HIS	4.6
1	A	214	ILE	4.4
1	A	146	GLY	4.1
1	A	260	HIS	3.9
1	A	251	PRO	3.9
1	A	158	LEU	3.7
1	A	208	GLU	3.6
1	A	259	GLN	3.3
1	A	147	ALA	3.3
1	A	400	ALA	3.3
1	A	174	SER	3.2
1	A	29	GLY	3.2
1	A	27	ASP	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	45	ILE	2.9
1	A	171	PHE	2.9
1	A	169	ASP	2.9
1	A	228	LEU	2.7
1	A	43	ILE	2.5
1	A	160	ILE	2.5
1	A	401	SER	2.4
1	A	156	THR	2.4
1	A	249	ILE	2.3
1	A	44	TRP	2.3
1	A	195	PHE	2.3
1	A	438	PRO	2.2
1	A	232	LEU	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	FLC	A	501	13/13	0.94	0.27	84,91,95,95	0
2	ZN	A	500	1/1	0.98	0.04	54,54,54,54	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.