

Full wwPDB X-ray Structure Validation Report (i)

Jul 11, 2022 – 01:25 pm BST

PDB ID : 6ZBZ

Title : Small-molecule inhibitors of the PDZ domain of Dishevelled proteins interrupt

Wnt signalling

Authors: Roske, Y.; Heinemann, U.; Oschkinat, H.

Deposited on : 2020-06-09

Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.29

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0267$

CCP4 : 7.1.010 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.29

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	95	5% 87%	7% 5%
1	В	95	81%	15% • •
1	С	95	82%	9% • 7%
1	D	95	71%	21% • 7%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Dishevelled, dsh homolog 3 (Drosophila), isoform CRA_b.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	90	Total	С	N	О	S	0	4	0
1	A	90	693	438	115	136	4	0	4	0
1	В	93	Total	С	N	О	S	0	4	0
1	Б	90	717	449	122	141	5	0	4	0
1	С	88	Total	С	N	О	S	0	2	0
1		00	669	422	114	129	4	U		U
1	D	88	Total	С	N	О	S	0	1	0
1	ט	00	662	415	114	129	4	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	242	ALA	-	expression tag	UNP Q9UG07
A	336	SER	-	expression tag	UNP Q9UG07
В	242	ALA	-	expression tag	UNP Q9UG07
В	336	SER	-	expression tag	UNP Q9UG07
С	242	ALA	-	expression tag	UNP Q9UG07
С	336	SER	-	expression tag	UNP Q9UG07
D	242	ALA	-	expression tag	UNP Q9UG07
D	336	SER	-	expression tag	UNP Q9UG07


• Molecule 2 is 2-[(3,4-dimethylphenyl)sulfonylamino]-5-fluoranyl-benzoic acid (three-letter code: QE5) (formula: $C_{15}H_{14}FNO_4S$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		Atoms					ZeroOcc	AltConf
2	٨	1	Total	С	F	N	О	S	0	0
2	А	1	22	15	1	1	4	1	0	U
2	В	1	Total	С	F	N	О	S	0	0
2	Ъ	1	22	15	1	1	4	1	0	U
2	С	1	Total	С	F	N	О	S	0	0
2	C	1	22	15	1	1	4	1	0	U
2	D	1	Total	С	F	N	О	S	0	0
2	D	1	22	15	1	1	4	1		0

 \bullet Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	94	Total O 94 94	0	0
4	В	115	Total O 115 115	0	0
4	С	120	Total O 120 120	0	0
4	D	58	Total O 58 58	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dishevelled, dsh homolog 3 (Drosophila), isoform CRA b





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	56.84Å 70.06Å 87.18Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.59 - 1.60	Depositor
resolution (A)	34.59 - 1.60	EDS
% Data completeness	99.8 (34.59-1.60)	Depositor
(in resolution range)	99.8 (34.59-1.60)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.70 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
P.P.	0.175 , 0.223	Depositor
R, R_{free}	0.182 , 0.230	DCC
R_{free} test set	2328 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for twinning ²	$ < L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3228	wwPDB-VP
Average B, all atoms $(Å^2)$	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: QE5, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ $ \# Z > 5$		RMSZ	# Z > 5	
1	A	0.66	0/711	0.65	0/958	
1	В	0.66	0/736	0.66	0/991	
1	С	0.66	0/681	0.66	0/916	
1	D	0.67	0/670	0.67	0/897	
All	All	0.66	0/2798	0.66	0/3762	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	693	0	716	6	0
1	В	717	0	732	11	0
1	С	669	0	689	6	0
1	D	662	0	672	12	0
2	A	22	0	0	0	0
2	В	22	0	0	0	0
2	С	22	0	0	0	0
2	D	22	0	0	2	0
3	В	4	0	6	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	С	8	0	12	1	0
4	A	94	0	0	0	0
4	В	115	0	0	2	0
4	С	120	0	0	0	0
4	D	58	0	0	0	0
All	All	3228	0	2827	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A + 1	A + a 2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:D:253:ASN:O	1:D:253:ASN:ND2	2.13	0.81
1:D:323:VAL:HG21	2:D:401:QE5:NAN	1.99	0.77
1:B:301:GLN:OE1	1:B:306[A]:ASN:ND2	2.33	0.62
1:B:258:ASN:C	1:B:258:ASN:HD22	2.02	0.61
1:D:321:GLU:O	1:D:325:LYS:HG3	2.06	0.56
1:C:274:GLY:HA2	1:C:298:MET:SD	2.46	0.56
1:B:246:ASN:HB3	1:B:335:LYS:HB2	1.87	0.55
1:C:258:ASN:H	1:C:258:ASN:HD22	1.54	0.54
1:B:317:ARG:NH2	4:B:501:HOH:O	2.39	0.54
1:D:267:GLN:HE21	1:D:312:ASN:H	1.56	0.54
1:A:248:ILE:HA	1:B:306[A]:ASN:HD21	1.73	0.53
1:D:305:ILE:HG21	1:D:318:VAL:HG11	1.91	0.52
1:C:325:LYS:HD2	3:C:402:EDO:H11	1.92	0.51
1:D:271:ARG:HD2	1:D:309:ASN:HA	1.92	0.50
1:D:306:ASN:HD21	1:D:308:GLU:HB2	1.76	0.50
1:D:302:VAL:HG22	1:D:331:LEU:HD22	1.95	0.49
1:B:302:VAL:HG12	1:B:322:ILE:HD13	1.94	0.49
1:C:305:ILE:HD12	1:C:322:ILE:HD11	1.93	0.49
1:A:302:VAL:HG22	1:A:331[B]:LEU:HD12	1.96	0.47
1:B:253:ASN:HD22	1:B:255:GLU:H	1.63	0.47
1:A:268:SER:OG	1:A:309:ASN:HA	2.14	0.47
1:D:301:GLN:HB3	1:D:332:THR:HB	1.96	0.46
1:A:247:ILE:O	1:B:273:ASP:OD2	2.34	0.45
1:D:302:VAL:HG22	1:D:331:LEU:CD2	2.47	0.45
1:B:283:LYS:NZ	4:B:504:HOH:O	2.49	0.44
1:C:301:GLN:HB2	1:C:332:THR:HB	2.00	0.43
1:D:323:VAL:HG21	2:D:401:QE5:SAV	2.58	0.43
1:B:298[B]:MET:HE3	1:B:300:LEU:HD21	2.00	0.43

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$egin{array}{ll} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	Clash overlap (Å)
1:D:319:LEU:HD12	1:D:319:LEU:HA	1.87	0.42
1:C:302:VAL:HG21	1:C:319:LEU:HD21	2.02	0.41
1:B:258:ASN:C	1:B:258:ASN:ND2	2.72	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	89/95 (94%)	87 (98%)	2 (2%)	0	100	100
1	В	95/95 (100%)	92 (97%)	3 (3%)	0	100	100
1	С	86/95 (90%)	85 (99%)	1 (1%)	0	100	100
1	D	84/95 (88%)	80 (95%)	4 (5%)	0	100	100
All	All	354/380 (93%)	344 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	79/78 (101%)	78 (99%)	1 (1%)	69 50
1	В	80/78 (103%)	75 (94%)	5 (6%)	18 4

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	C	74/78~(95%)	73 (99%)	1 (1%)	67 47
1	D	72/78~(92%)	66 (92%)	6 (8%)	11 2
All	All	305/312 (98%)	292 (96%)	13 (4%)	34 9

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	306	ASN
1	В	245	LEU
1	В	253	ASN
1	В	258	ASN
1	В	321[A]	GLU
1	В	321[B]	GLU
1	С	258	ASN
1	D	253	ASN
1	D	273	ASP
1	D	282	MET
1	D	304	GLU
1	D	336[A]	SER
1	D	336[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	301	GLN
1	A	306	ASN
1	В	253	ASN
1	В	258	ASN
1	В	267	GLN
1	В	269	ASN
1	С	258	ASN
1	D	267	GLN
1	D	306	ASN
1	D	324	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trmo	Chain	Res	Link	Во	ond leng	ths	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	QE5	В	401	-	23,23,23	2.26	3 (13%)	34,34,34	1.74	3 (8%)
2	QE5	D	401	-	23,23,23	2.32	3 (13%)	34,34,34	1.71	2 (5%)
2	QE5	A	401	-	23,23,23	2.22	3 (13%)	34,34,34	1.72	2 (5%)
2	QE5	С	401	-	23,23,23	2.24	3 (13%)	34,34,34	1.71	2 (5%)
3	EDO	В	402	-	3,3,3	0.50	0	2,2,2	0.31	0
3	EDO	С	402	-	3,3,3	0.50	0	2,2,2	0.30	0
3	EDO	С	403	-	3,3,3	0.49	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QE5	В	401	-	-	0/15/15/15	0/2/2/2
2	QE5	D	401	-	-	1/15/15/15	0/2/2/2
2	QE5	A	401	-	-	0/15/15/15	0/2/2/2
2	QE5	С	401	-	-	0/15/15/15	0/2/2/2
3	EDO	В	402	-	-	1/1/1/1	-
3	EDO	С	402	-	-	1/1/1/1	-
3	EDO	С	403	-	-	0/1/1/1	-



All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
2	D	401	QE5	OAD-SAV	6.20	1.50	1.43
2	D	401	QE5	OAE-SAV	6.14	1.50	1.43
2	В	401	QE5	OAD-SAV	6.10	1.50	1.43
2	A	401	QE5	OAD-SAV	6.06	1.50	1.43
2	A	401	QE5	OAE-SAV	6.01	1.50	1.43
2	С	401	QE5	OAD-SAV	6.00	1.50	1.43
2	С	401	QE5	OAE-SAV	5.99	1.50	1.43
2	В	401	QE5	OAE-SAV	5.97	1.50	1.43
2	D	401	QE5	CAS-NAN	-5.63	1.33	1.42
2	В	401	QE5	CAS-NAN	-5.46	1.33	1.42
2	С	401	QE5	CAS-NAN	-5.37	1.33	1.42
2	A	401	QE5	CAS-NAN	-5.11	1.34	1.42

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	В	401	QE5	OAE-SAV-OAD	-8.31	109.34	119.55
2	A	401	QE5	OAE-SAV-OAD	-8.15	109.53	119.55
2	D	401	QE5	OAE-SAV-OAD	-8.08	109.62	119.55
2	С	401	QE5	OAE-SAV-OAD	-8.05	109.65	119.55
2	A	401	QE5	CAI-CAR-CAM	-2.77	119.70	123.29
2	С	401	QE5	CAI-CAR-CAM	-2.70	119.79	123.29
2	В	401	QE5	CAI-CAR-CAM	-2.61	119.90	123.29
2	D	401	QE5	CAI-CAR-CAM	-2.53	120.00	123.29
2	В	401	QE5	CAT-SAV-NAN	2.29	109.71	106.83

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	401	QE5	CAS-NAN-SAV-OAE
3	С	402	EDO	O1-C1-C2-O2
3	В	402	EDO	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	401	QE5	2	0
3	С	402	EDO	1	0




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	90/95~(94%)	0.13	5 (5%) 24 22	17, 24, 43, 53	0
1	В	93/95~(97%)	0.06	3 (3%) 47 44	17, 25, 49, 58	0
1	С	88/95 (92%)	-0.09	2 (2%) 60 59	17, 26, 42, 55	0
1	D	88/95 (92%)	0.48	9 (10%) 6 5	22, 42, 63, 67	0
All	All	359/380 (94%)	0.14	19 (5%) 26 24	17, 27, 53, 67	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	309	ASN	3.6
1	D	274	GLY	3.5
1	С	274	GLY	3.4
1	D	253	ASN	3.3
1	D	255	GLU	3.1
1	D	271	ARG	2.9
1	D	273	ASP	2.9
1	A	269	ASN	2.9
1	В	244	SER	2.7
1	D	326	PRO	2.7
1	A	245	LEU	2.6
1	D	308	GLU	2.5
1	В	271	ARG	2.4
1	A	268	SER	2.3
1	D	258	ASN	2.2
1	С	309	ASN	2.2
1	A	274	GLY	2.1
1	A	256	LYS	2.1
1	В	245	LEU	2.1



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	EDO	С	403	4/4	0.72	0.17	56,57,57,58	0
3	EDO	В	402	4/4	0.74	0.20	44,45,47,48	0
3	EDO	С	402	4/4	0.79	0.26	37,42,43,49	0
2	QE5	D	401	22/22	0.86	0.22	39,53,66,69	0
2	QE5	В	401	22/22	0.95	0.08	24,34,42,44	0
2	QE5	С	401	22/22	0.95	0.10	22,30,35,38	0
2	QE5	A	401	22/22	0.95	0.08	20,28,30,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



















6.5 Other polymers (i)

There are no such residues in this entry.

