



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 14, 2023 – 05:13 am GMT

PDB ID : 3ZF6
Title : Phage dUTPases control transfer of virulence genes by a proto-oncogenic G protein-like mechanism. (Staphylococcus bacteriophage 80alpha dUTPase D81A D110C S168C mutant with dUpNHpp).
Authors : Tormo-Mas, M.A.; Donderis, J.; Garcia-Caballer, M.; Alt, A.; Mir-Sanchis, I.; Marina, A.; Penades, J.R.
Deposited on : 2012-12-10
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3518 (2.60-2.60)

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DUTPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	169	1322	832	229	257	4	0	1	0

There are 37 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-33	MET	-	expression tag	UNP A4ZF98
A	-32	GLY	-	expression tag	UNP A4ZF98
A	-31	SER	-	expression tag	UNP A4ZF98
A	-30	SER	-	expression tag	UNP A4ZF98
A	-29	HIS	-	expression tag	UNP A4ZF98
A	-28	HIS	-	expression tag	UNP A4ZF98
A	-27	HIS	-	expression tag	UNP A4ZF98
A	-26	HIS	-	expression tag	UNP A4ZF98
A	-25	HIS	-	expression tag	UNP A4ZF98
A	-24	HIS	-	expression tag	UNP A4ZF98
A	-23	SER	-	expression tag	UNP A4ZF98
A	-22	SER	-	expression tag	UNP A4ZF98
A	-21	GLY	-	expression tag	UNP A4ZF98
A	-20	LEU	-	expression tag	UNP A4ZF98
A	-19	VAL	-	expression tag	UNP A4ZF98
A	-18	PRO	-	expression tag	UNP A4ZF98
A	-17	ARG	-	expression tag	UNP A4ZF98
A	-16	GLY	-	expression tag	UNP A4ZF98
A	-15	SER	-	expression tag	UNP A4ZF98
A	-14	HIS	-	expression tag	UNP A4ZF98
A	-13	MET	-	expression tag	UNP A4ZF98
A	-12	ALA	-	expression tag	UNP A4ZF98
A	-11	SER	-	expression tag	UNP A4ZF98
A	-10	MET	-	expression tag	UNP A4ZF98
A	-9	THR	-	expression tag	UNP A4ZF98
A	-8	GLY	-	expression tag	UNP A4ZF98
A	-7	GLY	-	expression tag	UNP A4ZF98

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLN	-	expression tag	UNP A4ZF98
A	-5	GLN	-	expression tag	UNP A4ZF98
A	-4	MET	-	expression tag	UNP A4ZF98
A	-3	GLY	-	expression tag	UNP A4ZF98
A	-2	ARG	-	expression tag	UNP A4ZF98
A	-1	GLY	-	expression tag	UNP A4ZF98
A	0	SER	-	expression tag	UNP A4ZF98
A	81	ALA	ASP	engineered mutation	UNP A4ZF98
A	110	CYS	ASP	engineered mutation	UNP A4ZF98
A	168	CYS	SER	engineered mutation	UNP A4ZF98

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Ni 2 2	0	0

- Molecule 3 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula: C₉H₁₃N₂O₈P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O P 20 9 2 8 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	50	Total O 50 50	0	0

SEQUENCE-PLOTS INFOmissingINFO

3 Data and refinement statistics i

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	87.46Å 87.46Å 87.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.71 – 2.60	Depositor
% Data completeness (in resolution range)	99.8 (35.71-2.60)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.51 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.224 , 0.273	Depositor
Wilson B-factor (Å ²)	59.3	Xtrriage
Anisotropy	0.000	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.063 for l,-k,h	Xtrriage
Total number of atoms	1394	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, UMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/1343	0.50	2/1813 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	167	SER	N-CA-CB	-6.50	100.76	110.50
1	A	166	GLY	N-CA-C	5.11	125.88	113.10

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1322	0	1309	14	5
2	A	2	0	0	0	0
3	A	20	0	11	0	0
4	A	50	0	0	3	1
All	All	1394	0	1320	14	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:GLU:OE2	1:A:159:GLU:HA	1.63	0.96
1:A:160:ARG:HG2	1:A:161:GLY:N	1.96	0.80
1:A:159:GLU:OE2	1:A:159:GLU:CA	2.30	0.78
1:A:160:ARG:HG2	1:A:161:GLY:H	1.49	0.75
1:A:160:ARG:CG	1:A:161:GLY:N	2.63	0.61
1:A:76:GLU:OE1	4:A:2030:HOH:O	2.17	0.60
1:A:99:ASP:O	1:A:100:LYS:HB2	2.10	0.52
1:A:160:ARG:HB2	1:A:160:ARG:CZ	2.45	0.46
1:A:42:LYS:HG2	1:A:93:LYS:HG3	1.98	0.45
1:A:97[B]:GLU:HG2	4:A:2039:HOH:O	2.17	0.44
1:A:160:ARG:CZ	1:A:160:ARG:CB	2.93	0.44
1:A:8:LYS:HB3	1:A:50:ALA:HB3	2.01	0.43
1:A:64:ARG:HD3	4:A:2019:HOH:O	2.18	0.43
1:A:163:LYS:HA	1:A:164:GLY:HA2	1.76	0.43

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:CYS:SG	1:A:168:CYS:SG[5_555]	1.08	1.12
1:A:167:SER:OG	4:A:2050:HOH:O[5_555]	1.85	0.35
1:A:24:ASP:OD1	1:A:160:ARG:NH2[5_555]	1.86	0.34
1:A:110:CYS:CB	1:A:168:CYS:SG[5_555]	1.92	0.28
1:A:134:ARG:NH1	1:A:170:VAL:O[5_555]	2.02	0.18

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

4.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.