

# wwPDB X-ray Structure Validation Summary Report (i)

#### Aug 30, 2022 – 04:46 pm BST

PDB ID	:	7ZII
Title	:	Crystal structure of human tryptophan hydroxylase 1 in complex with inhibitor
		KM-05-193
Authors	:	Schuetz, A.; Mallow, K.; Nazare, M.; Specker, E.; Heinemann, U.
Deposited on		
Resolution	:	1.63 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity		4 02b 467
·		
Mogul	:	1.8.4, CSD as $541$ be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.30
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.30

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	4693 (1.64-1.60)
Clashscore	141614	5002 (1.64-1.60)
Ramachandran outliers	138981	4888 (1.64-1.60)
Sidechain outliers	138945	4887 (1.64-1.60)
RSRZ outliers	127900	4609 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
			5%					
1	А	326	83%	5%	12%			



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2680 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Tryptophan 5-hydroxylase 1.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	287	Total 2325	C 1497	N 388	0 428	S 12	0	1	0

Chain	Residue	Modelled	Actual	Comment	Reference
А	76	MET	-	initiating methionine	UNP P17752
А	77	LYS	-	expression tag	UNP P17752
А	78	HIS	-	expression tag	UNP P17752
А	79	HIS	-	expression tag	UNP P17752
А	80	HIS	-	expression tag	UNP P17752
А	81	HIS	-	expression tag	UNP P17752
А	82	HIS	-	expression tag	UNP P17752
A	83	HIS	-	expression tag	UNP P17752
А	84	HIS	-	expression tag	UNP P17752
А	85	GLY	-	expression tag	UNP P17752
А	86	ALA	-	expression tag	UNP P17752
А	87	ALA	-	expression tag	UNP P17752
А	88	GLY	-	expression tag	UNP P17752
А	89	THR	-	expression tag	UNP P17752
А	90	SER	-	expression tag	UNP P17752
А	91	LEU	-	expression tag	UNP P17752
A	92	TYR	-	expression tag	UNP P17752
А	93	LYS	-	expression tag	UNP P17752
A	94	LYS	-	expression tag	UNP P17752
А	95	ALA	-	expression tag	UNP P17752
А	96	GLY	-	expression tag	UNP P17752
А	97	GLU	-	expression tag	UNP P17752
А	98	ASN	-	expression tag	UNP P17752
А	99	LEU	-	expression tag	UNP P17752
А	100	TYR	-	expression tag	UNP P17752
А	101	PHE	-	expression tag	UNP P17752
А	102	GLN	-	expression tag	UNP P17752

There are 29 discrepancies between the modelled and reference sequences:

Continued on next page...




Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
А	103	GLY	-	expression tag	UNP P17752
А	104	SER	-	expression tag	UNP P17752

• Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Fe 1 1	0	0


• Molecule 3 is 8-(5 {H}-[1,3]dioxolo[4,5-f]benzimidazol-6-ylmethyl)-7-(phenylmethyl)-3-prop yl-purine-2,6-dione (three-letter code: IVQ) (formula: C<sub>24</sub>H<sub>22</sub>N<sub>6</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	А	1	Total 34	C 24	N 6	0 4	0	0

• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).





Mol	Chain	Residues	Ator	ms	ZeroOcc	AltConf
4	А	1	Total 6	C O 3 3	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	314	Total         O           314         314	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tryptophan 5-hydroxylase 1





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	46.97Å 57.95Å 59.35Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $97.62^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	29.55 - 1.63	Depositor
Resolution (A)	46.55 - 1.63	EDS
% Data completeness	99.7 (29.55-1.63)	Depositor
(in resolution range)	99.7 (46.55 - 1.63)	EDS
R <sub>merge</sub>	0.07	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.50 (at 1.63 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
D D	0.167 , $0.192$	Depositor
$R, R_{free}$	0.168 , $0.191$	DCC
$R_{free}$ test set	1978 reflections $(5.00\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	17.8	Xtriage
Anisotropy	0.375	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for twinning <sup>2</sup>	$ L  > = 0.49, < L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2680	wwPDB-VP
Average B, all atoms $(Å^2)$	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.45% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IVQ, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Ch	Chain	Boi	nd lengths	Bond angles		
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.49	2/2391~(0.1%)	0.58	0/3231	

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	252[A]	CYS	CB-SG	-5.85	1.72	1.81
1	А	252[B]	CYS	CB-SG	-5.85	1.72	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2325	0	2276	12	0
2	А	1	0	0	0	0
3	А	34	0	0	0	0
4	А	6	0	8	0	0
5	А	314	0	0	1	0
All	All	2680	0	2284	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



7ZII
------

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:250:PHE:HD2	1:A:252[B]:CYS:HG	1.43	0.67
1:A:250:PHE:HD2	1:A:252[B]:CYS:SG	2.20	0.65
1:A:229:ILE:HD12	1:A:252[B]:CYS:SG	2.46	0.55
1:A:390:THR:O	1:A:394:LYS:HG3	2.07	0.54
1:A:263:PHE:CD1	1:A:338:ILE:HD12	2.49	0.48

The worst 5 of 12 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	А	284/326~(87%)	281 (99%)	3~(1%)	0	100 100	

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	256/285~(90%)	254~(99%)	2(1%)	81 69	

All (2) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	А	227	PHE
1	А	366	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		Chain	Dec	Tiple	Bo	ond leng	$\mathbf{ths}$	В	ond ang	les
Mol Type	Chain	nes	Link	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2	
3	IVQ	А	502	2	33,39,39	2.10	7 (21%)	$32,\!57,\!57$	2.12	7 (21%)
4	GOL	А	503	-	$5,\!5,\!5$	0.35	0	$5,\!5,\!5$	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mo	r l	ype	Chain	Res	Link	Chirals	Torsions	Rings
3	]	VQ	А	502	2	-	3/11/17/17	0/6/6/6
4	(	GOL	А	503	-	-	0/4/4/4	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
3	А	502	IVQ	CAJ-CBD	-5.40	1.33	1.41
3	А	502	IVQ	CAI-CBC	-4.81	1.34	1.41
3	А	502	IVQ	CAO-CAV	-4.27	1.41	1.51
3	А	502	IVQ	C5-C6	-4.15	1.34	1.45
3	А	502	IVQ	C2-N3	-3.63	1.35	1.37

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	А	502	IVQ	C6-N1-C2	-9.00	119.05	127.35
3	А	502	IVQ	O2-C2-N3	-3.05	120.10	122.00
3	А	502	IVQ	CAL-N3-C2	2.80	120.30	117.14
3	А	502	IVQ	N1-C2-N3	2.48	119.79	115.39
3	А	502	IVQ	CAM-OAT-CAX	-2.42	102.20	105.34

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	502	IVQ	N7-CAO-CAV-CAH
3	А	502	IVQ	N7-CAO-CAV-CAG
3	А	502	IVQ	C8-CAN-CAW-NAP


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.



The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$<$ RSRZ $>$	#RSRZ>2		$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	А	287/326~(88%)	0.02	16 (5%) 24	21	12, 20, 47, 78	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	266	PRO	5.6
1	А	264	TYR	5.4
1	А	348	HIS	4.3
1	А	366	ILE	4.2
1	А	130	ASP	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q < 0.9
4	GOL	А	503	6/6	0.83	0.24	$46,\!50,\!55,\!58$	0
3	IVQ	А	502	34/34	0.91	0.14	16,22,51,61	0
2	FE	А	501	1/1	1.00	0.08	14,14,14,14	0



7ZII

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



### 6.5 Other polymers (i)

There are no such residues in this entry.

