



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2023 – 11:57 AM EDT

PDB ID : 2ZL7
Title : Atomic resolution structural characterization of recognition of histo-blood group antigens by Norwalk virus
Authors : Choi, J.M.; Huston, A.M.; Estes, M.K.; Prasad, B.V.V.
Deposited on : 2008-04-02
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	 7% 89% 7% .
1	B	295	 10% 88% 7% ..
2	C	3	 33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NGA	C	3	X	-	-	-
5	ACT	A	520	-	-	X	-
5	ACT	B	521	-	-	X	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 5106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 58 kd capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	285	Total	C	N	O	S	0	1	0
			2154	1380	357	409	8			
1	B	286	Total	C	N	O	S	0	3	0
			2169	1390	359	412	8			

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)]beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	3	Total	C	N	O	0	0	0
			36	20	1	15			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
6	A	347	Total	O	0	0
			347	347		
6	B	389	Total	O	0	0
			389	389		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 58 kd capsid protein



- Molecule 1: 58 kd capsid protein



- Molecule 2: alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)]beta-D-galactopyranose



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	83.07Å 83.07Å 164.32Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.07 – 1.35 30.07 – 1.36	Depositor EDS
% Data completeness (in resolution range)	98.5 (30.07-1.35) 98.4 (30.07-1.36)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 1.36Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.171 , 0.188 0.178 , 0.195	Depositor DCC
R_{free} test set	6957 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	13.1	Xtrriage
Anisotropy	0.007	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.036 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5106	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, FUC, MG, CA, ACT, NGA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/2227	0.66	0/3054
1	B	0.48	0/2249	0.72	3/3084 (0.1%)
All	All	0.48	0/4476	0.69	3/6138 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	383	SER	N-CA-C	-8.51	88.02	111.00
1	B	291	ARG	NE-CZ-NH1	6.58	123.59	120.30
1	B	346	ASP	CB-CG-OD1	5.62	123.36	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	382	PRO	Peptide
1	B	413	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2154	0	2075	30	0
1	B	2169	0	2099	34	0
2	C	36	0	33	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	4	0	3	13	0
5	B	4	0	3	15	0
6	A	347	0	0	8	0
6	B	389	0	0	4	0
All	All	5106	0	4213	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:LEU:HD11	5:A:520:ACT:H3	1.25	1.17
1:B:304:LEU:HD11	5:B:521:ACT:H3	1.20	1.14
1:B:304:LEU:HD11	5:B:521:ACT:CH3	1.95	0.94
1:B:304:LEU:CD1	5:B:521:ACT:H3	1.97	0.94
1:A:370:VAL:CG1	6:A:690:HOH:O	2.17	0.91
1:A:304:LEU:CD1	5:A:520:ACT:H3	2.00	0.90
1:A:304:LEU:HD21	5:A:520:ACT:CH3	2.02	0.90
1:A:370:VAL:HG12	6:A:690:HOH:O	1.74	0.86
1:B:414:PHE:O	1:B:414:PHE:HD2	1.59	0.85
1:A:304:LEU:HD11	5:A:520:ACT:CH3	2.09	0.78
1:B:291:ARG:HD2	1:B:370:VAL:HG22	1.66	0.77
1:A:396:GLY:O	6:A:862:HOH:O	2.03	0.76
1:B:304:LEU:HD13	6:B:959:HOH:O	1.88	0.73
1:B:414:PHE:O	1:B:414:PHE:CD2	2.42	0.72
6:A:690:HOH:O	1:B:426:PRO:HG3	1.90	0.70
1:B:304:LEU:HD21	5:B:521:ACT:CH3	2.22	0.70
1:B:329:HIS:HD2	1:B:344:ASP:OD1	1.77	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:291:ARG:CD	1:B:370:VAL:HG22	2.25	0.67
1:A:304:LEU:HD21	5:A:520:ACT:H2	1.76	0.67
1:B:329:HIS:HE1	1:B:377:SER:OG	1.79	0.66
1:A:329:HIS:HD2	1:A:344:ASP:OD1	1.79	0.66
6:A:521:HOH:O	1:B:286:HIS:HE1	1.79	0.64
1:B:413:GLY:HA3	1:B:414:PHE:CD1	2.32	0.64
1:A:286:HIS:HE1	6:B:663:HOH:O	1.79	0.64
1:B:289:LYS:HE2	5:B:521:ACT:H1	1.79	0.63
1:A:370:VAL:HG11	6:A:690:HOH:O	1.93	0.63
1:A:329:HIS:HE1	1:A:377:SER:OG	1.82	0.61
1:B:396:GLY:O	6:B:1036:HOH:O	2.16	0.59
1:A:304:LEU:HD21	5:A:520:ACT:H1	1.83	0.59
1:A:304:LEU:CD2	5:A:520:ACT:CH3	2.79	0.58
1:A:304:LEU:CG	5:A:520:ACT:H3	2.34	0.58
1:A:289:LYS:HE2	5:A:520:ACT:H1	1.84	0.58
1:B:286:HIS:HD2	6:B:662:HOH:O	1.87	0.57
1:A:286:HIS:HD2	6:A:537:HOH:O	1.87	0.56
1:A:314:GLY:O	1:A:355:HIS:HD2	1.87	0.56
1:A:399:ILE:H	1:A:399:ILE:HD12	1.70	0.55
1:A:289:LYS:CE	5:A:520:ACT:H1	2.37	0.53
1:A:399:ILE:HD12	1:A:399:ILE:N	2.23	0.53
1:B:289:LYS:CE	5:B:521:ACT:H1	2.38	0.52
1:B:304:LEU:HD21	5:B:521:ACT:H2	1.92	0.51
1:B:304:LEU:CD1	5:B:521:ACT:CH3	2.75	0.50
1:B:414:PHE:H	1:B:415:GLY:HA2	1.77	0.50
1:A:239:SER:HB3	1:A:496:ILE:HG23	1.93	0.50
1:A:289:LYS:NZ	5:A:520:ACT:H1	2.27	0.49
1:B:239:SER:HB3	1:B:497:ILE:HG23	1.95	0.49
6:A:720:HOH:O	1:B:444:HIS:HD2	1.95	0.49
1:A:346:ASP:OD2	1:A:381:HIS:HD2	1.95	0.49
1:B:413:GLY:O	1:B:416:GLU:HB2	2.13	0.49
1:A:304:LEU:CG	5:A:520:ACT:CH3	2.91	0.48
1:B:399:ILE:HD12	1:B:399:ILE:C	2.34	0.48
1:A:331:ASN:HD21	1:A:340:GLN:NE2	2.11	0.48
1:B:304:LEU:CG	5:B:521:ACT:H3	2.45	0.46
1:B:304:LEU:HD21	5:B:521:ACT:H1	1.97	0.45
1:B:304:LEU:CG	5:B:521:ACT:CH3	2.95	0.45
1:A:249:PRO:HG2	5:B:521:ACT:H2	1.98	0.44
1:B:304:LEU:CD2	5:B:521:ACT:CH3	2.95	0.43
1:A:474:LYS:HG3	1:A:512:LEU:HD21	2.01	0.42
1:B:232:THR:OG1	1:B:444:HIS:HE1	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:399:ILE:HD12	1:B:400:THR:N	2.35	0.42
1:B:289:LYS:NZ	5:B:521:ACT:H1	2.35	0.41
1:A:304:LEU:CD1	5:A:520:ACT:CH3	2.84	0.41
1:B:291:ARG:HD2	1:B:370:VAL:CG2	2.43	0.41
1:A:249:PRO:CG	5:B:521:ACT:H2	2.51	0.41
1:B:346:ASP:OD2	1:B:381:HIS:HD2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/295 (96%)	276 (98%)	5 (2%)	1 (0%)	34	12
1	B	287/295 (97%)	280 (98%)	6 (2%)	1 (0%)	41	18
All	All	569/590 (96%)	556 (98%)	11 (2%)	2 (0%)	34	12

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	414	PHE
1	A	339	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/250 (97%)	243 (100%)	0	100	100
1	B	246/250 (98%)	244 (99%)	2 (1%)	81	59
All	All	489/500 (98%)	487 (100%)	2 (0%)	91	81

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	414	PHE
1	B	452	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	286	HIS
1	A	300	ASN
1	A	329	HIS
1	A	340	GLN
1	A	355	HIS
1	A	381	HIS
1	A	444	HIS
1	B	286	HIS
1	B	329	HIS
1	B	381	HIS
1	B	386	GLN
1	B	444	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GAL	C	1	2	12,12,12	0.51	0	17,17,17	0.94	1 (5%)
2	FUC	C	2	2	10,10,11	0.41	0	14,14,16	0.42	0
2	NGA	C	3	2	14,14,15	2.35	2 (14%)	17,19,21	2.07	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	C	1	2	-	0/2/22/22	0/1/1/1
2	FUC	C	2	2	-	-	0/1/1/1
2	NGA	C	3	2	1/1/5/7	3/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3	NGA	O7-C7	8.29	1.42	1.23
2	C	3	NGA	C7-N2	2.58	1.43	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	NGA	O7-C7-N2	-6.14	110.66	121.95
2	C	3	NGA	O7-C7-C8	-3.91	114.80	122.06
2	C	1	GAL	O3-C3-C2	-2.58	104.39	110.35
2	C	3	NGA	C8-C7-N2	-2.43	111.98	116.10

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	3	NGA	C1


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	3	NGA	O7-C7-N2-C2
2	C	3	NGA	C1-C2-N2-C7
2	C	3	NGA	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACT	B	521	-	3,3,3	0.71	0	3,3,3	0.97	0
5	ACT	A	520	-	3,3,3	0.66	0	3,3,3	1.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	521	ACT	15	0
5	A	520	ACT	13	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/295 (96%)	0.65	21 (7%) 14 16	8, 13, 27, 34	0
1	B	286/295 (96%)	0.61	29 (10%) 7 8	8, 13, 29, 38	0
All	All	571/590 (96%)	0.63	50 (8%) 10 11	8, 13, 28, 38	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	414	PHE	11.4
1	A	399	ILE	9.9
1	B	399	ILE	8.6
1	B	488	ALA	8.4
1	B	510	PHE	8.3
1	A	400	THR	8.2
1	B	452	THR	7.7
1	A	509	PHE	7.4
1	B	414	PHE	7.2
1	B	400	THR	6.8
1	A	452	THR	6.7
1	B	487	GLY	5.6
1	B	486	ASN	5.0
1	A	486	ASN	4.6
1	B	489	SER	4.2
1	B	413	GLY	4.1
1	A	515	VAL	3.9
1	A	398	SER	3.9
1	B	383	SER	3.9
1	B	465	ASP	3.8
1	A	465	ASP	3.8
1	B	397	SER	3.7
1	A	397	SER	3.7
1	B	490	SER	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	451	PRO	3.5
1	A	488	ALA	3.5
1	A	453	VAL	3.5
1	B	384	GLY	3.4
1	B	451	PRO	3.1
1	B	466	THR	2.9
1	A	261	GLN	2.9
1	B	398	SER	2.9
1	B	468	ARG	2.8
1	B	494	GLN	2.7
1	A	489	SER	2.7
1	A	259	ASN	2.7
1	B	464	PRO	2.7
1	B	415	GLY	2.6
1	B	410	TYR	2.5
1	B	491	GLY	2.5
1	A	468	ARG	2.5
1	A	464	PRO	2.5
1	B	509	ARG	2.5
1	A	487	GLY	2.5
1	A	466	THR	2.5
1	B	516	VAL	2.4
1	B	330	ILE	2.2
1	B	386	GLN	2.1
1	B	493	GLN	2.1
1	A	350	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GAL	C	1	12/12	0.93	0.12	13,16,18,19	0
2	FUC	C	2	10/11	0.94	0.10	13,15,17,17	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NGA	C	3	14/15	0.94	0.11	11,12,15,16	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CA	A	1	1/1	0.95	0.23	42,42,42,42	0
5	ACT	A	520	4/4	0.95	0.11	17,18,18,19	0
5	ACT	B	521	4/4	0.95	0.10	17,17,17,18	0
4	MG	A	519	1/1	0.98	0.06	18,18,18,18	0
4	MG	B	520	1/1	0.98	0.10	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.