



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 12:12 pm BST

PDB ID : 4ZN0  
Title : Structure of the NADPH-dependent thioredoxin reductase from  
Methanosarcina mazei  
Authors : Buey, R.M.; de Pereda, J.M.; Balsera, M.  
Deposited on : 2015-05-04  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	 2% 81% 11% 8%
1	B	311	 2% 82% 7% 10%
1	C	311	 5% 83% 6% 10%
1	D	311	 11% 81% 6% 13%

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 15485 atoms, of which 7463 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thioredoxin reductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	287	4137	1326	2036	360	410	5	0	0	0
1	B	279	3982	1291	1950	339	397	5	0	0	0
1	C	279	3804	1252	1838	332	378	4	0	0	0
1	D	270	3494	1172	1639	322	357	4	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q8PUI1
A	-1	SER	-	expression tag	UNP Q8PUI1
A	0	HIS	-	expression tag	UNP Q8PUI1
A	1	MET	-	expression tag	UNP Q8PUI1
A	2	ALA	-	expression tag	UNP Q8PUI1
A	3	SER	-	expression tag	UNP Q8PUI1
B	-2	GLY	-	expression tag	UNP Q8PUI1
B	-1	SER	-	expression tag	UNP Q8PUI1
B	0	HIS	-	expression tag	UNP Q8PUI1
B	1	MET	-	expression tag	UNP Q8PUI1
B	2	ALA	-	expression tag	UNP Q8PUI1
B	3	SER	-	expression tag	UNP Q8PUI1
C	-2	GLY	-	expression tag	UNP Q8PUI1
C	-1	SER	-	expression tag	UNP Q8PUI1
C	0	HIS	-	expression tag	UNP Q8PUI1
C	1	MET	-	expression tag	UNP Q8PUI1
C	2	ALA	-	expression tag	UNP Q8PUI1
C	3	SER	-	expression tag	UNP Q8PUI1
D	-2	GLY	-	expression tag	UNP Q8PUI1
D	-1	SER	-	expression tag	UNP Q8PUI1
D	0	HIS	-	expression tag	UNP Q8PUI1

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	1	MET	-	expression tag	UNP Q8PUI1
D	2	ALA	-	expression tag	UNP Q8PUI1
D	3	SER	-	expression tag	UNP Q8PUI1

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	22	Total O 22 22	0	0
2	B	23	Total O 23 23	0	0
2	C	9	Total O 9 9	0	0
2	D	14	Total O 14 14	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	117.14Å 182.50Å 152.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.89 – 2.60 46.89 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.89-2.60) 100.0 (46.89-2.60)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.08 (at 2.61Å)	Xtrriage
Refinement program	PHENIX dev_1839	Depositor
R, $R_{free}$	0.217 , 0.241 0.224 , 0.241	Depositor DCC
$R_{free}$ test set	2412 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	61.0	Xtrriage
Anisotropy	0.432	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 56.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	15485	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/2132	0.45	0/2899
1	B	0.23	0/2063	0.43	0/2813
1	C	0.24	0/1996	0.43	0/2728
1	D	0.24	0/1882	0.46	0/2564
All	All	0.24	0/8073	0.45	0/11004

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2101	2036	2030	31	0
1	B	2032	1950	1947	18	0
1	C	1966	1838	1836	11	0
1	D	1855	1639	1635	24	0
2	A	22	0	0	0	0
2	B	23	0	0	2	0
2	C	9	0	0	0	0
2	D	14	0	0	2	0
All	All	8022	7463	7448	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.



The worst 5 of 73 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:96:THR:HG22	1:D:99:GLY:O	1.35	1.22
1:D:96:THR:HG23	1:D:99:GLY:H	1.27	0.99
1:C:279:ARG:NH1	1:C:292:ASP:OD2	1.96	0.98
1:A:282:PRO:HG3	1:B:252:GLU:OE1	1.66	0.94
1:D:6:ASP:HB2	1:D:29:ASP:O	1.75	0.85

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/311 (91%)	267 (94%)	16 (6%)	0	100	100
1	B	275/311 (88%)	264 (96%)	11 (4%)	0	100	100
1	C	273/311 (88%)	258 (94%)	15 (6%)	0	100	100
1	D	263/311 (85%)	253 (96%)	10 (4%)	0	100	100
All	All	1094/1244 (88%)	1042 (95%)	52 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/255 (82%)	197 (94%)	12 (6%)	20	41
1	B	200/255 (78%)	199 (100%)	1 (0%)	88	96
1	C	182/255 (71%)	174 (96%)	8 (4%)	28	53
1	D	152/255 (60%)	150 (99%)	2 (1%)	69	86
All	All	743/1020 (73%)	720 (97%)	23 (3%)	40	66

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	189	ASP
1	B	66	PHE
1	D	6	ASP
1	A	284	TRP
1	C	36	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	113	ASN
1	D	245	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	287/311 (92%)	0.56	6 (2%) 63 58	41, 59, 106, 132	0
1	B	279/311 (89%)	0.43	6 (2%) 62 56	45, 63, 96, 122	0
1	C	279/311 (89%)	0.49	16 (5%) 23 18	54, 72, 105, 127	0
1	D	270/311 (86%)	0.77	35 (12%) 3 2	44, 76, 111, 136	0
All	All	1115/1244 (89%)	0.56	63 (5%) 23 18	41, 66, 106, 136	0

The worst 5 of 63 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	5	TYR	4.8
1	D	103	ALA	4.5
1	C	37	GLU	4.4
1	D	10	ILE	4.4
1	D	32	ILE	4.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.