

Full wwPDB X-ray Structure Validation Report (i)

Jun 14, 2022 – 04:09 pm BST

PDB ID	:	7ZXF
Title	:	Pfs48/45 bound to monoclonal antibodies 10D8 and $85RF45.1$
Authors	:	Ko, K.T.; Lennartz, F.L.; Higgins, M.K.
Deposited on	:	2022-05-20
Resolution	:	3.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.28.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.28.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 3.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	1089 (3.90-3.54)
Clashscore	141614	1012 (3.88-3.56)
Ramachandran outliers	138981	1114 (3.90-3.54)
Sidechain outliers	138945	1110 (3.90-3.54)
RSRZ outliers	127900	1020 (3.90-3.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain							
1	А	448	4%	59%	16%	25%				
2	В	445	2% 	9%	54%					
3	С	216	3%	81%		19%				
4	D	466	2% 35%	11%	54%					
5	Е	240	4%	75%	1	• 10%				



Mol	Chain	Length	Qua	lity of chain
6	F	4	25%	75%
6	G	4	50%	50%



7ZXF

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 9384 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Gametocyte surface protein P45/48.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	337	Total 2738	C 1765	N 432	0 524	S 17	0	0	0

• Molecule 2 is a protein called 85RF45.1 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	В	206	Total 1573	C 1002	N 259	0 304	S 8	85	0	0

• Molecule 3 is a protein called 85RF45.1 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	С	216	Total 1657	C 1025	N 284	O 339	S 9	0	0	0

• Molecule 4 is a protein called 10D8 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	214	Total 1626	C 1023	N 273	O 320	S 10	34	0	0

• Molecule 5 is a protein called 10D8 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	Е	217	Total 1690	C 1061	N 278	0 344	${f S}7$	50	0	0

• Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluco pyranose.





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	F	4	Total 50	C 28	N 2	O 20	0	0	0
6	G	4	Total 50	C 28	N 2	O 20	0	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Gametocyte surface protein P45/48





 $\bullet \ Molecule \ 6: \ alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose \\ eta-D-glucopyranose \ (1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose \ (1-4)-2-acetamido-2-deoxy-beta-D-glucopyra$

Chain F: 25%

75%



<mark>NAG1</mark> NAG2 BMA3 MAN4

 $\bullet \ Molecule \ 6: \ alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose$

Chain G: 50% 50%

NAG1 NAG2 BMA3 MAN4



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants	156.88Å 156.88Å 148.76Å	Deperitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	70.16 - 3.72	Depositor
Resolution (A)	70.16 - 3.72	EDS
% Data completeness	84.6 (70.16-3.72)	Depositor
(in resolution range)	84.6 (70.16-3.72)	EDS
R_{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.27 (at 3.77 Å)	Xtriage
Refinement program	BUSTER 1.19_4092, PHENIX 1.19_4092	Depositor
P. P.	0.281 , 0.298	Depositor
Λ, Λ_{free}	0.289 , 0.316	DCC
R_{free} test set	790 reflections (4.66%)	wwPDB-VP
Wilson B-factor $(Å^2)$	189.9	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for $twinning^2$	$ L > = 0.45, < L^2 > = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9384	wwPDB-VP
Average B, all atoms $(Å^2)$	216.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
IVIOI	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.25	0/2798	0.47	0/3783
2	В	0.24	0/1614	0.48	0/2206
3	С	0.24	0/1696	0.48	0/2311
4	D	0.24	0/1667	0.49	0/2275
5	Е	0.25	0/1731	0.47	0/2350
All	All	0.25	0/9506	0.48	0/12925

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2738	0	2672	44	0
2	В	1573	0	1535	29	0
3	С	1657	0	1578	25	0
4	D	1626	0	1586	34	0
5	Е	1690	0	1623	24	0
6	F	50	0	43	0	0
6	G	50	0	43	0	0
All	All	9384	0	9080	149	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (149) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom_1	Atom_2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
3:C:36:TRP:HB2	3:C:49:ILE:HB	1.69	0.74	
1:A:229:LEU:HB2	1:A:232:LYS:HD2	1.70	0.73	
4:D:37:VAL:HG12	4:D:47:TRP:HA	1.72	0.71	
3:C:155:ASN:HD22	3:C:191:HIS:HB3	1.56	0.70	
4:D:148:LEU:HD13	4:D:203:VAL:HG21	1.73	0.69	
5:E:153:LYS:HB2	5:E:201:GLU:HG3	1.78	0.66	
1:A:123:LEU:HD13	1:A:154:ILE:HD11	1.76	0.66	
4:D:23:ALA:HA	4:D:78:THR:HG22	1.79	0.65	
1:A:102:GLU:HB2	1:A:107:LYS:HG3	1.79	0.64	
1:A:295:ILE:HG12	1:A:339:LEU:HD23	1.80	0.64	
2:B:22:CYS:HB3	2:B:79:LEU:HB3	1.79	0.64	
3:C:132:ASN:HA	3:C:185:SER:HB3	1.79	0.64	
1:A:85:ARG:HA	1:A:143:THR:HG22	1.81	0.63	
3:C:6:GLN:NE2	3:C:91:CYS:SG	2.62	0.62	
1:A:182:TYR:HE2	1:A:222:LEU:HB2	1.65	0.62	
2:B:125:PRO:HB3	2:B:151:TYR:HB3	1.80	0.62	
1:A:109:TYR:HE1	1:A:115:ARG:HA	1.65	0.61	
4:D:148:LEU:HB2	4:D:191:VAL:HG13	1.81	0.61	
2:B:128:TYR:HB2	2:B:147:LEU:HB3	1.83	0.61	
2:B:71:SER:HB3	2:B:80:TYR:HB2	1.82	0.61	
3:C:19:LYS:HB3	3:C:75:LEU:HD22	1.84	0.60	
5:E:44:GLN:HG3	5:E:50:PRO:HG3	1.85	0.58	
4:D:91:THR:HG23	4:D:120:THR:HA	1.86	0.58	
4:D:33:ALA:HA	4:D:72:ARG:HH12	1.69	0.58	
1:A:253:LYS:HD2	1:A:255:LEU:HD13	1.86	0.58	
1:A:184:HIS:HB3	1:A:221:GLU:HA	1.86	0.57	
1:A:217:LEU:HD11	1:A:286:LEU:HD23	1.86	0.57	
1:A:320:ASP:HB2	1:A:323:ALA:HB2	1.86	0.57	
2:B:40:ALA:HB3	2:B:43:LYS:HD2	1.88	0.55	
4:D:173:VAL:HG22	4:D:191:VAL:HG23	1.87	0.55	
4:D:129:PRO:HB3	4:D:155:TYR:HB3	1.88	0.55	
3:C:134:ALA:HB3	3:C:183:LEU:HB2	1.89	0.55	
1:A:126:ILE:HG13	1:A:151:PRO:HD3	1.88	0.55	
2:B:205:HIS:HB3	2:B:210:THR:HG22	1.87	0.55	
2:B:194:TRP:CG	2:B:195:PRO:HA	2.41	0.55	
1:A:250:ILE:HG23	1:A:257:ILE:HB	1.88	0.55	
5:E:199:THR:HG23	5:E:214:SER:HB2	1.89	0.54	



	lo uo pugo	Interatomic	Clash		
Atom-1	Atom-2	distance (Å)	overlap (Å)		
4:D:83:MET:HB3	4:D:86:LEU:HD21	1.90	0.54		
1:A:154:ILE:HG23	1:A:179:VAL:HB	1.89	0.53		
1:A:376:ILE:HD13	1:A:406:THR:HG21	1.89	0.53		
1:A:415:ASP:OD2	3:C:32:ASN:ND2	2.39	0.53		
4:D:34:MET:HB3	4:D:79:LEU:HD22	1.91	0.53		
5:E:45:LYS:HD3	5:E:90:ALA:HB2	1.91	0.53		
1:A:221:GLU:H	1:A:260:ALA:HB3	1.73	0.53		
3:C:120:THR:HB	3:C:139:LEU:HB2	1.91	0.53		
4:D:204:THR:HA	4:D:219:LYS:HA	1.91	0.52		
2:B:51:ILE:HB	2:B:70:ILE:HD13	1.92	0.52		
2:B:194:TRP:CD1	2:B:195:PRO:HA	2.44	0.52		
2:B:173:PRO:HD3	3:C:176:MET:HE3	1.90	0.52		
5:E:167:ASN:O	5:E:167:ASN:ND2	2.38	0.52		
1:A:125:LEU:HD23	1:A:128:TYR:HB3	1.92	0.51		
2:B:198:THR:HG23	2:B:215:LYS:HE3	1.92	0.51		
2:B:199:VAL:HG12	2:B:216:ILE:HD13	1.93	0.51		
1:A:341:GLY:HA3	1:A:395:LEU:HB3	1.92	0.50		
4:D:11:LEU:HD11	4:D:122:SER:HB2	1.94	0.50		
4:D:12:VAL:HG21	4:D:86:LEU:HD13	1.94	0.50		
5:E:89:LEU:HD13	5:E:172:GLN:HB2	1.94	0.50		
5:E:126:PRO:HD3	5:E:138:VAL:HG22	1.94	0.50		
2:B:5:VAL:HB	2:B:23:VAL:HB	1.95	0.49		
3:C:30:GLY:H	3:C:69:ARG:HH12	1.61	0.48		
5:E:72:GLY:HA3	5:E:77:PHE:HA	1.94	0.48		
1:A:287:LYS:HD3	4:D:57:TYR:CE1	2.49	0.48		
2:B:3:GLN:HB3	2:B:25:SER:HB2	1.96	0.48		
2:B:12:LEU:HD11	2:B:18:LEU:HB2	1.95	0.48		
1:A:235:PHE:CD1	1:A:273:CYS:HB2	2.49	0.48		
5:E:144:ASN:ND2	5:E:176:ASP:OD2	2.47	0.48		
4:D:5:VAL:HG23	4:D:23:ALA:HB3	1.96	0.48		
4:D:178:ALA:HB2	4:D:187:LEU:HD23	1.96	0.48		
1:A:293:LYS:HD2	1:A:307:LYS:O	2.13	0.47		
4:D:6:GLU:OE2	4:D:114:GLY:HA3	2.13	0.47		
4:D:177:PRO:HD2	5:E:168:SER:HB3	1.95	0.47		
5:E:70:GLY:HA2	5:E:79:LEU:HA	1.95	0.47		
5:E:31:TYR:HB2	5:E:34:ASN:HB3	1.96	0.47		
2:B:99:ASP:HB3	2:B:106:PHE:HD1	1.80	0.47		
1:A:269:VAL:O	1:A:287:LYS:HG2	2.15	0.47		
1:A:339:LEU:HD11	1:A:395:LEU:HD13	1.95	0.47		
1:A:105:PHE:O	1:A:130:ILE:HG21	2.15	0.46		
5:E:139:VAL:HA	5:E:183:SER:O	2.15	0.46		



		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:109:TYR:CE1	1:A:115:ARG:HA	2.47	0.46	
2:B:19:LYS:HE2	2:B:80:TYR:HB3	1.96	0.46	
1:A:108:VAL:HG12	1:A:110:THR:HG23	1.98	0.46	
3:C:151:ALA:HB3	3:C:197:GLN:HG3	1.98	0.46	
2:B:104:ASP:N	2:B:104:ASP:OD1	2.48	0.46	
5:E:201:GLU:HB2	5:E:210:PRO:HB2	1.97	0.46	
1:A:132:GLU:HB2	1:A:142:ARG:HE	1.81	0.45	
3:C:153:LYS:HB2	3:C:195:THR:OG1	2.16	0.45	
4:D:6:GLU:OE2	4:D:96:CYS:HB3	2.16	0.45	
1:A:246:SER:HB2	1:A:259:LYS:O	2.17	0.45	
1:A:416:LYS:HE3	3:C:33:TYR:CD1	2.52	0.45	
5:E:148:LYS:HE2	5:E:169:TRP:HB3	1.98	0.45	
3:C:36:TRP:HA	3:C:90:PHE:O	2.17	0.45	
5:E:43:GLN:HG3	5:E:90:ALA:HB3	1.99	0.45	
1:A:293:LYS:O	1:A:309:THR:HB	2.17	0.45	
2:B:99:ASP:N	2:B:99:ASP:OD1	2.48	0.45	
4:D:47:TRP:HZ2	4:D:50:THR:HG22	1.82	0.45	
4:D:52:SER:O	4:D:72:ARG:NH1	2.49	0.45	
4:D:28:THR:HB	4:D:31:ASN:HB2	1.99	0.44	
4:D:35:SER:HA	4:D:50:THR:HA	1.99	0.44	
1:A:299:ASN:HB2	1:A:310:PHE:CD2	2.51	0.44	
3:C:15:GLY:HA2	3:C:80:ASN:HA	2.00	0.44	
4:D:107:LEU:HG	5:E:97:TYR:HB2	1.98	0.44	
5:E:167:ASN:HB2	5:E:169:TRP:CZ3	2.52	0.44	
1:A:79:PHE:HD2	1:A:179:VAL:HG13	1.82	0.44	
2:B:99:ASP:HB3	2:B:106:PHE:CD1	2.53	0.44	
2:B:104:ASP:HB3	3:C:94:TYR:HB3	1.98	0.44	
3:C:136:LEU:HD23	3:C:136:LEU:HA	1.86	0.44	
5:E:34:ASN:C	5:E:34:ASN:HD22	2.20	0.44	
1:A:157:PHE:CE1	1:A:176:HIS:HB2	2.53	0.43	
2:B:16:ARG:HA	2:B:16:ARG:NH1	2.32	0.43	
5:E:156:ILE:HD11	5:E:185:LEU:HD21	1.99	0.43	
2:B:151:TYR:CE1	2:B:183:LEU:HD23	2.54	0.43	
4:D:181:GLN:N	4:D:184:LEU:O	2.26	0.43	
1:A:386:ASP:HB3	1:A:393:ILE:HB	2.01	0.43	
3:C:24:ARG:NE	3:C:26:THR:O	2.51	0.43	
4:D:20:VAL:HG23	4:D:81:LEU:HB3	2.00	0.42	
4:D:126:THR:HG22	4:D:157:PRO:HD3	2.00	0.42	
3:C:135:THR:HA	3:C:181:LEU:O	2.19	0.42	
4:D:13:LYS:HD2	4:D:14:PRO:HD2	2.01	0.42	
5:E:201:GLU:HB3	5:E:212:VAL:HG23	2.02	0.42	



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance $(Å)$	overlap (Å)
4:D:134:LEU:HD11	4:D:151:LEU:HB2	2.02	0.42
1:A:182:TYR:HD1	1:A:183:PRO:HD2	1.83	0.42
3:C:62:ARG:HB3	3:C:79:ASP:O	2.20	0.42
2:B:9:GLY:H	2:B:18:LEU:HD21	1.86	0.41
5:E:8:PRO:O	5:E:108:THR:HG22	2.21	0.41
5:E:144:ASN:ND2	5:E:178:THR:OG1	2.41	0.41
1:A:102:GLU:OE2	1:A:117:GLU:HG2	2.20	0.41
2:B:154:GLU:HG2	2:B:155:PRO:HB3	2.02	0.41
3:C:37:TYR:HE2	3:C:92:HIS:HB3	1.85	0.41
1:A:46:LYS:HG3	1:A:85:ARG:HB2	2.03	0.41
1:A:336:TYR:O	1:A:399:VAL:HA	2.20	0.41
1:A:343:ASN:OD1	1:A:391:ASP:HB3	2.21	0.41
4:D:122:SER:HA	4:D:123:PRO:HD3	1.92	0.41
3:C:30:GLY:HA3	3:C:69:ARG:HH22	1.86	0.41
2:B:34:MET:HB2	2:B:51:ILE:HG22	2.03	0.41
1:A:332:SER:HB2	1:A:426:ASP:OD2	2.20	0.41
3:C:117:PRO:HD3	3:C:200:HIS:CD2	2.55	0.41
1:A:213:LEU:HD23	1:A:284:ILE:HG23	2.03	0.41
2:B:196:SER:OG	2:B:197:GLN:N	2.54	0.41
3:C:195:THR:HG22	3:C:208:THR:HB	2.02	0.41
4:D:13:LYS:HD3	4:D:123:PRO:HA	2.02	0.41
4:D:151:LEU:HD21	4:D:153:LYS:HB2	2.02	0.41
1:A:182:TYR:CD1	1:A:183:PRO:HD2	2.56	0.40
2:B:23:VAL:HG22	2:B:78:THR:HG22	2.02	0.40
3:C:86:GLU:HG3	3:C:108:THR:HA	2.03	0.40
5:E:21:MET:HB2	5:E:79:LEU:HB3	2.03	0.40
4:D:178:ALA:HA	4:D:187:LEU:HB3	2.02	0.40
4:D:216:VAL:HG12	4:D:218:LYS:HG3	2.03	0.40
1:A:293:LYS:HB2	1:A:293:LYS:HE2	1.73	0.40
2:B:149:LYS:HA	2:B:182:THR:HG23	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	327/448~(73%)	292~(89%)	35 (11%)	0	100 100
2	В	202/445~(45%)	186~(92%)	16 (8%)	0	100 100
3	С	214/216~(99%)	205~(96%)	9 (4%)	0	100 100
4	D	210/466~(45%)	195~(93%)	15 (7%)	0	100 100
5	Е	215/240~(90%)	203 (94%)	12 (6%)	0	100 100
All	All	1168/1815~(64%)	1081 (93%)	87 (7%)	0	100 100

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	319/419~(76%)	314~(98%)	5(2%)	62 79
2	В	177/400~(44%)	174 (98%)	3~(2%)	60 78
3	С	193/193~(100%)	193 (100%)	0	100 100
4	D	186/417~(45%)	186 (100%)	0	100 100
5	Ε	194/214~(91%)	190~(98%)	4 (2%)	53 74
All	All	1069/1643~(65%)	1057~(99%)	12 (1%)	73 85

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	110	THR
1	А	190	ASN
1	А	234	CYS
1	А	244	TYR
1	А	354	PHE
2	В	99	ASP
2	В	101	ARG



Continued from previous page...

Mol	Chain	Res	Type
2	В	104	ASP
5	Е	34	ASN
5	Е	44	GLN
5	Е	167	ASN
5	Е	173	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	С	82	GLN
3	С	155	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	ol Type Chain Reg		ain Deg Link		Bo	Bond lengths		Bond angles		
Moi Type Chain	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
6	NAG	F	1	1,6	$14,\!14,\!15$	0.60	0	17,19,21	0.87	0
6	NAG	F	2	6	14,14,15	1.16	1(7%)	17,19,21	1.32	2 (11%)
6	BMA	F	3	6	11,11,12	1.18	1 (9%)	15,15,17	1.45	2 (13%)
6	MAN	F	4	6	11,11,12	0.77	0	15,15,17	1.47	2 (13%)
6	NAG	G	1	1,6	14,14,15	1.61	1 (7%)	17,19,21	1.14	2 (11%)
6	NAG	G	2	6	14,14,15	0.24	0	17,19,21	0.67	0



Mol Type	Turne	vno Chain B		Tinle	Bond lengths			Bond angles		
	туре	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
6	BMA	G	3	6	11,11,12	0.78	0	$15,\!15,\!17$	0.88	0
6	MAN	G	4	6	11,11,12	0.62	0	$15,\!15,\!17$	0.95	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	F	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	F	2	6	-	3/6/23/26	0/1/1/1
6	BMA	F	3	6	-	1/2/19/22	0/1/1/1
6	MAN	F	4	6	-	0/2/19/22	1/1/1/1
6	NAG	G	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	G	2	6	-	3/6/23/26	0/1/1/1
6	BMA	G	3	6	-	2/2/19/22	0/1/1/1
6	MAN	G	4	6	-	2/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
6	G	1	NAG	O5-C1	-5.83	1.34	1.43
6	F	2	NAG	O5-C1	-4.10	1.37	1.43
6	F	3	BMA	C1-C2	2.60	1.58	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
6	F	4	MAN	C1-O5-C5	4.54	118.35	112.19
6	F	2	NAG	C4-C3-C2	3.43	116.04	111.02
6	F	3	BMA	C1-C2-C3	-3.22	105.71	109.67
6	F	3	BMA	O5-C1-C2	-2.85	106.37	110.77
6	F	2	NAG	C3-C4-C5	2.48	114.65	110.24
6	G	4	MAN	C1-O5-C5	2.37	115.41	112.19
6	G	1	NAG	C1-O5-C5	-2.36	108.99	112.19
6	G	1	NAG	C3-C4-C5	2.28	114.30	110.24
6	F	4	MAN	O2-C2-C3	-2.22	105.69	110.14
6	G	4	MAN	O2-C2-C3	-2.17	105.78	110.14

There are no chirality outliers.



7	7VF
1	$\Delta \Lambda \Gamma$

Mol	Chain	Res	Type	Atoms
6	G	3	BMA	O5-C5-C6-O6
6	G	3	BMA	C4-C5-C6-O6
6	G	2	NAG	O5-C5-C6-O6
6	G	2	NAG	C4-C5-C6-O6
6	G	4	MAN	C4-C5-C6-O6
6	G	4	MAN	O5-C5-C6-O6
6	F	2	NAG	C1-C2-N2-C7
6	F	3	BMA	O5-C5-C6-O6
6	F	1	NAG	C3-C2-N2-C7
6	F	2	NAG	C3-C2-N2-C7
6	G	1	NAG	C3-C2-N2-C7
6	G	2	NAG	C3-C2-N2-C7
6	F	2	NAG	O5-C5-C6-O6
6	F	1	NAG	C1-C2-N2-C7
6	G	1	NAG	C1-C2-N2-C7

All (15) torsion outliers are listed below:

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	F	4	MAN	C1-C2-C3-C4-C5-O5

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry (i)

There are no ligands in this entry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	337/448~(75%)	0.28	19 (5%) 24 20	158, 209, 264, 280	0
2	В	194/445~(43%)	0.02	8 (4%) 37 31	146, 201, 246, 256	1 (0%)
3	С	216/216~(100%)	0.07	7 (3%) 47 39	177, 223, 253, 260	0
4	D	209/466~(44%)	0.15	11 (5%) 26 23	156, 209, 258, 295	0
5	Е	211/240~(87%)	-0.02	10 (4%) 31 26	180, 231, 261, 282	0
All	All	1167/1815~(64%)	0.12	55 (4%) 31 26	146, 217, 257, 295	1 (0%)

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	Е	43	GLN	6.4
2	В	184	THR	5.5
5	Е	110	LEU	4.4
5	Е	88	ASP	4.1
3	С	121	MET	4.0
5	Е	148	LYS	3.8
4	D	129	PRO	3.7
4	D	220	ILE	3.6
2	В	185	SER	3.6
1	А	399	VAL	3.4
1	А	338	HIS	3.2
1	А	250	ILE	3.1
1	А	398	ILE	3.0
3	С	120	THR	3.0
1	А	251	TYR	3.0
2	В	149	LYS	2.9
4	D	131	VAL	2.9
2	В	127	VAL	2.9
4	D	133	PRO	2.8
3	С	137	VAL	2.7



Mol	Chain	Res	Type	RSRZ
2	В	125	PRO	2.6
3	С	209	VAL	2.6
5	Е	92	TYR	2.6
5	Е	65	PRO	2.5
1	А	397	GLY	2.5
1	А	222	LEU	2.5
2	В	177	GLN	2.5
4	D	162	VAL	2.5
1	А	156	PHE	2.5
4	D	110	MET	2.4
1	А	235	PHE	2.4
4	D	203	VAL	2.4
4	D	111	ASP	2.4
5	Е	21	MET	2.4
5	Е	42	TYR	2.3
1	А	152	LYS	2.3
5	Е	41	TRP	2.2
1	А	257	ILE	2.2
4	D	113	TRP	2.2
2	В	216	ILE	2.2
4	D	151	LEU	2.2
1	А	86	LEU	2.2
3	С	52	ASP	2.2
5	Е	13	VAL	2.2
2	В	148	VAL	2.2
1	А	249	ILE	2.2
4	D	175	THR	2.2
1	А	175	VAL	2.2
3	С	139	LEU	2.1
1	А	425	ILE	2.1
1	А	188	PHE	2.1
1	А	406	THR	2.1
1	А	339	LEU	2.1
3	С	194	PHE	2.1
1	A	126	ILE	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
6	BMA	F	3	11/12	0.69	0.19	247,252,259,262	0
6	NAG	G	2	14/15	0.82	0.15	226,261,274,277	0
6	BMA	G	3	11/12	0.83	0.13	251,258,266,270	0
6	NAG	G	1	14/15	0.84	0.12	216,245,270,277	0
6	NAG	F	1	14/15	0.86	0.19	231,240,245,249	0
6	NAG	F	2	14/15	0.88	0.18	238,249,256,256	0
6	MAN	G	4	11/12	0.91	0.18	261,265,270,271	0
6	MAN	F	4	11/12	0.95	0.15	253,257,266,269	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

