



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2023 – 12:14 PM EDT

PDB ID : 1ZZ8  
Title : Crystal Structure of FeII HppE in Complex with Substrate Form 2  
Authors : Higgins, L.J.; Yan, F.; Liu, P.; Liu, H.W.; Drennan, C.L.  
Deposited on : 2005-06-13  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	198	 6% 77% 18% . .
1	B	198	 6% 72% 22% . . .
1	C	198	 3% 73% 22% . .

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4515 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hydroxypropylphosphonic Acid Epoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	192	1445	914	246	280	5	0	0	0
1	B	193	1451	918	250	278	5	0	0	0
1	C	192	1446	914	248	279	5	0	0	0

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	B	1	Total	Fe	0	0
			1	1		
2	C	1	Total	Fe	0	0
			1	1		

- Molecule 3 is (S)-2-HYDROXYPROPYLPHOSPHONIC ACID (three-letter code: S0H) (formula: C<sub>3</sub>H<sub>9</sub>O<sub>4</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			8	3	4	1		
3	B	1	Total	C	O	P	0	0
			8	3	4	1		
3	C	1	Total	C	O	P	0	0
			8	3	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	51	Total	O	0	0
			51	51		
4	B	31	Total	O	0	0
			31	31		
4	C	64	Total	O	0	0
			64	64		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hydroxypropylphosphonic Acid Epoxidase



- Molecule 1: Hydroxypropylphosphonic Acid Epoxidase



- Molecule 1: Hydroxypropylphosphonic Acid Epoxidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.65Å 111.65Å 152.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 30.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.3 (30.00-2.30) 96.4 (30.00-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.84 (at 2.31Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.225 , 0.252 0.221 , 0.252	Depositor DCC
$R_{free}$ test set	1852 reflections (4.31%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.8	Xtrriage
Anisotropy	0.133	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 39.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4515	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 31.13 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1656e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: FE2, S0H

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/1475	0.70	1/2007 (0.0%)
1	B	0.44	0/1481	0.71	2/2015 (0.1%)
1	C	0.44	0/1475	0.72	2/2006 (0.1%)
All	All	0.44	0/4431	0.71	5/6028 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	194	ILE	N-CA-C	-5.78	95.40	111.00
1	B	144	LEU	CA-CB-CG	5.26	127.40	115.30
1	C	121	VAL	N-CA-C	-5.18	97.02	111.00
1	B	135	ASN	N-CA-C	-5.01	97.48	111.00
1	A	135	ASN	N-CA-C	-5.00	97.49	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	83	ILE	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1445	0	1420	27	0
1	B	1451	0	1426	41	0
1	C	1446	0	1422	31	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	8	0	3	0	0
3	B	8	0	3	0	0
3	C	8	0	3	0	0
4	A	51	0	0	0	1
4	B	31	0	0	5	0
4	C	64	0	0	0	0
All	All	4515	0	4277	98	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:153:MET:HE2	1:B:166:LEU:HD11	1.47	0.95
1:A:91:PRO:HG2	1:A:107:CYS:HB2	1.50	0.92
1:B:158:LYS:HD2	1:B:158:LYS:H	1.36	0.90
1:B:145:PHE:HB3	1:B:194:ILE:HG22	1.58	0.85
1:A:145:PHE:HB3	1:A:194:ILE:HG22	1.62	0.80
1:C:186:LYS:HE3	1:C:187:GLY:N	1.97	0.78
1:B:145:PHE:HB3	1:B:194:ILE:CG2	2.15	0.76
1:B:91:PRO:HG2	1:B:107:CYS:HB2	1.66	0.76
1:C:146:VAL:HG13	1:C:151:ILE:HD13	1.66	0.76
1:B:148:GLU:OE1	1:B:192:LYS:HD3	1.87	0.74
1:C:154:LYS:HG2	1:C:163:GLU:HG2	1.68	0.74
1:B:126:THR:HG22	4:B:1007:HOH:O	1.92	0.70
1:A:145:PHE:HB3	1:A:194:ILE:CG2	2.24	0.68
1:B:102:TYR:CE1	1:B:126:THR:HG21	2.30	0.67
1:C:156:GLY:O	1:C:179:PRO:HG2	1.95	0.66

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:158:LYS:HD2	1:B:158:LYS:N	2.10	0.66
1:A:122:VAL:HB	1:A:193:LEU:CD1	2.26	0.66
1:C:144:LEU:HB2	1:C:172:MET:HG2	1.78	0.65
1:B:134:PHE:HA	1:B:181:ALA:HB2	1.81	0.63
1:B:153:MET:CE	1:B:166:LEU:HD11	2.27	0.62
1:B:32:LEU:HD22	1:B:62:VAL:HG13	1.83	0.61
1:C:104:VAL:HB	1:C:123:ASP:HB2	1.83	0.61
1:B:148:GLU:HG3	1:B:192:LYS:HB3	1.83	0.60
1:A:122:VAL:HB	1:A:193:LEU:HD11	1.83	0.59
1:C:91:PRO:HG2	1:C:107:CYS:HB2	1.84	0.59
1:A:144:LEU:HB2	1:A:172:MET:HG2	1.85	0.58
1:A:193:LEU:HD12	1:A:193:LEU:O	2.03	0.58
1:C:186:LYS:HE3	1:C:187:GLY:H	1.69	0.57
1:B:126:THR:CG2	4:B:1007:HOH:O	2.51	0.57
1:B:156:GLY:O	1:B:179:PRO:HG2	2.05	0.56
1:A:90:ARG:HB2	1:A:107:CYS:O	2.05	0.55
1:B:104:VAL:HG23	1:B:125:LEU:HD11	1.89	0.55
1:A:100:VAL:HG11	1:A:133:LYS:NZ	2.23	0.54
1:C:157:ASP:OD2	1:C:159:GLU:HB2	2.08	0.54
1:C:97:ARG:HG2	1:C:133:LYS:CB	2.38	0.53
1:C:130:ASP:C	1:C:132:ALA:N	2.62	0.52
1:C:97:ARG:HB2	1:C:102:TYR:HB2	1.91	0.52
1:B:23:LYS:HD2	1:B:23:LYS:N	2.23	0.52
1:C:150:GLU:HG2	1:C:185:ALA:HB2	1.91	0.52
1:C:58:ARG:HG2	1:C:58:ARG:HH11	1.74	0.52
1:C:151:ILE:HD11	1:C:191:ALA:HB1	1.92	0.52
1:C:107:CYS:HA	1:C:120:LEU:HD23	1.92	0.51
1:A:112:LYS:NZ	1:B:61:HIS:ND1	2.50	0.51
1:A:52:THR:OG1	1:A:55:GLN:HG3	2.11	0.51
1:B:142:GLU:HG2	1:B:144:LEU:HD22	1.93	0.51
1:A:117:LEU:HG	1:A:119:PRO:HD3	1.92	0.51
1:C:126:THR:HG22	1:C:128:ASN:H	1.74	0.51
1:C:146:VAL:HG13	1:C:151:ILE:CD1	2.39	0.51
1:A:100:VAL:HG11	1:A:133:LYS:HZ2	1.76	0.50
1:B:141:ASN:ND2	1:B:176:GLU:H	2.09	0.50
1:A:134:PHE:HA	1:A:181:ALA:HB2	1.92	0.50
1:B:150:GLU:HB3	1:B:185:ALA:HB2	1.94	0.50
1:C:117:LEU:HG	1:C:119:PRO:HD3	1.94	0.49
1:B:18:ARG:HA	1:B:21:GLN:HE21	1.77	0.49
1:B:35:GLU:OE2	1:B:39:THR:HG21	2.12	0.49
1:B:39:THR:O	1:B:42:ALA:HB3	2.12	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:134:PHE:HA	1:C:181:ALA:HB2	1.95	0.49
1:A:156:GLY:O	1:A:179:PRO:HG2	2.12	0.49
1:B:193:LEU:C	1:B:193:LEU:HD12	2.33	0.49
1:B:142:GLU:HG2	1:B:144:LEU:CD2	2.44	0.48
1:C:128:ASN:O	1:C:130:ASP:N	2.47	0.48
1:A:122:VAL:HB	1:A:193:LEU:HD12	1.95	0.47
1:B:98:ASP:O	1:B:100:VAL:HG23	2.14	0.47
1:B:85:GLN:OE1	1:B:108:LEU:HA	2.15	0.47
1:C:127:ASP:O	1:C:186:LYS:HA	2.14	0.46
1:A:137:GLY:HA3	1:A:177:HIS:O	2.14	0.46
1:B:158:LYS:HE2	1:B:179:PRO:CB	2.45	0.46
1:A:16:LYS:HB2	1:A:44:GLU:HG2	1.98	0.46
1:A:19:ARG:HG2	1:A:24:MET:O	2.16	0.46
1:B:97:ARG:NH2	4:B:1006:HOH:O	2.48	0.46
1:B:104:VAL:HB	1:B:123:ASP:HB2	1.97	0.46
1:A:84:ILE:CG2	1:A:85:GLN:N	2.79	0.45
1:B:128:ASN:C	1:B:128:ASN:HD22	2.18	0.45
1:B:141:ASN:HD22	1:B:176:GLU:H	1.62	0.45
1:C:97:ARG:HB2	1:C:102:TYR:CB	2.47	0.45
1:A:193:LEU:HD12	1:A:193:LEU:C	2.37	0.44
1:B:28:ALA:O	1:B:32:LEU:HB2	2.18	0.44
1:A:104:VAL:HG23	1:A:125:LEU:HD11	1.99	0.44
1:A:84:ILE:HG22	1:A:85:GLN:N	2.31	0.44
1:C:86:MET:O	1:C:89:GLU:HG2	2.17	0.44
1:B:117:LEU:HG	1:B:119:PRO:HD3	1.98	0.44
1:C:104:VAL:HG23	1:C:125:LEU:HD11	2.00	0.44
1:B:151:ILE:HD13	1:B:193:LEU:HD23	1.99	0.44
1:C:176:GLU:O	1:C:177:HIS:HB2	2.18	0.43
1:C:130:ASP:N	1:C:130:ASP:OD2	2.51	0.43
1:B:18:ARG:HB2	4:B:1026:HOH:O	2.17	0.43
1:B:158:LYS:H	1:B:158:LYS:CD	2.18	0.43
1:A:33:LEU:HD23	1:A:62:VAL:HG21	2.02	0.42
1:C:151:ILE:CD1	1:C:191:ALA:HB1	2.50	0.42
1:A:100:VAL:CG1	1:A:133:LYS:HZ3	2.33	0.41
1:C:36:THR:OG1	1:C:38:GLU:HG2	2.21	0.41
1:B:142:GLU:HB3	1:B:174:VAL:HG13	2.01	0.41
1:A:97:ARG:O	1:A:100:VAL:HG13	2.19	0.41
1:B:158:LYS:HE2	1:B:179:PRO:HB2	2.03	0.41
1:C:118:VAL:HB	1:C:197:ASN:HB2	2.01	0.41
1:C:127:ASP:HB3	1:C:189:GLY:O	2.21	0.41
1:A:63:LEU:HD12	1:A:63:LEU:HA	1.94	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:124:VAL:HG13	4:B:1007:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1047:HOH:O	4:A:1052:HOH:O[2_765]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/198 (96%)	178 (94%)	12 (6%)	0	100	100
1	B	191/198 (96%)	187 (98%)	4 (2%)	0	100	100
1	C	188/198 (95%)	181 (96%)	6 (3%)	1 (0%)	29	35
All	All	569/594 (96%)	546 (96%)	22 (4%)	1 (0%)	47	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	129	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	152/160 (95%)	145 (95%)	7 (5%)	27	38
1	B	151/160 (94%)	140 (93%)	11 (7%)	14	18
1	C	152/160 (95%)	146 (96%)	6 (4%)	32	46
All	All	455/480 (95%)	431 (95%)	24 (5%)	22	31

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	LEU
1	A	70	LEU
1	A	100	VAL
1	A	131	ASP
1	A	148	GLU
1	A	158	LYS
1	A	172	MET
1	B	32	LEU
1	B	80	ASP
1	B	82	VAL
1	B	97	ARG
1	B	101	ASP
1	B	126	THR
1	B	128	ASN
1	B	144	LEU
1	B	148	GLU
1	B	158	LYS
1	B	159	GLU
1	C	83	ILE
1	C	97	ARG
1	C	129	PRO
1	C	147	LEU
1	C	172	MET
1	C	186	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	21	GLN
1	B	45	ASN
1	B	99	ASN
1	B	128	ASN
1	B	141	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	106	ASN
1	C	160	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	S0H	A	1001	2	7,7,7	4.19	4 (57%)	8,10,10	2.30	3 (37%)
3	S0H	B	1003	2	7,7,7	4.21	4 (57%)	8,10,10	2.32	2 (25%)
3	S0H	C	1002	2	7,7,7	4.40	4 (57%)	8,10,10	2.36	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	S0H	A	1001	2	-	0/5/5/5	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	S0H	B	1003	2	-	0/5/5/5	-
3	S0H	C	1002	2	-	0/5/5/5	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1002	S0H	C1-C3	-9.34	1.10	1.51
3	A	1001	S0H	C1-C3	-9.14	1.11	1.51
3	B	1003	S0H	C1-C3	-9.11	1.11	1.51
3	C	1002	S0H	P1-C2	4.94	1.83	1.78
3	A	1001	S0H	P1-O13	-4.33	1.45	1.54
3	B	1003	S0H	P1-O13	-4.18	1.45	1.54
3	C	1002	S0H	P1-O13	-4.16	1.45	1.54
3	B	1003	S0H	P1-C2	4.01	1.83	1.78
3	A	1001	S0H	P1-C2	3.49	1.82	1.78
3	A	1001	S0H	P1-O14	2.75	1.56	1.50
3	B	1003	S0H	P1-O14	2.70	1.55	1.50
3	C	1002	S0H	P1-O14	2.51	1.55	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1003	S0H	O15-P1-O14	-5.22	98.59	112.39
3	C	1002	S0H	O15-P1-O14	-5.20	98.65	112.39
3	A	1001	S0H	O15-P1-O14	-4.88	99.49	112.39
3	C	1002	S0H	O13-P1-C2	3.03	114.15	106.74
3	A	1001	S0H	O13-P1-C2	2.74	113.45	106.74
3	B	1003	S0H	O13-P1-C2	2.50	112.87	106.74
3	A	1001	S0H	O15-P1-O13	2.05	114.08	108.08

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	192/198 (96%)	0.17	11 (5%) 23 30	26, 36, 52, 63	0
1	B	193/198 (97%)	0.27	11 (5%) 23 30	22, 37, 53, 57	0
1	C	192/198 (96%)	-0.03	5 (2%) 56 63	17, 31, 47, 58	0
All	All	577/594 (97%)	0.14	27 (4%) 31 38	17, 35, 52, 63	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	160	ASN	4.2
1	C	160	ASN	4.2
1	C	159	GLU	4.0
1	A	99	ASN	3.8
1	A	98	ASP	3.5
1	B	37	PRO	3.3
1	C	98	ASP	3.2
1	B	32	LEU	3.1
1	C	157	ASP	3.0
1	B	99	ASN	2.9
1	B	196	VAL	2.9
1	A	195	ALA	2.7
1	B	194	ILE	2.5
1	B	34	GLY	2.5
1	C	158	LYS	2.4
1	A	100	VAL	2.4
1	A	128	ASN	2.3
1	B	45	ASN	2.3
1	A	196	VAL	2.2
1	B	148	GLU	2.2
1	A	194	ILE	2.2
1	A	157	ASP	2.1
1	A	80	ASP	2.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	58	ARG	2.1
1	A	159	GLU	2.1
1	B	98	ASP	2.1
1	B	195	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q < 0.9
3	S0H	B	1003	8/8	0.97	0.16	25,31,32,32	0
3	S0H	A	1001	8/8	0.99	0.13	32,33,35,38	0
2	FE2	A	199	1/1	0.99	0.13	31,31,31,31	0
3	S0H	C	1002	8/8	0.99	0.17	29,33,34,35	0
2	FE2	C	199	1/1	1.00	0.14	29,29,29,29	0
2	FE2	B	199	1/1	1.00	0.14	29,29,29,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.